

The Cryosphere Discuss., 4, C1369–C1370, 2010  
www.the-cryosphere-discuss.net/4/C1369/2010/  
© Author(s) 2010. This work is distributed under  
the Creative Commons Attribute 3.0 License.



TCD

4, C1369–C1370, 2010

Interactive  
Comment

## ***Interactive comment on “Isotope hydrological studies on the perennial ice deposit of Saarahalle, Mammuthöhle, Dachstein Mts, Austria” by Z. Kern et al.***

**Z. Kern et al.**

zoltan.kern@gmail.com

Received and published: 16 December 2010

Original Short comment: The authors should get in contact with Lukas Plan at the KHA who recently successfully dated organic matter in Feenpalast. Although not directly connected to Saarahalle, this information would fit to the paper.

Author’s response: Thanks to Michael Behm for this short comment. We have contacted with Lukas Plan as suggested. We got the information that he found two pieces of wood in the deeper lying ice block between Saarahalle and Feenpalast and these are submitted for dating however results have not been published yet.

On the other hand, some radiocarbon dating has been established from a wood rem-  
C1369

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



nant found at the base of Feenpalast's ice block c. ten years ago (Mais and Pavuza, 2000). The sample (GrA-14417) gave  $695 \pm 35$  14C BP yr age. When this date was recalibrated on the most recent calibration dataset (intcal09, Reimer et al. 2009) the calibration provided two separate intervals for the two-sigma range (1259-1352 cal AD, 68.3%, and 1318-1390 cal AD 27.1%).

We will mention these information in the expanded site description section as earlier dating result from ice deposits of the Mammuthöhle-system. However, we remark that this date might have little relevance to the age of Saarahalle ice block and we expect even greater discrepancies between the aforesaid two on-going dates and the remote and deeper lying Saarahalle-ice body. In addition we remark, that nobody could found macroscopic organic material near or at the base of the Saarahalle so far. So we might have to melt ice to get the needed 200  $\mu\text{g}$  of dispersed organic material to test the radiocarbon dating of the Saarahalle ice. However it is really a great challenge (see May et al. this issue).

References: Mais von K. and Pavuza R.: Hinweise zu Höhlenklima und Höhleis in der Dachstein Mammuthöhle (Oberösterreich), Die Höhle, 51/4, 121-125, 2000

May, B., Wagenbach, D., Spötl, C., Dublyansky Y. and Liebl, J.: First investigations of an ice core from Eisriesenwelt cave (Austria), The Cryosphere Discuss, 4, 1525-1559, doi:10.5194/tcd-4-1525-2010, 2010

Reimer PJ, Baillie MGL, Bard E, Bayliss A, Beck JW, Blackwell PG, Bronk Ramsey C, Buck CE, Burr G, Edwards RL, Friedrich M, Grootes PM, Guilderson TP, Hajdas I, Heaton TJ, Hogg AG, Hughen KA, Kaiser KF, Kromer B, McCormac FG, Manning SW, Reimer RW, Richards DA, Southon J, Turney CSM, van der Plicht J, Weyhenmeyer C: IntCal09 and Marine09 radiocarbon age calibration curves, 0 - 50,000 years cal BP. Radiocarbon 51, 1111-1150, 2009

Interactive comment on The Cryosphere Discuss., 4, 1449, 2010.