

This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

Multi-decadal marine and land-terminating glacier recession in the Ammassalik region, Southeast Greenland

S. H. Mernild¹, J. K. Malmros², N. T. Knudsen³, and J. C. Yde⁴

¹Climate, Ocean, and Sea Ice Modeling Group, Los Alamos National Laboratory, New Mexico, USA

²Department of Geography and Geology, University of Copenhagen, Denmark

³Department of Geoscience, Aarhus University, Aarhus, Denmark

⁴Sogn og Fjordane University College, Sogndal, Norway

Received: 26 January 2012 – Accepted: 28 January 2012 – Published: 7 February 2012

Correspondence to: S. H. Mernild (mernild@lanl.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Landsat imagery was applied to elucidate glacier fluctuations of land and marine terminating outlet glaciers from the Greenland Ice Sheet (GrIS) and local land terminating glaciers and ice caps (GIC) peripheral to the GrIS in the Ammassalik region, Southeast Greenland, during the period 1972–2011. Data from 21 marine-terminating glaciers – including the glaciers Helheim, Midgaard, and Fenris –, the GrIS land-terminating margin, and 35 GIC were examined, and compared to observed atmospheric air temperatures, precipitation, and reconstructed ocean water temperatures. Here, we document that net glacier recession has occurred since 1972 in the Ammassalik region for all glacier types and sizes, except for three GIC; however, the land-terminating; however, the land-terminating GrIS and GIC reflect lower marginal and areal changes than the marine-terminating outlet glaciers. The mean annual land-terminating GrIS and GIC margin recessions were about 3 to 5 times lower than the GrIS marine-terminating values. The marine-terminating outlet glaciers had an average net frontal retreat for 1999–2011 of 0.098 km yr^{-1} , which was significantly higher than in previous sub-periods 1972–1986 and 1986–1999. For the marine-terminating GrIS the annual areal recession rate was decreasing since 1972, while increasing for the land-terminating GrIS since 1986. On average for the GIC a net frontal retreat for 1986–2011 of $0.010 \pm 0.006 \text{ km yr}^{-1}$ and a mean areal recession of around 1 % per year occurred (overall $27 \pm 24 \%$). Since 1986, five GIC melted away in the Ammassalik area, and one would therefore expect that GIC might melt substantially within the 21st century under ongoing climate warming.

1 Introduction

The Greenland Ice Sheet (GrIS) – land- and marine-terminating outlet glaciers on the periphery of the GrIS – and local land-terminating glaciers and ice caps (GIC)

TCD

6, 531–561, 2012

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



peripheral to the GrIS have undergone rapid changes over the last decades. Most notable changes are thinning, leading to destabilization and accelerated retreat of GrIS marine-terminating glaciers (Joughin et al., 2004, 2010; Howat et al., 2005, 2008; Rignot and Kanagaratnam, 2006; Moon and Joughin, 2008; Nick et al., 2009), and increasing mass loss, thinning, and retreat of land-terminating GIC (Yde and Knudsen, 2007; Mernild et al., 2011a; Radić and Hock 2011).

The underlying mechanisms of GrIS marine-terminating glacier dynamics remain unclear (Straneo et al., 2010; Johannessen et al., 2011). The thinning and frontal recession are likely due to changes in oceanic warm subsurface waters and atmospheric warming, where warm subsurface waters are suggested by Luthcke et al. (2006), Velicogna and Wahr (2006), Holland et al. (2008), van den Broeke et al. (2009), Velicogna (2009), Murray et al. (2010), Rignot et al. (2010), Straneo et al. (2010), Andresen et al. (2011) to play a significant role, as well as to penetration of snow and ice melt water to the glacier bed influencing the GrIS sliding and dynamic processes (Johannessen et al., 2011). The mechanisms suggested for land-terminating GIC recession are less complex. In the Ammassalik region, Southeast Greenland, studies of GIC have shown that mass loss and margin retreat have been driven mainly by higher surface temperatures (increasing surface ablation) and decreasing precipitation (decreasing snow accumulation) since the mid-1900s (Mernild et al., 2011a).

Despite the importance of glacier contribution to sea-level rise, Greenland marine-terminating glaciers have been sparsely analyzed by satellite and direct glaciological observations. Howat and Eddy (2011) have identified changes in ice-frontal positions from 210 GrIS marine-terminating glaciers with fronts wider than 1 km (spanning nearly four decades, 1972–2010). These results show a trend of accelerated recession, where 90% of the observed glaciers receded between 2000 and 2010. Box and Decker (2011) identified areal changes at 39 of the widest Greenland marine-terminating glacier (2000–2010). Collectively, the 39 glaciers lost a cumulative area of 1368 km². For the Ammassalik region, including the Sermilik Fjord (the largest fjord system in Southeast Greenland), only frontal changes of major GrIS

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



marine-terminating glaciers, such as Helheim, Fenris, and Midgaard glaciers, have been observed; however at least 21 marine-terminating glaciers have been identified (Fig. 1).

For the land-terminating GIC in Greenland, margin retreat has been sparsely observed, and the only currently published time series of whole-glacier in-situ mass balance observations (since 1995/1996) is from the Mittivakkat Gletscher, located in the Ammassalik region (WGMS, 2009; Mernild et al., 2011a), even though thousands of individual GIC are located in the land-strip between the GrIS and ocean, of which several of hundreds are situated in the Ammassalik region (Mernild et al., 2012b). Thus, there is a need for more information about contemporary glacier fluctuations of Greenlandic GIC and their coupling to climate change.

The ability to assess GrIS and GIC margin changes in e.g., the Ammassalik region has been improved through the use of Landsat imagery going back to 1972. The imagery gives us the possibility to illustrate “snapshots” and the averaged behavior of glacier changes for the past four decades for the identified marine-terminating glaciers, the GrIS land-terminating margin, and the GIC during a period of climate warming. The average multi-decadal glacier recession in the Ammassalik region (65° N, 37° W), was examined, rather than the annual range of variability, even though recent observations suggest that major changes in the dynamics of Greenland marine-terminating glaciers take place over timescales of 3–10 yr (Howat et al., 2007; Nick et al., 2009; Andresen et al., 2011; Johannessen et al., 2011), rather than over several decades or centuries as previously believed (Truffer and Fahnestock, 2007).

Here, we examine net frontal position and area fluctuations of 21 marine-terminating glaciers, land-terminating glacier frontal positions for parts of the GrIS, and of 35 GIC on approximately decadal scale for the Ammassalik region (a region including the thermodynamic transition zone from the North Atlantic Ocean into the Arctic Ocean through the Denmark Strait), during the last four decades (1972–2011) using multi spectral Landsat satellite imagery data. Changes were considered in the context of meteorological observations and reconstructed ocean water temperatures time series. As part

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the study the cumulative net area changes for the marine-terminating glaciers was examined. Finally, we investigate differences in marginal change rates between the marine and the terrestrial glacier environments.

2 Data and methods

5 The Landsat and Advanced Spaceborne Thermal and Reflection Radiometer (ASTER) Global Digital Elevation Model Version 2 (GDEM v2) scenes were selected and acquired through WIST (<https://wist.echo.nasa.gov/api/>; EOSDIS, 2009): Landsat 1 carried the four-band Multispectral Scanner (MSS), Landsat 5 carried the seven-band Thematic Mapper (TM), and Landsat 7 carried the eight-band Enhanced Thematic Mapper Plus (ETM+), having a ground resolution of ~60, 30, and 30 m (15 m in the panchromatic band 8), respectively. The ASTER GDEM v2 has a ground resolution of 30 m (Table 1).

15 Data were obtained from seven relatively cloud free scenes (> 25% cloud free) covering the Ammassalik region at the end of the ablation period (mid-August through mid-September) for the time series analysis i.e., 1972, 1986, 1999, 2007, and 2011 (Table 1). For 1972 the GrIS and GIC land-terminating margin analyses were omitted due to difficulties in separating ice cover from snow cover – the year 1972 was one of the coldest years during the study period, and the year with the lowest satellite-derived melt extent cover for GrIS (Mernild et al., 2011b). For 2011, scenes from 2007 were used for filling gaps, due to the ETM+ sensor Scan Line Instrument malfunction. All imagery and features were projected in WGS84, UTM Zone 24N, and the selected scenes were radiometrically calibrated using the Landsat calibration tool in ENVI™ software package (<http://www.itvis.com/ProductServices/ENVI.aspx>), converting the band values to “At Surface Reflectance”. The individual bands (TM and ETM+ bands 1–5, and 7, and MSS bands 2–7) were standardized using the ENVI™ Dark Subtract (DS) tool before ratio and indices were calculated. Errors associated with the different scenes and sensors were ±30 m (MSS), ±15 m (TM), ±15 m (7.5 m panchromatic) (ETM+), and ±30 m (ASTER GDEM v2), and the error associated with the ASTER

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



GDEM v2 is expected to be $\sim \pm 12.5$ m vertically (Table 1) (Tachikawa et al., 2011).

The classification process used for the 1986, 1999, and 2011 scenes was based on a multi-criteria analysis involving the calculation of a set of indices: normalized difference snow index (NDSI; Dozier and Warren, 1982), normalized difference water index (NDWI; Gao, 1996), NDVI (normalized difference vegetation index; Rouse, 1973), and RATIO bands 3/5 TM/ETM+. The RATIO was added due to better performance than the NDSI index in mountainous areas (like the Ammassalik region) capturing ice and snow covered areas influenced by shadows (Paul, 2004). The NDVI was used to filter out vegetation and the NDWI to identify and filter out lakes in the margin area of the GrIS and GIC. The resulting classifications were converted to: (1) polygon files and cleaned up manually in ESRITM ArcMap by visual inspection; (2) poly-line files for visual presentation; and (3) to point shapes with a 30-m distribution along the lines and used to examine the mean and median center of the margin distribution for each year. The 1972-scene was not included in the classification process since the spectral bands of the Landsat 1 MSS sensor used different wavelengths and band distribution than both the TM and ETM+ sensors.

The GrIS and GIC margin positions were digitized for each of the years and the distance between the margins were calculated at the dominant glacier's flow direction using the centerline method (Howat and Eddy, 2011). The expected error related to both the classification and determination of the glacier margin positions closely follows the precision error mentioned in Table 1.

The observed meteorological data, air temperature and precipitation, were obtained from the Danish Meteorological Institute (DMI) station in Tasiilaq for 1972–2011, located less than 10 km from the Sermilik Fjord outlet.

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results and discussion

3.1 Marine-terminating glaciers

For the Ammassalik region, the 21 observed marine-terminating glaciers on average for the survey period 1972–2011 receded at a net rate of $0.053 \pm 0.113 \text{ km yr}^{-1}$ (here and below, the \pm standard deviation are included) with a median of 0.025 km yr^{-1} (Table 2). For 1999–2011, the mean net recession rate was $0.098 \pm 0.171 \text{ km yr}^{-1}$, which is in accordance with the mean net recession rate of 0.11 km yr^{-1} for 210 GrIS marine-terminating outlet glaciers (2000–2010) determined by Howat and Eddy (2011). For the sub-periods the fraction of marine-terminating glaciers in net recession increased from 57 % (1972–1986), 86 % (1986–1999), to 100 % (1999–2011), indicating complete marine-terminating glacier retreat for the Ammassalik region since 1999. For Southeast Greenland, Howat and Eddy (2011) estimated an increasing fraction of net retreat from 55 % (1972–1985), 73 % (1985–2000), to 89 % (2000–2010) ($n = 38$). For the last period, this was almost in the same order of magnitude as the fraction of glacier recession determined for the Ammassalik region.

On individual glacier scale, the glaciers Midgaard, Helheim, Tasis Sassik Fjord, Heims, and Fenris had the first-, second-, third-, fourth-, and fifth-highest net recession rates of 0.365 km yr^{-1} (this equals a net frontal recession of 14.3 km), 0.155 km yr^{-1} (6.0 km), 0.130 km yr^{-1} (5.1 km), 0.075 km yr^{-1} (2.8 km), and 0.065 km yr^{-1} (2.6 km), respectively (Fig. 2) for 1972–2011. All five glaciers are outlets from the GrIS receiving ice from the interior of the ice sheet, having the largest GrIS catchment areas within the Ammassalik region. Overall for the Ammassalik region, 80 % of the marine-terminating glaciers receded at rates less than 0.050 km yr^{-1} (Fig. 2). The mean net recession rates during the period 1972–2011 for the glaciers Midgaard, Helheim, and Fenris compared well with the recession rates determined by Howat and Eddy (2011) for 1972–2010.

For the sub-periods, the 21 observed marine-terminating glaciers had mean frontal recession rates of 0.031 ± 0.056 and $0.029 \pm 0.060 \text{ km yr}^{-1}$ for 1972–1986 and 1986–1999, respectively (significant similar, 97.5 % quartile). For 1999–2011, the mean

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



recession rate increased to $0.098 \pm 0.171 \text{ km yr}^{-1}$, which was significantly higher (95 % quartile) than the two previous sub-periods, and about 3.2 times higher than the 1972–1986 rate. The same trend occurred for the median recession rates, where the 1999–2011 rate was 5.4 times higher than for 1972–1986 (Table 2). The high mean rates for 1999–2011 were mainly due to the recession at the glaciers Midgaard, Helheim, F5, and Fenris, whereas both Midgaard Gletscher and Helheim Gletscher were outside the 75 % percentiles (Figs. 2 and 3a). Also, for 1972–1986 and 1986–1999 Midgaard Gletscher and Helheim Gletscher were considered as outliers. The maximum net advance rate in the dataset was observed for Helheim Gletscher of 0.100 km yr^{-1} (1986–1999). A maximum net recession rate was observed for Midgaard Gletscher of 0.700 km yr^{-1} (1999–2011) (Figs. 1 and 3a). For the Ammassalik region in general, the mean recession rate increased since 1972 (Fig. 3a), as was also reported by Howat and Eddy (2011) for the GrIS outlet glaciers, where a transition was described from stability and small fluctuations in glacier frontal positions (1972–1985) to moderate widespread recession in the southeast and western parts of GrIS (1985–2000), followed by an accelerated net recession in all regions of the ice sheet (2000–2010).

Since 1972, the 21 marine-terminating glaciers showed a cumulative net area loss of 281 km^2 , with mean annual area loss rates of $7.2 \text{ km}^2 \text{ yr}^{-1}$ (Fig. 4). The largest individual marine-terminating area loss occurred at Midgaard Gletscher of 130 km^2 (equal to 46 % of the cumulative net area exposure in the Ammassalik region), followed by Helheim Gletscher of 67 km^2 (equal to 24 %). About 70 % of the glaciers had an area loss less than 4 km^2 during the period 1972–2011. Also, the 21 observed marine-terminating glaciers had a trend towards lower annual area loss rates (Fig. 4): for the sub-periods 1972–1986, 1986–1999, and 1999–2011 the mean area loss rates were $8.5 \text{ km}^2 \text{ yr}^{-1}$ (equal to 118 km^2), $6.8 \text{ km}^2 \text{ yr}^{-1}$ (88 km^2), and $6.3 \text{ km}^2 \text{ yr}^{-1}$ (75 km^2), respectively (Fig. 4), indicating decreasing area loss rates since 1972. Box and Decker (2011) measured area change at 39 of the widest GrIS marine-terminating glaciers based on Moderate Resolution Imaging Spectroradiometer (MODIS) between 2000 and 2010, including the glaciers Midgaard, Helheim, and Fenris. Overall for the

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



39 glaciers, they exposed a cumulated net area of 1368 km². On individual glacier scale the observed area loss rates for the glaciers Midgaard, Helheim, and Fenris were -2.9, -2.0, and -0.4 km² yr⁻¹ for 1999–2011, respectively, compared to -3.6, -2.5, and -0.3 km² yr⁻¹ for 2000–2010 determined by Box and Decker (2011). The Landsat derived area loss rates were in the same range as Box and Decker (2011) (97.5 % quartile), and both studies indicated that the clearest pattern of area exposure happened at the Midgaard Gletscher in the Sermilik Fjord, even compared with the entire Southeast Greenland according to Box and Decker (2011).

Several authors (e.g., Holland et al., 2008; Murray et al., 2010; Straneo et al., 2010; Andresen et al., 2011) suggest that ocean warming and warm subsurface waters caused large changes to marine-terminating glacier frontal positions and ice discharge activity. In recent decades the 21 marine-terminating glaciers in the Ammassalik region have experienced increasing frontal recession rates and decreasing area exposure rates that are synchronous with both increasing mean annual air temperature (MAAT) (~ 0.06 °C yr⁻¹, significant at $p < 0.01$, where p is level of significance; from the DMI meteorological station in Tasiilaq) and reconstructed annual ocean water temperature (~ 0.12 °C yr⁻¹, significant $p < 0.025$; at 400 m depth in the Irminger Sea penetrating into the Sermilik Fjord and exposing the lower part of glaciers such as Helheim and Fenris to warm waters with temperatures up to 4 °C (Johannessen et al., 2011)) (Figs. 3b and 4). The mean glacier retreat was more widespread for 1999–2011 (approximately the first decade of the 21st century), than for the earlier sub-periods (1972–1986 and 1986–1999). The observed accelerated recession of the marine-terminating glaciers in the Ammassalik region coincided with the onset of a warming trend in the sub-polar North Atlantic Ocean (Myers et al., 2007; Straneo et al., 2010), likely initiated by the influx of warmer deep water originating in the Irminger Sea (Holland et al., 2008; Hanna et al., 2009). This supports the hypothesis that ocean warming associated with shifts in the Irminger and East Greenland currents caused increasing submarine melt at the ice-ocean interface. However, oceanographic studies have demonstrated that although subtropical ocean waters reach glacial fjords in Southeast Greenland, there

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



decreased ~ 20 % in total length from 686 km (1986) to 544 km (2011), because the 2011-margin was less curved – had fewer land-terminating outlets – than in 1986 and 1999. When the area recession is compared to changes in margin length, the largest GrIS area length recession ratio occurred at the elevation > 800 m a.s.l., most pronounced for the period 1999–2011 (Fig. 6). The spatial area recession seems to be highly influenced by local topography, hypsometry, and its shadow effects, climate variability, glacier dynamic processes within the GrIS, increasing ELA elevation (the ELA is the spatially averaged elevation of the equilibrium line, defined as the set of points on the glacier surface where the net mass balance is zero), and the margin elevation distribution, where approximately 20 % of the margin was located between 701–800 m a.s.l. (1986–2011) (Fig. 6).

The mean net recession rate of the GrIS land-terminating margin was $0.018 \pm 0.009 \text{ km yr}^{-1}$ (equal to a net recession of 0.443 km) for 1986–2011, comprised of a mean recession rate of 0.010 km yr^{-1} (0.127 km) for 1986–1999, and 0.026 km yr^{-1} (0.316 km) for 1999–2011 (Table 3). This land-terminating recession rate for the GrIS over this period of 1986–2011 is about three times lower the mean rate of recession of the marine-terminating GrIS. Sohn et al. (1998) measured recession rates of the GrIS land-terminating margin near Jakobshavn Isbræ, West Greenland, of $0.016\text{--}0.040 \text{ km yr}^{-1}$, averaging 0.026 km yr^{-1} 1962–1992. This shows the recession rate can be expected to be within this order of magnitude along many parts of the GrIS land-terminating margin.

Net GrIS land-terminating marginal recession for the Ammassalik region, including increasing area exposure, occurred for the period 1972–2011, during a period of increasing MAAT ($\sim 0.06 \text{ }^\circ\text{C yr}^{-1}$) and decreasing annual precipitation ($-7.0 \text{ mm water equivalent (w.e.) yr}^{-1}$, significant at $p < 0.025$; Fig. 3b) – probably heading towards future warmer and dryer conditions in the region (Mernild et al., 2011a). The average increase in MAAT generally favors surface ablation (evaporation, sublimation, and melt), and an earlier start of the ablation season by decreasing the “cold content” of the snowpack (Bøggild et al., 2005; Mernild et al., 2011), whereas a decrease in annual

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



precipitation may lead to earlier exposure of glacier ice melt and summer firn surface of previous years (having a lower albedo than fresh snow promoting increased solar absorption). Therefore, the combination of increasing air temperature and decreasing precipitation is likely to increase ablation and GrIS margin thinning and recession, and if MAAT and precipitation continue to follow these trends, then it is expected that the GrIS land-terminating margin will continue its recession, leading to increased area exposure. However, changes in the hypsometric distribution along the GrIS margin may influence recession rates on a decadal scale.

3.3 Land-terminating glaciers and ice caps

Peripheral to the GrIS 35 land-terminating GIC were chosen (representative of GIC conditions in size and elevation) (Figs. 1 and 7) - both large ($>10\text{ km}^2$) and small ($<10\text{ km}^2$) GIC to assess area exposure for the Ammassalik region (Fig. 7a). Glacier extent was estimated for 1986, 1999, and 2011 based on Landsat imagery. The 35 GIC are non-surging glaciers located south of the East Greenland surge cluster (Jiskoot et al., 2003). For the Ammassalik region, the observed GIC indicated a relative mean area exposure of $4 \pm 18\%$ for 1986–1999, and $27 \pm 24\%$ for 1986–2011 (Fig. 7b), which is equal to a mean net area exposure rate of $0.04\text{ km}^2\text{ yr}^{-1}$ per glacier (or $1.06\% \text{ yr}^{-1}$ per glacier). For small GIC ($n = 32$) the net area exposure rate was on average $1.08\% \text{ yr}^{-1}$, and for large GIC ($n = 3$) was a comparable rate of $0.87\% \text{ yr}^{-1}$. For 1986–1999 eleven individual GIC (around 30%, mostly below 2 km^2) had a net increase in area, while for 1986–2011 it was only three GIC all facing towards the west (L11, L14, and L9; and around 10% – all less than 1 km^2) (Fig. 7b). As illustrated in Fig. 7c, GIC having a mean elevation height (defined as the maximum plus minimum elevations divided by two) higher than 685 m a.s.l. had in general a net area increase from 1986 to 1999, while glaciers with a mean elevation lower than 685 m a.s.l. had a net area decrease (based on the significant linear regression; $r^2 = 0.31$; $p < 0.01$). The height of 685 m a.s.l. was around the observed average ELA of 690 m a.s.l. at the Mittivakkat Gletscher in the late 1990s (Knudsen and Hasholt, 2004; Mernild et al.,

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2011a; Table 2). For 1999–2011 the linear regression shown in Fig. 7c indicates the opposite trend for GIC in the Ammassalik region: an increase in area recession for GIC at high elevation ranges, and vice versa. This shift in trend occurred simultaneously with an increase in the average observed ELA for the Mittivakkat Gletscher to 750 m a.s.l. Overall for 1986–2011, the observed GIC faced a general net area loss that was highest at low elevations, and vice versa (based on the linear regression; Fig. 7c). However, as previously mentioned three minor GIC (<1 km²) had a net area gain during this 25-yr period, indicating that glacier fluctuations may vary on local scales.

Since the GIC on average had a mean net area exposure rate of about 1 % yr⁻¹, it may be expected that GIC in the Ammassalik region could melt substantially in the 21st century under ongoing climate change. For the period 1986–2011 there are examples of five glaciers which completely melted away: all located at different mean elevations within the region from 500 to 1130 m a.s.l., and with different aspects facing from east to west (Table 4). The recession seems therefore not to be limited to low elevated areas only, but more likely to occur for north facing GIC. Also, in Fig. 8, examples of eight GIC are shown to illustrate the spatial changes in margin location from 1986 to 2011.

For the largest GIC – the Mittivakkat Gletscher (26.3 km² in 2011) the area extent had diminished about 18 % since 1986 (lower than the mean GIC area exposure for the Ammassalik region of 27 ± 24 %). The terminus has retreated by 1.6 km (0.015 km yr⁻¹) since the maximum extent of the Little Ice Age around 1900, by 1.3 km (0.017 km yr⁻¹) since 1931 (Humlum and Christiansen, 2008; Mernild et al., 2011a), and by 0.3 km (0.013 km yr⁻¹) since 1986, which is almost of the same magnitude as the GIC of the Ammassalik region's mean net recession rate of 0.010 ± 0.006 km yr⁻¹ (1986–2011), and of the regional GrIS land-terminating margin of 0.018 ± 0.009 km yr⁻¹ (see Sect. 3.2): the mean Ammassalik GIC land-terminating recession rate (1986–2011) is about five times lower than the mean GrIS marine-terminating recession rate. Also, for Mittivakkat Gletscher the annual mass balance measured continuously since 1995/96 illustrates a 16-yr average mass loss of 0.970 ± 0.190 m w.e. yr⁻¹, indicating that the glacier is significantly out of balance with the current climate. The glacier will likely lose

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



at least 70 % of its current area extent and 80 % of its volume even in the absence of further climate changes (Mernild et al., 2011a). Since the initiation of the mass balance observation program in 1995/1996, Mittivakkat Gletscher had in 14 out of 16 yr a negative surface mass balance, while the general climatic trend in the region has been towards higher temperatures, less winter precipitation, and more negative glacier mass balances and continuous marginal recession (Fig. 8). Consecutive record glacier mass loss occurred for the years 2009/2010 and 2010/2011 of -2.16 and -2.45 m w.e. yr^{-1} , respectively. The 2011 mass loss was not only the largest annual loss of volume in the history of the mass balance observational program, but also the largest annual loss in simulations of glacier mass balance changes back to 1898 (Mernild et al., 2008). The marginal recession and mass balance observations suggest that recent Mittivakkat mass losses, which have been driven largely by higher surface temperatures and less solid precipitation, are representative of the broader region, which includes the 35 observed GIC in Figs. 7 and 8 – glaciers of different sizes and elevation ranges. This is confirmed since the Mittivakkat Gletscher net area exposure rate closely follows the average rates for the Ammassalik region.

Glacier fluctuations and area exposure have been studied in other parts of Greenland. North of the Ammassalik region, in Central East Greenland (68° N– 72° N) land-terminating GIC peripheral to the GrIS have receded at a mean rate of 0.010 km yr^{-1} for a wide range of glacier sizes (2002–2009) (Kargel et al., 2011). Also, on Disko Island in West Greenland, Yde and Knudsen (2007) estimated mean GIC terminus retreat rates of 0.008 km yr^{-1} for non-surging GIC, and 0.020 km yr^{-1} for quiescent phase surge-type GIC, reflecting a higher non-climatic-driven recession rate after glacier surges. These studies are in accordance with the findings for the Ammassalik region and indicate that the current mean recession rate for GIC in Greenland (probably excluding North Greenland where no data is currently available) is likely to be on the order of 0.008 – 0.010 km yr^{-1} .

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Summary and conclusion

The satellite observations show net glacier recession since 1972 for the Ammassalik region for all glacier types and sizes, both at the marine-terminating and land-terminating GrIS, and land-terminating GIC. However, the land-terminating GrIS and GIC reflect slower area exposure rates than the faster marine-terminating outlet glaciers. This could likely be due to a combination of effects, as the marine-terminating GrIS was influenced by the onset of a sea-water warming trend in the North Atlantic Ocean (Myers et al., 2007; Straneo et al., 2010), and by atmospheric impacts from regional trends in MAAT and precipitation, while GIC were only influenced by the latter. For the marine-terminating GrIS outlet glaciers the mean annual area exposure rate has decreased since 1972, whereas it has increased for the land-terminating GrIS margin since 1986, even though both parts of the GrIS have undergone substantial area changes in the past decades. The observed land-terminating GrIS and GIC indicate a net area recession of 4% (equal to an area exposure rate of $0.15\% \text{ yr}^{-1}$) and $27 \pm 24\%$ (around $1\% \text{ yr}^{-1}$), respectively, and margin recession rates of 0.018 ± 0.009 and $0.010 \pm 0.006 \text{ km yr}^{-1}$. These mean net margin recession rates are about 3 to 5 times lower than the GrIS marine-terminating margin rates. If these GIC recession trends are extrapolated it would indicate that a substantial amount of the GIC in the Ammassalik region might melt away within the 21st century under ongoing climate warming. So far, five GIC in the Ammassalik region have melted away since 1986.

Acknowledgements. The authors thank Matthew Hecht and Mauri Pelto for their insightful review of an early version of this paper. We also thank Ola M. Johannessen for providing the reconstructed ocean water data. This work was partly supported by the Climate Change Prediction Program and by the Scientific Discovery for Advanced Computing (SciDAC) program within the US Department of Energy's Office of Science, Los Alamos National Laboratory (LANL) Director's Fellowship, and LANL Institute for Geophysics and Planetary Physics (LANL is operated under the auspices of the National Nuclear Security Administration of the US Department of Energy under Contract No. DE-AC52-06NA25396), and partly from the European Community's Seventh Framework Programme under grant agreement No. 262693. SHM and

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



JKM compiled the satellite images, analyzed the data, and wrote the manuscript. NTK and JCY contributed to the discussion of results and writing of the text.

References

- Andresen, C. S., Straneo, F., Rieberggaard, M. H., Bjørk, A. A., Andersen, T. J., Kuijpers, A., Nørgaard-Pedersen, N., Kjær, K. H., Schjøth, F., Weckstrom, K., and Ahlstrøm A. P.: Rapid response of Helheim Glacier in Greenland to climate variability over the past century, *Nat. Geosci.*, 5, 37–41, doi:10.1038/NCEO1349, 2011.
- Bøggild, C. E., Forsberg, R., and Reeh, N.: Melt: water in a transect across the Greenland ice sheet, *Ann. Glaciol.*, 40, 169–173, 2005.
- Box, J. E. and Decker D. T.: Greenland marine-terminating glacier area changes: 2000–2010, *Ann. Glaciol.*, 52, 91–98, 2011.
- Dozier, J. and Warren, S. G.: Effect of viewing angle on the infrared brightness temperature of snow, *Water Resour. Res.*, 18, 1424–1434, 1982.
- EOSDIS: Earth Observing System Data and Information System (EOSDIS). 2009. Earth Observing System ClearingHouse (ECHO)/Warehouse Inventory Search Tool (WIST) Version 10.X. Greenbelt, MD: EOSDIS, Goddard Space Flight Center (GSFC) NASA, 2009.
- Gao, B.-C: NDWI – a normalized difference water index for remote sensing of vegetation liquid water from space, *Remote Sens. Environ.*, 58, 257–266, 1996.
- Haerberli, W., Gartner-Roer, I., Hoelzle, M., Paul, F., and Zemp, M. (Eds.): WGMS: Glacier Mass Balance Bulletin. Bulletin No. 10 (2006–2007), ICSU(WDS)/IUGG(IACS)/UNEP/UNESCO/WMO, World Glacier Monitoring Service, Zurich, 96 pp., 2009.
- Hanna, E., Cappelen, J., Fettweis, X., Huybrechts, P., Luckman, A., and Ribergaard, M. H.: Hydrologic response of the Greenland ice sheet: the role of oceanographic warming, *Hydrol. Process.*, 23, 7–30, 2009.
- Hansen, J., Ruedy, R., Sato, M., and Lo, K.: Global surface temperature change, *Rev. Geophys.*, 48, RG4004, doi:10.1029/2010RG00345, 2010.
- Holland, D. M., Thomas, R. H., de Young, B., Ribergaard, M. H., and Lyberth, B.: Acceleration of Jakobshavn Isbrae triggered by warm subsurface ocean waters. *Nat. Geosci.*, 1, 659–662, 2008.

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Howat, I. M. and Eddy, A.: Multi-decadal retreat of Greenland's marine-terminating glaciers, *J. Glaciol.*, 57, 389–396, 2011.
- Howat, I. M., Joughin, I., Tulaczyk, S., and Gogineni, S.: Rapid retreat and acceleration of Helheim Glacier, East Greenland, *Geophys. Res. Lett.*, 32, L22502, doi:10.1029/2005GL024737, 2005.
- Howat, I. M., Joughin, I., and Scambos, T. A.: Rapid changes in ice discharge from Greenland outlet glaciers, *Science*, 315, 1559–1561, 2007.
- Howat, I. M., Joughin, I., Fahnestock, M., Smith, B. E., and Scambos, T.: Synchronous retreat and acceleration of southeast Greenland outlet glaciers 2000–2006: ice dynamic and coupling to climate, *J. Glaciol.*, 54, 646–660, 2008.
- Humlum, O. and Christiansen, H. H.: Geomorphology of the Ammassalik Island, SE Greenland, *Geogr. Tidsskr.*, 108, 5–20, 2008.
- Jiskoot, H., Murray, T., and Luckman, A.: Surge potential and drainage-basin characteristics in East Greenland, *Ann. Glaciol.*, 36, 142–148, 2003.
- Johannessen, O. M., Korabely, A., Miles, V., Miles, M. W., and Solberg, K. E.: Interaction between the warm subsurface Atlantic water in the Sermilik Fjord and Helheim Glacier in Southeast Greenland, *Surv. Geophys.*, 32, 387–396, doi:10.1007/s10712-011-9130-6, 2011.
- Joughin, I., Abdalati, W., and Fahnestock, M. A.: Large fluctuations in speed on Greenland's Jakobshavn Isbrae glacier, *Nature*, 432, 608–610, 2004.
- Joughin, I., Smith, B. E., Howat, I. M., Scambos, T., and Moon, T.: Greenland flow variability from ice-sheet-wide velocity mapping, *J. Glaciol.*, 56, 415–430, 2010.
- Kargel, J. S., Ahlstrøm, A. P., Alley, R. B., Bamber, J. L., Benham, T. J., Box, J. E., Chen, C., Christoffersen, P., Citterio, M., Cogley, J. G., Jiskoot, H., Leonard, G. J., Morin, P., Scambos, T., Sheldon, T., and Willis, I.: Brief Communication: Greenland's shrinking ice cover: “fast times” but not that fast, *The Cryosphere Discuss.*, 5, 3207–3219, doi:10.5194/tcd-5-3207-2011, 2011.
- Knudsen, N. T. and Hasholt, B.: Mass balance observations at Mittivakkat Glacier, Southeast Greenland 1995–2002, *Nord. Hydrol.*, 35, 381–390, 2004.
- Luthcke, S., Zwally, H. J., Abdalati, W., Rowlands, D. D., Ray, R. D., Nerem, R. S., Lemoine, F. G., McCarthy, J. J., and Chinn, D. S.: Recent Greenland mass loss by drainage system from satellite gravity observations, *Science*, 314, 1286–1289, 2006.
- Mernild, S. H., Kane, D. L., Hansen, B. U., Jakobsen, B. H., Hasholt, B., and Knudsen, N. T.: Climate, glacier mass balance, and runoff (1993–2005) for the Mittivakkat Glacier catchment,

Multi-decadal marine and land-terminating glacier recessionS. H. Mernild et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Multi-decadal marine and land-terminating glacier recessionS. H. Mernild et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ammassalik Island, SE Greenland, and in a long term perspective (1898–1993), *Hydrol. Res.*, 39, 239–256, 2008.

Mernild, S. H., Knudsen, N. T., Lipscomb, W. H., Yde, J. C., Malmros, J. K., Hasholt, B., and Jakobsen, B. H.: Increasing mass loss from Greenland's Mittivakkat Gletscher, *The Cryosphere*, 5, 341–348, doi:10.5194/tc-5-341-2011, 2011a.

Mernild, S. H., Mote, T., and Liston, G. E.: Greenland Ice Sheet surface melt extent and trends, 1960–2010, *J. Glaciol.*, 57, 621–628, 2011b.

Mernild, S. H., Seidenkrantz, M.-S., Chylek, P., Liston, G. E., and Hasholt, B.: Climate-driven fluctuations in freshwater to Sermilik Fjord, East Greenland, during the last 4000 years, *Holocene*, 22(2), 155–164, doi:10.1177/0959683611431215, 2012a.

Mernild, S. H., Hanna, E., Knudsen, N. T., Yde, J. C., and Seidenkrantz, M.-S.: The climate impact on observed land terminating glaciers and ice caps in Northern North Atlantic, *J. Climate*, in review, 2012b.

Moon, T. K. and Joughin, I.: Retreat and advance of Greenland tidewater glaciers from 1992 to 2007, *J. Geophys. Res.*, 113, F02022, doi:10.1029/2007JF000927, 2008.

Murry T., Scharrer, K., James, T. D., Dye, S. R., Hanna, E., Booth, A. D., Selmes, N., Luckman, A., Hughes, A. L. C., Cook, S., and Huybrechts, P.: Ocean regulation hypothesis for glacier dynamics in southeast Greenland and implications for ice sheet mass changes, *J. Geophys. Res.*, 115, F03026. doi:10.1029/2009JF001522, 2010.

Myers, P., Kulan, N., and Ribergaard, M. H.: Irminger water variability in the West Greenland current, *Geophys. Res. Lett.*, 34, L17601, doi:10.1029/2007GL030419, 2007.

Nick, F., Vieli, M. A., Howat, I. M., and Joughin, I.: Large-scale changes in Greenland outlet glacier dynamics triggered at the terminus, *Nat. Geosci.*, 2, 110–114, doi:10.1038/NGEO394, 2009.

Paul, F.: *The New Swiss Glacier Inventory 2000 – Application of remote sensing and GIS*, PhD Thesis, Department of Geography, University of Zurich, 198 pp., 2004.

Radić, V. and Hock, R.: Regionally differentiated contribution of mountain glaciers and ice caps to future sea-level rise, *Nat. Geosci.*, 4, 91–94, 2011.

Rignot, E. and Kanagaratnam, P.: Changes in the velocity structure of the Greenland Ice Sheet, *Science*, 311, 986–990, 2006.

Rignot, E., Knoppes, M., and Velicogna, I.: Rapid submarine melting of the calving faces of West Greenland glaciers. *Nat. Geosci.*, 3, 187–191, 2010.

Rouse, J. W., Haas, R. H., Schell, J. A., and Deering, D. W.: Monitoring vegetation systems in

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the Great Plains with ERTS, Third ERTS Symposium, NASA SP-351 I, 309–317, 1973.

Sohn, H.-G., Jezek, K. C., van der Veen, C. J.: Jakobshavn Glacier, West Greenland: 30 years of spaceborne observations, *Geophys. Res. Lett.*, 25, 2699–2702, 1998.

5 Straneo, F., Hamilton, G. S., Sutherland, D. A., Sterns, L. A., Davidson, F., Hammill, M. O., Stenson, G. B., and Rosing-Asvid, A.: Rapid circulation of warm subtropical waters in a major glacial fjord in East Greenland, *Nat. Geosci.*, 3, 182–222, 2010.

Tachikawa, T., Kaku, M., Iwasaki, A., Gesch, D., Oimoen, M., Zhang, Z., Danielson, J., Krieger, T., Curtis, B., Haase, J., Abrams, M., Crippen, R., and Carabaja, C.: ASTER Global Digital Elevation Model Version 2 – Summary of Validation Results. NASA Land Processes Distributed Active Archive Center and the Joint Japan-US ASTER Science Team, available at: https://igskmncnwb001.cr.usgs.gov/aster/GDEM/Summary_GDEM2_validation_report_final.pdf, 2011.

10 Truffer, M. and Fahnestock, M.: Rethinking ice sheet time scales, *Science*, 315, 1508–1510, 2007.

15 van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E., van de Berg, W., van Meijgaard, E., Velicogna, I., and Wouters, B.: Partitioning recent Greenland mass loss, *Science*, 326, 984–986, 2009.

Velicogna, I.: Increasing rates of ice mass loss from the Greenland and Antarctic ice sheets revealed by GRACE, *Geophys. Res. Lett.*, 36, L19503, doi:10.1029/2009GL040222, 2009

20 Velicogna, I. and Wahr, J.: Acceleration of Greenland ice mass loss in spring 2004, *Nature*, 443, 329–331, 2006.

Walsh, K. M., Howat, I. M., Ahn, Y., and Enderlin, E. M.: Changes in the marine-terminating glaciers of central east Greenland and potential connections to ocean circulation, 2000–2010, *The Cryosphere Discuss.*, 5, 2865–2894, doi:10.5194/tcd-5-2865-2011, 2011.

25 Yde, J. C. and Knudsen, N. T.: 20th-century glacier fluctuation on Disko Island (Qeqertarsuaq), Greenland, *Ann. Glaciol.*, 46, 209–214, 2007.

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Table 1. Satellite platform, sensors, and band information and scenes used in the analysis.

Platform	Landsat 1	Landsat 5	Landsat 7	Terra
Sensor and bands	Landsat MSS (bands 2–7)	Landsat TM (bands 1–5, and 7)	Landsat ETM+ (bands 1–5, and 7)	ASTER GDEM Version 2
Ground resolution	~60 m	30 m	30 m, 15 m pancromatic	30 m
Precision error	±30 m (horizontal)	±15 m (horizontal)	±15 m, 7.5 m pancromatic (horizontal)	~ ±12.5 m (vertical)
Scenes	M1251013.01319720907	L5231014.01419860911	L71231014.01419990907 L71232013.01320110814 L71232014.01420110814 L71232014.01420070904 L71231014.01420070913	20111017145633.1819068388
Survey years and dates	7 Sep 1972	11 Sep 1986	7 Sep 1999 13 Sep 2007 14 Aug 2011	–

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Table 2. Frontal position change rate statistics for the period 1972–2011 for the GrIS marine-terminating glaciers in the Ammassalik region, mean annual air temperature and precipitation anomaly from the DMI meteorological station in Tasiilaq, and mean annual reconstructed ocean water temperature anomaly at 400 m depth in the Irminger Sea (Johannessen et al., 2011).

Range	Number of GrIS marine-terminating glaciers	Mean annual air temperature anomaly and standard deviation ($^{\circ}\text{C yr}^{-1}$)	Mean annual precipitation anomaly and standard deviation (mm w.e. yr^{-1})	Mean annual reconstructed ocean water temperature anomaly and standard deviation ($^{\circ}\text{C yr}^{-1}$)	Glacier fraction in net retreat (%)	Mean change rate and standard deviation (km yr^{-1})	Median change rate (km yr^{-1})	Maximum advance rate (km yr^{-1})	Maximum recession rate (km yr^{-1})
1972–1986	21	-0.63 ± 0.78	69 ± 212	-0.12 ± 0.30	57	-0.031 ± 0.056	-0.007	0.005	-0.223
1986–1999	21	-0.33 ± 0.63	-17 ± 238	-0.06 ± 0.49	86	-0.029 ± 0.060	-0.009	0.100	-0.187
1999–2011	21	1.08 ± 0.59	-62 ± 163	$0.23 \pm 0.43^*$	100	-0.098 ± 0.171	-0.038	-0.003	-0.758
1972–2011	21	–	–	–	100	-0.053 ± 0.113	-0.025	0.100	-0.758

* Data are missing from 1995 and 1996, and only present to 2009.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Table 3. GrIS land-terminating margin-position change rate statistics for each survey year and period for the Ammassalik region.

Years	GrIS margin length for the highlighted section in Ammassalik region (km)	GrIS area for the section for the highlighted section in Ammassalik region (km ²)	Area in percentage related to the 1986 area (%)
1986	686	1166	100
1999	589	1153	99
2011	544	1124	96
	GrIS mean margin change rate (km yr ⁻¹)	GrIS mean margin change (km)	
1986–1999	–0.010	–0.127	
1999–2011	–0.026	–0.316	
1986–2011	–0.018	–0.443	

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Table 4. Characteristics of land-terminating GIC L33–L37 in the Ammassalik region, which have melted away during the period 1986–2011 (see Fig. 1 for location of the doomed glaciers).

GIC	Minimum elevation (m a.s.l.)	Maximum elevation (m a.s.l.)	Mean elevation (m a.s.l.)	Aspect	Area 1986 (km ²)	Area 1999 (km ²)	Area 2011 (km ²)
L33	507	792	650	E	0.178	0.050	0
L34	462	538	500	E	0.041	0.034	0
L35	846	910	878	W	0.036	0	0
L36	777	896	837	W	0.076	0.016	0
L37	1107	1145	1126	SW	0.018	0.029	0

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. The Ammassalik region, including the Sermilik Fjord and parts of the southeastern sector of the GrIS. The marine-terminating glaciers margins are marked with positions (lines) for each of the survey years 1972 (red color), 1986 (green), 1999 (blue), and 2011 (yellow) (the location names are written in turquoise). The 35 local land-terminating glaciers and ice caps (GIC) peripheral to the GrIS are written in yellow, where eight are marked with shaded areas – marginal changes for those eight GIC and the GrIS are detailed illustrated. The GIC marked with L33–L37 are examples of GIC melted away during 1986–2011. The inset figure indicates the general location of the Ammassalik region in Southeast Greenland (source: Landsat 7 ETM+ Mosaik, 7 September 1999/9 September 2000).

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.




Fig. 2. Satellite-derived net frontal recession for the 21 marine-terminating glaciers in the Ammassalik region for the sub-periods 1972–1986 (green), 1986–1999 (blue), and 1999–2011 (yellow). For glacier locations, see Fig. 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. (a) Box plots of frontal-position change for the sample of 21 marine-terminating glaciers in the Ammassalik region with measurements in each of the 1972–1986, 1986–1999, and 1999–2011 survey periods. The edges of the boxes denote 25 % and 75 % percentiles and the vertical line mean. Data points outside this range are considered outliers and are plotted as crosses and labeled; **(b)** Mean annual air temperature anomaly (observed at the DMI meteorological station in Tasiilaq), mean annual ocean water temperature anomaly at 400 m depth in the Irminger Sea (Johannessen et al., 2011), mean annual precipitation anomaly (uncorrected) (observed at the DMI meteorological station in Tasiilaq) and standard deviations are shown.

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.




Fig. 4. Mean annual area change rates and cumulative net area change for the 21 marine-terminating glaciers for the Ammassalik region 1972–2011.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc



Printer-friendly Version

Interactive Discussion




Greenland Ice Sheet 1986 - 2011
(land-terminating margin)

Elevation (m)



ASTER GDEM v2

Margin change



Landsat 7, September 1999

Fig. 5. The location of the GrIS land-terminating margin for the survey years 1986 (green), 1999 (blue), and 2011 (yellow) in the Ammassalik region, and the marginal changes between the survey years. The topography has black and gray shaded colors (source: ASTER GDEM v2 and Landsat 7).

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.




Fig. 6. GrIS land-terminating area recession rates for different elevation intervals, Ammassalik region, for the sub-periods 1986–1999 (light gray) and 1999–2011 (dark gray), area recession rate related to changes in margin length for 1986–1999 (black dashed line, long dashed) and 1999–2011 (black dashed line, short dashed), and the percentage of margin elevation for the different elevation intervals for the years 1986 (green), 1999 (blue), and 2011 (yellow) (see Fig. 5 for location of the margin).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. (a) Land-terminating GIC area for the 35 observed GIC (Mittivakkat, Hobbs, Tinit, and L1–L32 glaciers) for 1986, 1999, and 2011 (the listed glaciers are illustrated as an example in Fig. 8); (b) relative GIC area change since 1986 (the named glaciers are the ones with area increase); and (c) relative GIC area change in relation to variations in mean elevation GIC height for 1986–1999, 1999–2011, and overall for 1986–2011.

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 8. The margin location of eight land-terminating GIC (peripheral to the GrIS) in the Ammassalik region: Hobbs, Mittivakkat, Tinit, and L1–L5 glaciers for the survey years 1986 (green), 1999 (blue), and 2011 (yellow) estimated from Landsat images. The location of the individual glaciers is shown in Fig. 1 (source: Landsat 7 ETM+ Mosaik, 7 September 1999/9 September 2000).

Multi-decadal marine and land-terminating glacier recession

S. H. Mernild et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

