

The growth of
sublimation crystals
and surface hoar

J.-C. Gallet et al.

This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

The growth of sublimation crystals and surface hoar on the Antarctic plateau

J.-C. Gallet^{1,2,3}, F. Domine^{1,2,4}, J. Savarino^{1,2}, M. Dumont⁵, and E. Brun⁶

¹CNRS, LGGE, 38041 Grenoble, France

²Université Grenoble Alpes, LGGE, 38041 Grenoble, France

³Norwegian Polar Institute, Tromsø, Norway

⁴Takuvik Joint International Laboratory and Department of Chemistry, Université Laval and CNRS, Quebec, Canada

⁵Météo-France-CNRS, CNRM-GAME UMR 3589, CEN, Saint-Martin d'Hères, France

⁶Météo-France-CNRS, CNRM-GAME UMR 3589, Toulouse, France

Received: 26 November 2013 – Accepted: 29 November 2013

– Published: 13 December 2013

Correspondence to: F. Domine (florent.domine@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

On the Antarctic plateau, the budget of water vapor and energy is in part determined by precipitation, but these are so low that the dynamic of snow crystal growth and sublimation at the surface can be important factors. At Dome C (75° S, 123° E), we have frequently observed the growth of crystals on the snow surface under calm sunny weather. Here, we present the time variations of specific surface area and density of these crystals. Using the detailed snow model Crocus, we conclude that these crystals were very likely due to the nighttime formation of surface hoar crystals and to the daytime formation of sublimation crystals. These latter crystals form by processes similar to those involved in the formation of frost flowers on young sea ice. The formation of these crystals impact the albedo, mass and energy budget of the Antarctic plateau. In particular, the specific surface area variations of the surface layer can induce an instantaneous forcing of up to -10 W m^{-2} at noon, resulting in a surface temperature drop of 0.45 K.

1 Introduction

Snow is the most reflective surface on Earth. It can covers up to 50 % of land masses in the Northern Hemisphere in winter (Robinson et al., 1993), and nearly all of Greenland and Antarctica year round. Snow therefore considerably affects the energy budget of the Earth.

At temperatures below 0°C, snow is a porous medium made of air, ice and small amounts of impurities. Because of the large surface area of the ice–air interface, snow is thermodynamically unstable and is therefore in constant evolution, the most visible modifications being changes in the shapes and sizes of snow grains. The set of physical processes responsible for these modifications is called metamorphism (Colbeck, 1982). Metamorphism also affects macroscopic physical properties such as density, thermal conductivity, and albedo.

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In the visible, snow albedo is mostly dependant on its impurity content while it depends on the size of the snow grains in the infrared part of the solar spectrum (Warren, 1982). The size of the snow grains usually increases during metamorphism (Taillandier et al., 2007; Flanner and Zender, 2006) so that in general snow albedo decreases over time. Understanding snow grain size variations is therefore crucial for an accurate calculation of the Earth's energy budget.

Given the complex shapes of snow particles, defining grain size unambiguously has been difficult (Aoki et al., 2000). However, it has been shown that the albedo of non-spherical particles was adequately represented by a collection of spherical particles having the same surface to volume (S/V) ratio as the non-spherical particles, and the size of those spheres define the optical radius r_{opt} unambiguously (Grenfell and Warren, 1999). For optical calculations, r_{opt} of snow crystals has been used successfully. In practice, r_{opt} can be obtained by measuring the specific surface area (SSA) of the snow, defined as the surface area of the air–ice interface per unit mass: $SSA = S/M = S/(\rho_{ice} V)$, where ρ_{ice} is the ice density, 917 kg m^{-3} at 0°C . Treating snow crystals as spheres, we then have $SSA = 3/(\rho_{ice} r_{opt})$, and SSA measurements have been used successfully for albedo calculations (Gallet et al., 2011). Snow SSA is expressed in units of $\text{m}^2 \text{ kg}^{-1}$. A compilation of measured values ranges from $2 \text{ m}^2 \text{ kg}^{-1}$ ($r_{opt} = 1.64 \text{ mm}$) for melt-freeze crusts to $156 \text{ m}^2 \text{ kg}^{-1}$ ($r_{opt} = 21.0 \mu\text{m}$) for fresh dendritic crystals (Domine et al., 2007). Recently, a value of $223 \text{ m}^2 \text{ kg}^{-1}$ ($r_{opt} = 14.7 \mu\text{m}$) has been measured for diamond dust crystals (i.e. clear sky precipitation) in the Arctic (Domine et al., 2011b).

On a daily time-scale, variations of the factors affecting the energy budget of the Antarctic ice cap (Van As et al., 2005; Van Den Broeke et al., 2006) can result in sublimation and condensation of water vapor at the surface. These physical processes result in modifications of the size and shape of snow crystals, and therefore of their SSA. Since SSA determines albedo, this variable contributes to daily albedo variations. Other critical variables that cause daily albedo variations include the solar zenith angle (Warren, 1982). Previous field studies have indeed observed daily variations in snow albedo in Antarctica, (McGuffie and Henderson-Sellers, 1985; Pirazzini, 2004)

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



but detailed interpretations in terms of quantitative snow surface properties could not be made, because among other factors snow SSA was not measured. On the Antarctic plateau, far from any source of impurities (Warren and Clarke, 1990), snow albedo can in theory be calculated using snow SSA only, in the case of a semi-infinite homogeneous layer. In the case of a complex stratified snowpack, the SSA, thickness and density of each layer must be known (Zhou et al., 2003).

During summer campaigns at Dome Concordia (DC, 75°06' S, 123°20' E, 3233 m a.s.l.) on the east Antarctic plateau, under calm and sunny weather, we frequently observed the rapid growth of snow crystals on the surface, in the absence of detectable precipitation. These crystals grow slowly and can persist for several weeks at a time. They usually disappear when they are blown off by strong winds. We initially thought that they were surface hoar crystals, which very commonly grow on any snow surface that cools radiatively in the absence of strong winds (Cabanès et al., 2002). To study the impact of these crystals on the radiative budget of the surface, we undertook to measure their SSA and density. From our initial results, we inferred that these crystals probably did not result only from surface hoar formation, and that sublimation crystals, as observed by (Gow, 1965), (Orheim, 1968) and (Weller, 1969), were probably forming. We therefore used the Crocus snow model (Vionnet et al., 2012) and the formalism of (Style and Worster, 2009) to test the possibility that sublimation crystals were forming on the Antarctic plateau. We report here these field and modeling studies and attempt to derive implications for the energy and mass balance of the Antarctic plateau.

2 Experimental and modeling methods

2.1 Experimental

The study took place at the French–Italian base of DC, on 18–19 January 2009. An area of about 16 m² was used for this study. A fairly flat and planar area was chosen to

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reduce the effects of surface roughness (i.e. sastrugi). SSA measurements were done using the DUFISSS instrument (DUAl Frequency Integrating Sphere for Snow SSA measurements) described in (Gallet et al., 2009). Briefly, the principle of the method is to measure snow hemispherical reflectance at 1310 or 1550 nm with an integrating sphere. The SSA-reflectance relationship was determined from simultaneous SSA measurements using CH₄ adsorption (Legagneux et al., 2002). The method can be used on all snow types and its accuracy is 10 % at the 95 % confidence level. In all cases, triplicate samples were measured, to minimize errors and the effects of spatial variability. To measure the SSA of surface snow with a 1 cm resolution, the sampling protocol of (Gallet et al., 2009) was modified. Briefly, snow was delicately sampled with a spatula and placed into the sample holder, a cylindrical cup 63 mm in diameter and 25 mm deep. Since in all cases the snow was soft and non-cohesive, this sampling did not result in structural modifications. The excess snow was shaved off with a spatula, so that the sample presented a perfectly flat surface. Other aspects of the protocol are as described in (Gallet et al., 2009). The sample holder was placed against the nadir port of the sphere, where it was illuminated with a 1310 nm laser diode. The reflected light was measured with an InGaAs photodiode. The signal was calibrated with six reflectance standards.

Measuring the density of 1 cm-thick snow layers is extremely delicate. Here, we weighed the sample holder used for SSA measurements. Great care was taken not to compact or otherwise perturb snow structure, which was possible because snow grains were only very loosely bonded. Evaluating a measurement error using this method is delicate, because we certainly introduce systematic errors. Typical uncertainty in snow density measurements made by weighing a known volume are about 10 % (Conger and McClung, 2009). Given the extra process of transferring snow into a container with a spatula, we estimate that the overall uncertainty in our density measurements was 20 %. Visual snow observations were performed using an 8X magnifying glass. Efforts were done to carefully identify the various crystal types found and their proportions.

Data for meteorological variables are available at DC. An IR sensor yielded snow surface temperature. Air temperature at 2 m and wind speed are also observed with an Automatic Weather Station (AWS). Errors on these measurements are caused, among other things, by radiative heating of the shelter housing the thermometer (Genthon et al., 2010) and inaccuracy in the reference black body of the IR sensor. Overall, we estimate that the accuracy is 1.5°C on the air and snow skin temperatures. We also used temperature, humidity and wind speed data at 4 m height from the 45 m height tower where sensors are ventilated as detailed (Genthon et al., 2010).

2.2 Modeling

In order to investigate the relative impact of various surface processes leading to condensation or sublimation of vapor water at the snow–air interface, we used the detailed snowpack model Crocus (Vionnet et al., 2012). Crocus is a multilayered one-dimensional snowpack model with an elaborate description of the energy budget of the snowpack including surface energy and mass balance. Each layer is characterized by its density, thickness, size and shape of grains, temperature and liquid water content. Snow metamorphism in Crocus is represented by semi-empirical variables and is simulated for each layer using the temperature, the temperature gradient and the liquid water content. The water vapor exchanges inside the snowpack, i.e. from one layer to the next, are not explicitly accounted for in this model. However, water vapor exchanges between the snow surface and the atmosphere are taken into account. Solar radiation penetration is computed as a function of snow optical radius and age, as a surrogate for snow darkening due to the deposition of aerosols, a process which is irrelevant at Dome C. Snow darkening was thus not accounted for in our simulation. Crocus takes as input the meteorological variables that constrain the surface mass and energy balance of the snowpack: air temperature, specific humidity, wind speed, shortwave and longwave incoming radiation, and precipitation amount and phase.

Brun et al. (2011) have carried out Crocus simulations at Dome C. In their study two types of simulations were done, i.e. stand alone simulations forced by local meteorolog-

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ical variables over a period of ten days and coupled simulations with the atmospheric model AROME. They showed that the model reproduces the observed downward propagation of the diurnal heat wave into the upper 50 cm of the snowpack and the snow surface temperature reasonably well.

5 In this study, we used the meteorological variables provided by ERA-interim reanalysis (Dee et al., 2011). These data are provided on a 3 h time step and have been processed as in (Brun et al., 2013). Operational analyses from the European Center for Medium-Range Weather Forecasts (ECMWF) similar to ERA-interim have been evaluated at Dome C with respect to field measurements of temperature, humidity and wind
10 speed at different levels on a 45 m tower by (Genthon et al., 2010). They concluded on a small warm bias of temperature relative to observations for all height. The amplitude of the diurnal cycle of relative humidity was also overestimated. Consequently, for January 2009, we used wind speed, humidity and temperature measured (ventilated sensors) 4 m above ground at the tower instead of ERA-interim reanalysis.

15 A first Crocus simulation was initialized with a 30 m thick snowpack with similar snow physical properties as in (Brun et al., 2011). This simulation was performed during 2 yr (1 January 2007 to 13th January 2009) to initialize properly the deep snowpack temperature profile using ERA-interim forcing data. A simulation from 13 to 20 January 2009 was then run where temperature, humidity and wind speed were taken from
20 measurements at 4 m height. In addition, the density of the upper 2 cm of the simulated snowpack was changed to the mean density values measured during the period, i.e., 141 kg m^{-3} measured for the first centimeter instead of 98 kg m^{-3} modeled and 236 kg m^{-3} for the second centimeter instead of 210 kg m^{-3} .

25 Crocus features neither a detailed treatment of condensation or sublimation of water vapor in terms of snow micro-structural properties, nor a representation of water vapor exchange between layers. Thus the purpose of the model run presented here is not an attempt to reproduce the daily variations in surface SSA but rather an exploration of the physical conditions leading to such variations.

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results

3.1 Experimental

Our January 2009 study took place under clear sky conditions. The surface was covered with crystals that closely resembled sublimation and surface hoar crystals as described in (Gow, 1965), (Orheim, 1968) and (Weller, 1969). Observations (Fig. 1) revealed clusters of crystals presenting both faceted and rounded parts. Sharp points, sometimes with dendrite-like outgrowths, were often sticking out of those clusters. Faceted crystals, that resembled typical surface hoar crystals, were often observed between and on those clusters. Faceted-rounded crystals, following the terminology of (Fierz et al., 2009), were also observed. A quantitative estimation of the proportion of such mixture is not very realistic, and we will explain the reasons in this work. During this study, the clusters covered the whole surface as far as the eye could see, and definitely were not limited to the area around the base.

SSA and density data are shown in Fig. 2a, and the numerical values are detailed in Table 1. The SSA of the first cm increases in the evening of the 18th (event I.1) and also at noon on the 19th (event I.2) respectively by 23 % and 13 % relative to the previous measurement. SSA decreases at midday on both days (events D.1 and D.2), respectively by 17 and 20 %. For the second cm, SSA remains almost constant on the 18th. On the 19th, a 30 % increase is first observed, followed by a 30 % decrease, and finally an increase in the very last part of the study. The observed variations are greater than the measurement uncertainty of 10 %.

The density of the first cm varies around 140 kg m^{-3} , while that of the second cm is around 240 kg m^{-3} . In general density and SSA appear anti-correlated. If we look in more detail at the density measurements between 18 and 19 January, we observe a decrease of 20 % and 30 %, respectively for the first and the second cm over the first 24 h of the study. This is greater or equal than our estimated accuracy of 20 %.

Figure 2b shows that the air temperature peaked around 16:00 and was minimal around 04:00 (dashed curve). During this study, the wind speed was low (green curve)

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and the sky remained clear, except for occasional thin cirrus covering less than 10% of the sky. Snow skin temperature (black curve) varied between -31°C at noon and -47°C between 02:00 and 03:00.

3.2 Simulated surface mass and energy balance

5 The snow surface temperature simulated by Crocus was evaluated with respect to the measurements of the IR sensor (Fig. 2b). The mean bias over the three days of simulation is 0.22 K, the root mean square deviation is 2.01 K, and most of the discrepancies happen at night. Given the potential errors on measuring temperature from IR emission, in part due to errors on snow emissivity (Bintanja and Vandenbroeke, 1995),
10 this shows that Crocus provides a reliable estimates of the snow surface temperature. A similar conclusion was reached by (Brun et al., 2011).

Figure 3 presents the modeled vertical profiles of temperature for the upper layers of the snowpack for different times of the day for 18 and 19 January 2009. The evolution of the vertical profile is similar for both days with maximum surface temperature around noon. During the day, the warmest point of the profile is located about 1 cm below the snow surface. This is a direct consequence of the coupled effect of solar radiation which penetrates inside the snowpack and radiative cooling under clear sky conditions that occurs at the surface (Colbeck, 1989). This induces a strong temperature gradient around noon in the uppermost part of the snowpack.

20 Atmospheric latent and sensible heat fluxes together with the net radiation budget at the snow surface are reported in Fig. 4 for the three days of simulations. Sensible and radiative fluxes are positive in the day time so that the atmosphere expectedly heats up during the day, as well as the snow surface due to a positive radiation budget. The latent heat flux also shows a diurnal cycle, with night time condensation alternating
25 with daytime sublimation. Although sublimation leads to mass loss, it can under certain conditions also lead to the formation of frost flowers over young sea-ice (Style and Worster, 2009; Domine et al., 2005; Perovich and Richter-Menge, 1994) or sublimation crystals over the snow surface (Gow, 1965; Weller, 1969), as observed here.

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To investigate the environmental conditions favorable for various types of ice crystals formation, we used the classification described by Style and Worster (2009) which includes six regimes of possible behavior of an air-ice system in terms of water vapor exchange (Fig. 3 of Style and Worster, 2009). These regimes depend on the saturation ratio of the atmosphere a few meters above the surface (a region called far field by Style and Worster, 2009), $\pi = p_{\infty}/p_{\text{sat}}(T_{\infty})$, where p_{∞} is the water vapor pressure of the far field and $p_{\text{sat}}(T_{\infty})$ is the saturation pressure at the far-field temperature, and of the difference between the far field air temperature, T_{∞} and the temperature of the surface, T_i . Note that at Dome C, the air temperature is always below freezing so that the saturation ratio is calculated with respect to solid ice. In our case, the atmosphere is always undersaturated ($p_{\infty}/p_{\text{sat}}(T_{\infty}) < 1$) at 4 m above the surface. Therefore, only four of the six regimes described by (Style and Worster, 2009) in their Table 2 need to be considered. The environmental conditions needed for each of these regimes can be described as a function of the value of $\mu = (T_{\infty} - T_i)/T_{\infty}$ and $m = ML/(RT_{\infty})$ where $ML = 50.9 \text{ kJ mol}^{-1}$ is the molar heat of sublimation of ice and $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ is the gas constant. These conditions are reported in Table 2.

Figure 5 depicts the typical snow and air temperature profiles during daytime for a clear sky day together with profiles of vapor pressure and saturation vapor pressure. As shown in this schematic diagram, a supersaturation region can exist just above the surface where the vapor pressure is greater than the saturation vapor pressure (Style and Worster, 2009). The local supersaturation in the air just above the ice happens because of the strong non linear relationship between temperature and saturation vapor pressure. This local supersaturation can induce water vapor fluxes responsible for the formation of sublimation crystals. This process has been frequently reported on sea ice, with the formation of large crystals called frost flowers (Perovich and Richter-Menge, 1994; Domine et al., 2005). The typical height of this supersaturation region shown in Fig. 5, a few mm at most, is not accessible to measurements currently carried out at Dome C. Consequently, to investigate the possibility of sublimation crystals formation, we use the criteria of Style and Worster (2009) and compare the saturation

ratio of the far field, taken in our study as the 4 m measurements, and the snow surface temperature simulated by Crocus.

Figure 6 depicts the saturation ratio ($\pi = p_{\infty}/p_{\text{sat}}(T_{\infty})$) vs. $(1 + m\mu)e^{-m\mu}$. The blue dots corresponds to the different times of the day for 17 January 2009, the red dots the same for 18 January, and the green diamonds for 19 January. The black solid line is the $y = x$ line which separates regimes V (grayed area) and VI (white area). Dots are only plotted if the surface is warmer than the air, which only occurs during daytime. Regime VI (Table 2) is characterized by unsaturated far field and surface temperature higher than far field temperature. This corresponds to the regime required for the formation of sublimation crystals (Style and Worster, 2009).

Figure 6 shows that regime VI occurs on the morning (8 to 12 h) of 18 January, when the air humidity is maximal and the snow is warmest. Our simulations therefore predict that sublimation crystals can indeed form, but only on the morning of 18 January, while we observed their formation also on 17 and 19 January. We explain this discrepancy as follows. Conditions for which sublimation crystals can form are very sensitive to the values of the temperature difference between the snow and the air and to the air relative humidity. In their work, (Style and Worster, 2009) investigated the possibility of frost flowers formation over ice. In this study, we observed the formation of sublimation crystals over snow, which is a porous material while ice is not. Consequently in our case, the water vapor can originate from below the snow surface. As seen on Figs. 3 and 5, during the day, the warmest point of the snowpack is located around 1 cm below the surface. This warm point facilitates the release of water vapor through ice sublimation and the formation of a supersaturation region just above the surface.

It may then be argued that not only snow surface temperature but also the temperature of the warmest point of the uppermost snowpack needs to be considered when assessing the conditions favorable for the formation of sublimation crystals. Style and Worster (2009) did not need to consider this because the ice surface they modeled is not porous and the water vapor source is indeed the ice–air interface. The temperatures of the snow surface and of the warmest point below the surface are determined,

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



among other factors, by the thermal conductivity of snow. In our modeling study, the thermal conductivity was computed as a function of density following Yen's parameterization (Yen, 1981). Although the parameterization has proven to be adequate for numerous snowpack conditions, density is not a perfect predictor of snow thermal conductivity, and snow mechanical properties also come into play, with harder snows being more conductive, for a given density (Domine et al., 2011a). Here, the surface snow at Dome C was very soft, likely implying a lower thermal conductivity than predicted by (Yen, 1981). This would generate a higher temperature below the snowpack surface. These considerations would imply that regime VI is met more often than predicted in Fig. 6 and would further increase the possibility of formation of sublimation crystals.

To illustrate these considerations, new simulations were thus run starting on 13 January with modified densities, using 100 kg m^{-3} for the first centimeter and 200 kg m^{-3} for the second centimeter which are lower than base run densities and lead to a lowered thermal conductivity. The Style and Worster' regimes met for the different hours of the day are depicted in Fig. 7. These calculations were done in two ways. First, as previously, the snow temperature used was the surface snow temperature (blue diamonds). Second, the snow temperature used was that 1 cm below the surface (snow warmest point, light blue square).

Figure 7 shows that condensation happens each night in all the cases (regime IV), confirming that surface hoar crystals do form. Regime VI (formation of sublimation crystals) also happens each day around noon when considering that the vapor exchanges are initiated at the warmest point of the snowpack and using a modified density lower than the field value. While using field measured density, regime VI still occurs on 17 and 18 January.

In summary, besides the commonly described formation of surface hoar crystals, our analysis predicts that sublimation crystals form every day of our study period, if the formalism of (Style and Worster, 2009) is adapted to the conditions applicable to a porous snow surface. Formation can take place under clear sky and low wind because it allows a maximum cooling at the snow surface and heating below that surface so that a high

tribute the remaining 11 % and are responsible for a variation of 0.007 in albedo. This is not negligible as it leads to an instantaneous local forcing of 6 W m^{-2} at 14:30. Between 18 January 14:30 and 19 January noon, the SSA increase due to the growth of sublimation crystals leads to an albedo increase of 0.012 and to a forcing of 10 W m^{-2} .

Using Crocus, we calculate that increasing the albedo by 0.012 over 24 h on 19 January leads to a maximum surface temperature drop of 0.45 K and an average drop over the day of 0.33 K. This result is similar to predicted by (Picard et al., 2012) on the effect of increased precipitation on the surface temperature of the Antarctic plateau, confirming that snow SSA is a critical factor in the energy budget of Antarctica.

Note that this calculation does not take into account the albedo change due to changes in surface roughness (Hudson and Warren, 2007). As seen in Fig. 1, crystals formation induces a change of surface roughness. The structures are very small, a few mm, and the effect of such small structures on albedo have never been quantified, even though it might reach a few percent (Warren et al., 1998). This should be also quantified in future work.

Our observations show significant exchanges of water vapor between the snow and the atmosphere. We have demonstrated that some of the crystals formed were sublimation crystals, while others were surface hoar crystals, showing that water vapor fluxes were important and in both directions, affecting the surface mass balance in a complex manner. Figure 4 shows that sublimation predominates over condensation, so that it may be deduced that water vapor exchanges lead to an overall mass loss. However, uncertainties in the calculations do not allow such a clear-cut conclusion. Indeed, this result is based on simulations by Crocus, which does not treat the formation of sublimation crystals and the resulting modifications in snow SSA and roughness. Therefore we cannot at this stage quantify with great accuracy the surface energy and mass budgets under the clear sky and low wind conditions of this study. Future work will be devoted to this task and in particular to measure the albedo and mass balance of the snow. For an accurate mass balance, the density of snow layers can be mea-

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

sured, but investigating the top 2 cm is likely not sufficient, as deeper layers may also contribute to sublimating fluxes.

It would also be interesting to quantify the fractions of depth hoar and sublimation crystals that formed. Looking at the crystal morphology probably is not sufficient for identification. At Alert, in the Canadian high Arctic, (Cabanès et al., 2002) observed feather-like crystals that were positively identified as surface hoar and that had many similarities with those of Fig. 1. Sublimation crystals are therefore difficult to differentiate from surface hoar, and indeed (Gow, 1965) seems to have mixed some sublimation crystals with surface hoar (which he calls hoar frost). From Fig. 4, most of the latent heat fluxes are indicative of sublimation, a conclusion similar to that of (Weller, 1969), from which we deduce that most of our crystal mass is probably from sublimation.

Furthermore since for surface hoar crystals, the matter comes from the atmosphere while from sublimation crystals it comes from the snow this may generate an isotopic signal (Neumann et al., 2008; Steen-Larsen et al., 2013) that might be usable to quantify water vapor exchanges and therefore improve our understanding of the processes involved in the mass balance of the East Antarctic ice sheet.

5 Conclusions

This study, performed under conditions of clear sky and low wind frequently encountered in summer on the Antarctic plateau, highlights the highly dynamic character of the snow surface of the Antarctic plateau under these conditions. We conclude that water vapor exchanges between the snow surface and the atmosphere take place in both directions. In the daytime, upward fluxes lead to the growth of sublimation crystals that form on the surface.

This dynamic interface has many important consequences for the mass and energy budgets of the Antarctic plateau. The growth of depth hoar and sublimation crystals dramatically changes the snow structure and surface roughness, affecting turbulent fluxes and hence energy and mass exchanges. Furthermore, the formations of crys-

TCD

7, 5971–5999, 2013

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tals modify snow SSA, density and again roughness, and therefore albedo. We have calculated that the albedo change was up to 0.012, which represents a local forcing of -10 W m^{-2} at noon.

Acknowledgements. This work was funded by the French Polar Institute (IPEV) under grant Nite DC, n°1011. We thank G. Picard for discussions on energy and mass exchanges at the surface. We are grateful to L. Arnaud for providing us with the snow surface temperature data and the BSRN network for the radiation data. We thank all the technical staff of Concordia station for their help during the campaign. C. Genthon kindly provided data at 4 m from the tower. JCG was also supported through the project “Long range transport of Black Carbon and its effect on snow albedo in North East China and in the Arctic” funded by the Norwegian Research council (Norklima program, project 193717).

References

- Aoki, T., Fukabori, M., Hachikubo, A., Tachibana, Y., and Nishio, F.: Effects of snow physical parameters on spectral albedo and bidirectional reflectance of snow surface, *J. Geophys. Res.*, 105, 10219–10236, 2000.
- Bintanja, R. and Vandenbroeke, M. R.: The surface-energy balance of antarctic snow and blue ice, *J. Appl. Meteorol.*, 34, 902–926, doi:10.1175/1520-0450(1995)034<0902:tseboa>2.0.co;2, 1995.
- Brun, E., Six, D., Picard, G., Vionnet, V., Arnaud, L., Bazile, E., Boone, A., Bouchard, A., Genthon, C., Guidard, V., Le Moigne, P., Rabier, F., and Seity, Y.: Snow/atmosphere coupled simulation at Dome C, Antarctica, *J. Glaciol.*, 57, 721–736, 2011.
- Brun, E., Vionnet, V., Boone, A., Decharme, B., Peings, Y., Valette, R., Karbou, F., and Morin, S.: Simulation of northern Eurasian local snow depth, mass and density using a detailed snowpack model and meteorological reanalysis, *J. Hydrometeorol.*, 14, 203–214, doi:10.1175/jhm-d-12-012.1, 2013.
- Cabanes, A., Legagneux, L., and Domine, F.: Evolution of the specific surface area and of crystal morphology of Arctic fresh snow during the ALERT 2000 campaign, *Atmos. Environ.*, 36, 2767–2777, 2002.
- Colbeck, S. C.: An overview of seasonal snow metamorphism, *Rev. Geophys.*, 20, 45–61, 1982.

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Colbeck, S. C.: Snow-crystal growth with varying surface temperatures and radiation penetration, *J. Glaciol.*, 35, 23–29, 1989.
- Conger, S. M. and McClung, D. M.: Comparison of density cutters for snow profile observations, *J. Glaciol.*, 55, 163–169, 2009.
- 5 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kallberg, P., Kohler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011.
- 10 Domine, F., Taillandier, A. S., Simpson, W. R., and Severin, K.: Specific surface area, density and microstructure of frost flowers, *Geophys. Res. Lett.*, 32, L13502, doi:10.1029/2005gl023245, 2005.
- 15 Domine, F., Taillandier, A. S., and Simpson, W. R.: A parameterization of the specific surface area of seasonal snow for field use and for models of snowpack evolution, *J. Geophys. Res.*, 112, F02031, doi:10.1029/2006jf000512, 2007.
- 20 Domine, F., Bock, J., Morin, S., and Giraud, G.: Linking the effective thermal conductivity of snow to its shear strength and its density, *J. Geophys. Res.*, 116, F04027, doi:10.1029/2011JF002000, 2011a.
- Domine, F., Gallet, J. C., Barret, M., Houdier, S., Voisin, D., Douglas, T., Blum, J. D., Beine, H., and Anastasio, C.: The specific surface area and chemical composition of diamond dust near Barrow, Alaska, *J. Geophys. Res.*, 116, D00R06, doi:10.1029/2011JD016162, 2011b.
- 25 Fierz, C., Armstrong, R. L., Durand, Y., Etchevers, P., Greene, E., McClung, D. M., Nishimura, K., Satyawali, P. K., and Sokratov, S. A.: The International classification for seasonal snow on the ground UNESCO-IHP, Paris/ACS Contribution N°1, 80, 2009.
- Flanner, M. G. and Zender, C. S.: Linking snowpack microphysics and albedo evolution, *J. Geophys. Res.*, 111, D12208, doi:10.1029/2005jd006834, 2006.
- 30 Gallet, J.-C., Domine, F., Zender, C. S., and Picard, G.: Measurement of the specific surface area of snow using infrared reflectance in an integrating sphere at 1310 and 1550 nm, *The Cryosphere*, 3, 167–182, doi:10.5194/tc-3-167-2009, 2009.

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gallet, J.-C., Domine, F., Arnaud, L., Picard, G., and Savarino, J.: Vertical profile of the specific surface area and density of the snow at Dome C and on a transect to Dumont D'Urville, Antarctica – albedo calculations and comparison to remote sensing products, *The Cryosphere*, 5, 631–649, doi:10.5194/tc-5-631-2011, 2011.

5 Genthon, C., Town, M. S., Six, D., Favier, V., Argentini, S., and Pellegrini, A.: Meteorological atmospheric boundary layer measurements and ECMWF analyses during summer at Dome C, Antarctica, *J. Geophys. Res.-Atmos.*, 115, D05104, doi:10.1029/2009jd012741, 2010.

Gow, A. J.: On the accumulation and seasonal stratification of snow at the South Pole, *J. Glaciol.*, 5, 467–477, 1965.

10 Grenfell, T. C. and Warren, S. G.: Representation of a nonspherical ice particle by a collection of independent spheres for scattering and absorption of radiation, *J. Geophys. Res.*, 104, 31697–31709, 1999.

Hudson, S. R. and Warren, S. G.: An explanation for the effect of clouds over snow on the top-of-atmosphere bidirectional reflectance, *J. Geophys. Res.*, 112, doi:10.1029/2007jd008541, 2007.

15 Legagneux, L., Cabanes, A., and Domine, F.: Measurement of the specific surface area of 176 snow samples using methane adsorption at 77 K, *J. Geophys. Res.*, 107, 4335, doi:10.1029/2001jd001016, 2002.

McGuffie, K. and Henderson-Sellers, A.: The diurnal hysteresis of snow albedo, *J. Glaciol.*, 31, 188–189, 1985.

20 Neumann, T. A., Albert, M. R., Lomonaco, R., Engel, C., Courville, Z., and Perron, F.: Experimental determination of snow sublimation rate and stable-isotopic exchange, *Ann. Glaciol.*, 49, 2008, 49, 1–6, 2008.

Orheim, O.: Surface snow metamorphosis on the Antarctic plateau, in: *Norsk Polarinstitut Arbok 1966*, Norsk Polarinstitut, Oslo, 84–91, 1968.

25 Perovich, D. K. and Richter-Menge, J. A.: Surface characteristics of lead ice, *J. Geophys. Res.*, 99, 16341–16350, 1994.

Picard, G., Domine, F., Krinner, G., Arnaud, L., and Lefebvre, E.: Inhibition of the positive snow-albedo feedback by precipitation in interior Antarctica, *Nature Climate Change*, 2, 795–798, doi:10.1038/nclimate1590, 2012.

30 Pirazzini, R.: Surface albedo measurements over Antarctic sites in summer, *J. Geophys. Res.*, 109, doi:10.1029/2004jd004617, 2004.

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ricchiuzzi, P., Yang, S. R., Gautier, C., and Sowle, D.: SBDART: A research and teaching software tool for plane-parallel radiative transfer in the Earth's atmosphere, *B. Am. Meteorol. Soc.*, 79, 2101–2114, 1998.

Robinson, D. A., Dewey, K. F., and Heim, R. R.: Global snow cover monitoring – an update, *B. Am. Meteorol. Soc.*, 74, 1689–1696, 1993.

Stamnes, K., Tsay, S. C., Wiscombe, W., and Jayaweera, K.: Numerically stable algorithm for discrete-ordinate-method radiative-transfer in multiple-scattering and emitting layered media, *Appl. Opt.*, 27, 2502–2509, 1988.

Steen-Larsen, H. C., Masson-Delmotte, V., Hirabayashi, M., Winkler, R., Satow, K., Prié, F., Bayou, N., Brun, E., Cuffey, K. M., Dahl-Jensen, D., Dumont, M., Guillevic, M., Kipfstuhl, J., Landais, A., Popp, T., Risi, C., Steffen, K., Stenni, B., and Sveinbjörnsdóttir, A.: What controls the isotopic composition of Greenland surface snow?, *Clim. Past Discuss.*, 9, 6035–6076, doi:10.5194/cpd-9-6035-2013, 2013.

Style, R. W. and Worster, M. G.: Frost flower formation on sea ice and lake ice, *Geophys. Res. Lett.*, 36, L11501, doi:10.1029/2009gl037304, 2009.

Taillandier, A. S., Domine, F., Simpson, W. R., Sturm, M., and Douglas, T. A.: Rate of decrease of the specific surface area of dry snow: Isothermal and temperature gradient conditions, *J. Geophys. Res.*, 112, F03003, doi:10.1029/2006jf000514, 2007.

Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geosci. Model Dev.*, 5, 773–791, doi:10.5194/gmd-5-773-2012, 2012.

Warren, S. G.: Optical-properties of snow, *Rev. Geophys.*, 20, 67–89, 1982.

Warren, S. G. and Clarke, A. D.: Soot in the atmosphere and snow surface of Antarctica, *J. Geophys. Res.*, 95, 1811–1816, 1990.

Warren, S. G., Brandt, R. E., and Hinton, P. O.: Effect of surface roughness on bidirectional reflectance of Antarctic snow, *J. Geophys. Res.-Planets*, 103, 25789–25807, 1998.

Weller, G.: The heat and mass balance of snow dunes on the central Antarctic plateau, *J. Glaciol.*, 8, 277–284, 1969.

Yen, Y.-C.: Review of thermal properties of snow, ice, and sea ice, United States Army Corps of Engineers, Hanover, NH, USACRREL Report 81–10, 1–27, 1981.

Zhou, X. B., Li, S. S., and Stamnes, K.: Effects of vertical inhomogeneity on snow spectral albedo and its implication for optical remote sensing of snow, *J. Geophys. Res.*, 108, D4738, doi:10.1029/2003jd003859, 2003.

Table 1. SSA and density values used in DISORT for albedo calculations for the 18–19 January. Standard deviations are calculated from triplicates measurements, for density and SSA.

First centimeter				
Date and Time (GMT + 08:00)	Mean SSA $\text{m}^2 \text{kg}^{-1}$	SSA standard deviation	Mean density kg m^{-3}	Density standard deviation
18 Jan 2009 11:00	40.0	1.2	156	8
18 Jan 2009 14:30	33.3	0.5	165	4
19 Jan 2009 00:00	40.9	1.8	147	8
19 Jan 2009 04:30	39.7	2.3	145	10
19 Jan 2009 11:30	46.3	1.0	125	5
19 Jan 2009 17:10	36.8	0.7	130	6
19 Jan 2009 21:00	37.0	1.1	133	4
20 Jan 2009 01:20	37.8	0.9	133	4

Second centimeter				
Date and Time (GMT + 08:00)	Mean SSA $\text{m}^2 \text{kg}^{-1}$	SSA standard deviation	Mean density kg m^{-3}	Density standard deviation
18 Jan 2009 11:00	25.6	1.2	290	6
18 Jan 2009 14:30	25.4	0.7	258	3
19 Jan 2009 00:00	24.7	0.6	235	16
19 Jan 2009 04:30	26.5	0.9	236	13
19 Jan 2009 11:30	32.1	1.1	202	13
19 Jan 2009 17:10	23.2	0.7	215	7
19 Jan 2009 21:00	25.1	0.3	227	10
20 Jan 2009 01:20	28.3	0.6	231	11

**The growth of
sublimation crystals
and surface hoar**

J.-C. Gallet et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Table 2. Possible regimes for water vapor exchange as a function of environmental conditions for a unsaturated atmosphere. This table is adapted from the Table 1 of Style and Worster (2009).

Regime	Temperature conditions	Surface characteristics	Environmental conditions
III	Air warmer than surface	Condensation Supersaturated	$T_{\infty} > T_i$ and $\pi > (1 + m\mu)e^{-m\mu}$
IV	Air warmer than surface	Condensation Unsaturated	$T_{\infty} > T_i$ and $\pi < (1 + m\mu)e^{-m\mu}$ and $\pi > e^{-m\mu}$
V	Surface warmer than air	Evaporation Unsaturated	$T_{\infty} < T_i$ and $\pi < e^{-m\mu}$ or $T_{\infty} > T_i$ and $\pi < (1 + m\mu)e^{-m\mu}$
VI	Surface warmer than air	Evaporation Supersaturated	$T_{\infty} < T_i$ and $\pi > (1 + m\mu)e^{-m\mu}$

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




Fig. 1. Sublimation crystals and surface hoar crystals on the surface of the snowpack at Concordia station on 18 January 2009.

TCD

7, 5971–5999, 2013

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.




Fig. 2. Time series of physical variables on 18 and 19 January 2009. Time is local time, GMT + 08:00. **(a)** Snow SSA and density values. **(b)** Meteorological variables.

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.




Fig. 3. Simulated vertical temperature profiles for the upper layer of the snowpack on 18 June 2009 (a) and on 19 June 2009 (b) for several local times (LT = UTC + 8).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.




Fig. 4. Simulated energy fluxes at the air–snow interface: atmospheric latent and sensible heat on top panel; radiative fluxes at snow surface on bottom panel. Positive atmospheric latent heat implies snow sublimation; positive sensible heat leads to transfer of energy from the snow to the atmosphere. Positive net radiation would be leading to heating of the snow surface.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 5. Schematic diagram of temperature and water vapour pressure for the air-snow system. The figure is adapted from Fig. 2 of Style and Worster (2009). The snow temperature is simulated by Crocus on 18 January at 11:00 LT. The blue and red dots and the blue diamond are calculated from 4 m measurements. The T , water vapour pressure and saturation pressure profiles between the snow surface and 4 m height are drawn from Fig. 2 of Style and Worster (2009).

The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.




Fig. 6. Atmosphere saturation ratio as a function of $(1 + m\mu)e^{-m\mu}$. The different markers correspond to the three days of the measurements period. Dots are only indicated if the surface is warmer than the far field. The white area corresponds to regime VI (Table 2) and the greyed area to regime V. Hours are local time.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.




Fig. 7. Determinations of the regimes of Style and Worster (2009) with modified density values and taking the snow temperature 1 cm below the surface. The blue diamonds correspond to a simulation with modified density and the snow temperature is taken as the surface temperature. The red crosses are also for modified density but the snow temperature is taken 1 cm below the surface. The green crosses are the results of simulations with measured density and snow temperature 1 cm below the surface.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


The growth of sublimation crystals and surface hoar

J.-C. Gallet et al.




Fig. 8. Broadband albedo, 18–19 January 2009 for local time (GMT + 08:00). Calculations were done using DISORT with two scenarios: SSA variations are always taken into account while the effect of the solar zenith angle (SZA) is either taken into account or not.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)