

Seasonal changes of
ice surface
characteristics

D. M. Chandler et al.

This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

Seasonal changes of ice surface characteristics and productivity in the ablation zone of the Greenland Ice Sheet

D. M. Chandler¹, J. D. Alcock², J. L. Wadham¹, S. L. Mackie³, and J. Telling¹

¹Bristol Glaciology Centre, School of Geographical Sciences, Bristol University, Bristol, UK

²4 Avenham Terrace, Preston, Lancashire, PR1 3SY, UK

³School of Earth Sciences, Bristol University, Bristol, UK

Received: 3 February 2014 – Accepted: 16 February 2014 – Published: 25 February 2014

Correspondence to: D. M. Chandler (davemchandler@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Field and remote sensing observations in the ablation zone of the Greenland Ice Sheet have revealed a diverse range of ice surface characteristics, primarily reflecting the variable distribution of fine debris (cryoconite). This debris reduces the surface albedo and is therefore an important control on melt rates and ice sheet mass balance. Meanwhile, studies of ice sheet surface biological processes have found active microbial communities associated with the cryoconite debris, which may themselves modify the cryoconite distribution. Due to the considerable difficulties involved with collecting ground-based observations of the ice surface, our knowledge of the physical and biological surface processes, and their links, remains very limited. Here we present data collected at a field camp established in the ice sheet ablation zone at 67° N, occupied for almost the entire melt season (26 May–10 August 2012), with the aim of gaining a much more detailed understanding of the physical and biological processes occurring on the ice surface. These data sets include quadrat surveys of surface type, measurements of ice surface ablation, and in-situ biological oxygen demand incubations to quantify microbial activity. In addition, albedo at the site was retrieved from AVHRR remote sensing data. Observations of the areal coverage of different surface types revealed a rapid change from complete snow cover to the “summer” ice surface of patchy debris (“dirty ice”) and cryoconite holes. There was significant correlation between surface albedo, cryoconite hole coverage and cryoconite hole productivity during the main melt season, but microbial activity in “dirty ice” was not correlated with albedo and varied widely throughout the season. While this link suggests the potential for a remote-sensing approach to monitoring cryoconite hole biological processes, very wide seasonal and spatial variability in net surface productivity demonstrates the need for caution when extrapolating point measurements of biological processes to larger temporal or spatial scales.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

The ablation zone of the Greenland Ice Sheet displays a remarkable variety of surface characteristics that vary widely in time and space. The varying distribution of ice, snow, melt water and sediment across the surface strongly influences the surface energy balance, and therefore the melt rate and mass balance (e.g. Konzelmann and Braithwaite, 1995; Knap and Oerlemans, 1996; Bøggild et al., 2010). Recent studies have also shown links between sediment distribution and biological activity, indicative of feedback between microbial processes and albedo (e.g. Anesio et al., 2009; Uetake et al., 2010).

Due to the logistical demands of observing the ice sheet surface on foot, which is particularly problematic in the ablation zone where rivers, slush and crevasses make surface travel difficult, much of our present knowledge of its surface properties and processes has been gained through short 1–2 day field visits, automated data acquisition (primarily at automatic weather stations: AWSs) and satellite remote sensing. Consequently there are very few data sets on seasonal changes in surface characteristics and none on seasonal changes in biological activity on the ice sheet. Many more observations are needed to accurately estimate the contribution of observed biological activity to global chemical cycles, ideally leading to a method by which such activity can be monitored remotely.

In 2012 a field camp was established centrally in the ice sheet ablation zone at 1030 m elevation, 35 km from the western margin, with the aim of acquiring a continuous 10 week time series of data relating to surface characteristics and processes. The camp was occupied continuously from 26 May to 10 August, covering most of the melt season. Resulting data on surface ablation, surface type and surface productivity (as quantified by biological oxygen demand (BOD) experiments) are compared with a satellite remote-sensing time series of surface albedo for the same location. Using these data sets, we identify the characteristics of physical and biological surface processes at the field site and assess the extent to which these can be quantified remotely

TCO

8, 1337–1382, 2014

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in surface albedo retrievals. We finally examine in more detail the seasonal changes in surface productivity, which are presently unknown in Greenland. To our knowledge, this is the longest record of ground-based surface observations yet to be collected on the ice sheet.

1.1 Ablation zone ice surface characteristics

During the melt season, the ice sheet surface in the ablation zone displays a wide range of features commonly found on terrestrial glaciers, including crevasses, rivers, lakes, cryoconite holes and snow remnants. A variable amount of fine surface debris is also present, for example as dispersed particles, as a thin layer in cryoconite holes, or as thick deposits (Fig. 1). This fine debris, usually referred to as cryoconite, is mostly composed of fine mineral particles with a small (< 10%) organic matter content (Bayley, 1891; Hodson, 2010; Wientjes et al., 2011). Typical particle sizes are of order 10^{-5} to 10^{-4} m (data from several studies reported in Table 1 of Hodson et al., 2010). During the summer, debris causes some parts of the ablation zone to appear notably darker than the surrounding ice in satellite images (Wientjes et al., 2011). These dark zones can be clearly observed each year during the melt season in true-colour MODIS AERONET imagery of west Greenland (<http://lance-modis.eosdis.nasa.gov/imagery/subsets/?project=aeronet&subset=Kangerlussuaq>). The lower albedo of the dark region in the ablation zone has previously been interpreted as arising from the accumulation of surface melt water (Oerlemans and Vugts, 1993; Knap and Oerlemans, 1996). Whether debris- or water-related, this zone of lower albedo has a direct impact on surface melt rates and therefore on ice sheet mass balance.

As a consequence of the variable surface characteristics and debris cover, the ice surface albedo can range widely, from > 80% (fresh snow) to typically 40–60% (bare ice) and as low as 10% for cryoconite debris (Knap and Oerlemans, 1996; Bøggild et al., 2010). However, bare ice containing cryoconite holes has been found to have a higher albedo than bare ice with a lower mass loading of dispersed debris, because

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the depth of the cryoconite holes keeps the debris partially hidden at most incidence angles (Bøggild et al., 2010). Seasonal changes in albedo have been identified in the ablation zone, where a sharp then gradual decrease in albedo in the early melt season is associated with the melting of highly-reflective seasonal snow and the subsequent increase in surface water and/or exposure of surface debris (Knap and Oerlemans, 1996; Stroeve et al., 2001; Wientjes et al., 2011). Although none of these studies has presented detailed time series lasting longer than a year, consistency between the different years and locations investigated suggests the reported seasonal patterns are widespread and persistent.

In addition to the different ice types and patchy debris cover, the ice surface is also characterised by its small-scale topography. For example, the variable action of surface melt leads to the formation of the “lumpy” topography in the lower ablation zone, with peak-to-peak height differences between valleys and ridges increasing down-glacier and reaching 5 m close to the margin in west Greenland (Fig. 2). This topography includes the characteristic “domes”, “hummocks” and “gullies” described by Smeets and van den Broeke (2008) towards the end of the melt season at 67° N. Development of these topographical features and the change from snow to bare ice lead to large spatial and seasonal changes in the aerodynamic roughness length of several orders of magnitude. This affects the sensible and latent heat fluxes into the ice but has been rarely measured in the ablation zone (Duynderke and van den Broeke, 1994; Smeets and van den Broeke, 2008). Remote sensing of surface roughness has been shown to be possible using a Multi-angle Imaging SpectroRadiometer (MISR) in the Sermeq Kujalleq (Jacobshavn) region of west Greenland (Nolin et al., 2002; Nolin and Payne, 2007) but seasonal changes in roughness measured using this technique have not yet been reported. Roughness is not considered further in the present paper but should be addressed in future studies.

1.2 Ablation zone ice surface biological processes

The lack of any obvious plant or animal life on the ice sheet gives the impression that the surface is devoid of biological activity. However, studies of surface debris on the Greenland Ice Sheet have revealed the existence of active microbial communities including bacteria, cyanobacteria and green algae (Steinböck, 1936; Hodson et al., 2010; Stibal et al., 2012). Much of the activity was originally believed to be associated with the fine debris concentrated in a thin layer on the bottom of cryoconite holes (Anesio et al., 2009), but activity in algae present over much of the remaining surface in the ablation zone has recently been reported (Cook et al., 2012).

Cryoconite holes are widespread in the ablation zone of the Greenland Ice Sheet as well as in many other glaciated areas worldwide, and may form an important component of regional chemical cycles (e.g., Hodson et al., 2007; Anesio et al., 2009; Cook et al., 2012; Telling et al., 2012b). Since the Greenland Ice Sheet's melt zone has (potentially) a much greater surface area of microbial communities than that of any other terrestrial ice mass outside Antarctica, the role of the ice sheet's surface debris in the global carbon cycle has attracted increasing attention. Measurements of ecosystem productivity by in-situ incubation experiments (Table 1) have revealed active communities in the few field sites visited on the ice sheet to-date. Incubation experiments have now seen widespread use on many other glaciers, where similarly high levels of biological activity have been observed (Telling et al., 2010). The most detailed study in Greenland collected productivity measurements along a 70 km transect across the ablation zone in west Greenland at 67° N (Stibal et al., 2012; Cook et al., 2012), close to the field site studied in the present paper. Stibal et al. (2012) found a peak in cryoconite hole carbon storage rate ($22 \pm 4.8 \mu\text{gC day}^{-1}$ per unit mass of debris) at 11 km from the margin (761 m a.s.l.) and subsequent decrease to near zero by 65 km (1446 m a.s.l.). The most active part of the transect in terms of carbon storage rate per unit area was $\sim 7 \text{ mgC m}^{-2} \text{ day}^{-1}$ at the ice margin (399 m a.s.l.) where debris cover is very high (Stibal et al., 2012). Interestingly, measurements of surface algae productivity

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on the debris-poor ice between holes suggested the carbon storage by surface algae might be considerably higher than the above figures for cryoconite holes (Yallop et al., 2012), but the different method (^{14}C uptake; 1 h incubation rather than 24 h) introduced considerable uncertainty in the estimates. Attempts have been made to calculate ice sheet-scale carbon fluxes from Greenland's cryoconite holes and surface algae by extrapolating existing data (Hodson, 2010; Cook, 2012); however, the great variability in carbon fluxes measured at different field sites (Table 1) and a lack of knowledge regarding seasonal variability requires many more measurements to be made before realistic extrapolation is possible.

In addition to its biogeochemical interest, biological activity in cryoconite debris can modify the characteristics of the debris itself, for example by generating dark organic material or by causing small grains to clump together into larger, less mobile particles. These processes alter the albedo of the cryoconite holes (Langford et al., 2010; Takeuchi, 2010) and are therefore of great interest because of their potential effect on surface albedo, one of the key factors controlling ice sheet mass balance. We speculate that microbial processes will also affect the mobility of debris within the surface drainage system, and although this question remains open at present, it could be addressed using the time-lapse technique employed recently to observe cryoconite particle motion in Svalbard (Irvine-Fynn et al., 2011).

Given the importance of ice surface characteristics in controlling surface melt (and thus mass balance), and the increasing evidence for links between biological activity and surface characteristics, there is clearly a need to gain a better understanding of ice sheet surface processes in the ablation zone. In this paper, changes in the areal coverage and associated biological activity over most of a melt season are described for different surface types identified in the ablation zone of the western Greenland Ice Sheet at 67°N . The resulting time series reveal the seasonal patterns in much greater detail than has been achieved in previous field studies.

2 Methods

Field data were collected during the 2012 summer melt season at 66.97° N, 049.27° W, in the ablation zone of the ice sheet at 1030 m altitude and approximately 35 km from the western margin (Fig. 3). The field site is on the western edge of the “dark” band of the ablation zone identified by Wientjes et al. (2011). Site locations for the surveys and incubation experiments (see below) were all located down-stream from, and on the opposite side of, a surface melt water stream passing adjacent to the field camp, to minimise any interference or contamination from human activity.

2.1 Surface characteristics and ablation

Daily ice ablation was recorded by measuring the distance between the ice surface and the tops of five 40 cm plastic poles installed in holes drilled 50–100 cm into the ice. Installing the stakes into the ice below the surface avoided the lateral melting that normally occurs around poles left protruding from the ice. Stakes were arranged in a “+” shape with 4 outer stakes and 1 in the centre, at ~ 2 m spacing. The ice surface was often very rough, so for consistency each measurement was taken to be the vertical distance between the stake top and a 25 cm hacksaw blade laid centrally across the top of the hole in an east–west orientation. The initial surface slope at the site was flat (less than 10°), but by August the site had become a lump with the central and northern stakes on its flat top and the eastern, southern and western stakes on sides sloping outwards at 10–20°. Stakes were re-drilled before the stake had started to protrude from the hole.

Surface characteristics were monitored frequently from 1 June to 11 August (total 18 surveys) in five 2.7 m × 2.7 m quadrats situated within a region of roughly 60 m × 40 m (Fig. 4a and d). Each quadrat comprised a 10 × 10 grid with 30 cm spacing. Quadrat locations were marked by drilling ~ 50 cm deep holes in the ice at each corner of a 3 m × 3 m square, to ensure the same area of ice was surveyed each time. To carry out a survey, two nylon ropes (both marked with the positions of the hole edges and then

TCD

8, 1337–1382, 2014

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



at 30 cm intervals) were laid out in parallel between each pair of holes. The ropes were kept tight by attaching steel weights that could hang freely in the drilled holes. A rigid 3 m pole, also marked at 30 cm intervals, was laid between the first pair of markers on the ropes. The surface type was recorded immediately in front of each mark along the pole, before moving the pole to the next pair of rope markers and recording the second row of surface types, and so on. There were 6 surface types present at the field site, of which five were recorded in the quadrats (Table 2; Fig. 4). All surveys were carried out by the same pair of observers, to ensure consistency in the classification.

2.2 Productivity (biological oxygen demand)

Net ecosystem production (NEP) and respiration were quantified within each of the different surface environments recorded in the quadrat surveys, at approximately weekly intervals, by measuring the rate of O₂ production in samples collected from the ice surface. The method is described in detail by Telling et al. (2010). One sample was collected for each surface type present, and divided into 6 glass 250 mL biological oxygen demand (BOD) bottles. Ice samples were melted before pouring into the BOD bottles, by collecting the ice in a Whirlpak bag and immersing the bag in luke-warm water. These melting samples were left in a black plastic box, typically for 2–3 h, until completely melted (partially melted samples were not used). For samples representing cryoconite holes, debris from the base of the hole was collected using a 100 mL syringe fitted with a ~20 cm plastic tube, and the area of debris removed from the hole was selected to match the area of the BOD bottle base (Fig. 5a). This ensured that the debris thickness in the bottle was representative of that in the hole. These bottles were then filled with water collected from the same hole as the debris. The debris typically settled out onto the base of the bottle within 30 s.

The temperature and dissolved oxygen content of the water in each BOD bottle were measured using a PreSens Fibox3 fibreoptic oxygen meter (manufacturer's stated accuracy: ±1 %; PreSens, 2006), and the bottle was then immediately sealed with a glass stopper. Bottles were checked carefully to ensure no air bubbles were present.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Three out of the six bottles for each sample type were wrapped in foil to eliminate light (therefore measuring respiration only); the other three remained unwrapped to allow photosynthesis as well as respiration (to measure NEP). All bottles were placed in a cryoconite hole sufficiently deep to ensure complete immersion, in order to replicate typical light and temperature conditions as well as to prevent the bottle freezing and cracking. After 24 ± 1 h, the bottles were recovered and the dissolved oxygen content and temperature were re-measured.

In addition to these “standard” weekly incubation experiments, we carried out 4 additional experiments to better characterise temporal and spatial variability and errors in NEP rates, as follows. (1) To better quantify the variance in the four common environments (water, clean ice, dirty ice, cryoconite hole) we carried out additional incubations with 12 light and 12 dark replicates of just one sample type. (2) Temporal variability in cryoconite hole productivity within the 24 h incubation time was measured from 4–5 August by filling 32 bottles (16 light, 16 dark) with cryoconite hole debris and water, then recovering sets of bottles at 3 h intervals for 24 h. (3) Spatial variability and effects of cryoconite hole size were investigated on 6–7 August by choosing 5 sites and incubating light/dark pairs of samples from 3 holes of varying size (nominally “small”, “medium” and “large”) at each site. (4) Two standard incubations (4, 9 July) also included samples of orange-green material (Fig. 5b) sampled from the former bed of a shrinking lake (66.97° N, 049.21° W; 3 km east of the main field site).

The rate of oxygen production in the BOD bottles, expressed in $\text{mgL}^{-1} \text{h}^{-1}$, is equal to the difference between the start and end point dissolved oxygen concentrations, divided by the incubation time. In this experiment, two rates of oxygen production were measured for each sample type: $\Delta\text{O}_2^{(\text{light})}/\Delta t$ and $\Delta\text{O}_2^{(\text{dark})}/\Delta t$ for the light and dark bottles, respectively. We assume that respiration rates were independent of light conditions and therefore the same in the light and dark bottles. The rate of oxygen production by photosynthesis in the light bottles, representing that in the original environment, is therefore $P = (\Delta\text{O}_2^{(\text{light})} - \Delta\text{O}_2^{(\text{dark})})/\Delta t$.

To convert oxygen production into carbon storage, we assume photosynthesis can be represented as:



5 Respiration is taken to be the reverse reaction. In photosynthesis, this equation predicts that n moles of CO_2 are stored for each n moles of O_2 released. This assumption of a 1 : 1 stoichiometric ratio has been adopted in previous studies of cryoconite holes (Hodson et al., 2007, 2010; Anesio et al., 2009) but may not be exact: for example in Svalbard, Telling et al. (2010) found 1.24 ± 0.20 moles of O_2 were released for each
10 mole of CO_2 consumed during photosynthesis, while 0.80 ± 0.17 moles of CO_2 were released for each mole of O_2 consumed during respiration. Neglecting this uncertainty given the error bounds on the above ratios, a 1 : 1 molar ratio will yield 1.375 g of CO_2 (or 0.375 g of C) consumed per 1 g of O_2 released.

Carbon storage rates in the BOD bottles are more useful after conversion to a surface flux (e.g., $\text{mgCm}^{-2}\text{day}^{-1}$) for each sample type. To achieve this we required the
15 bottle volume V_B , bottle radius R_B , sample thickness (for ice samples) H_S , incubation duration T , water density ρ_W and sample density ρ_S . For water and cryoconite hole samples (CW, CS), the area of ice sheet A_S sampled to fill the bottle is $A_S = \pi R_B^2$. For ice samples, $A_S = V_B \rho_W / H_S \rho_S$. The C flux $F_{C,X}$ from each ice type X is then
20 $F_{C,X} = \Delta\text{O}_2^{(\text{light})} \times 0.375 V_B / (A_S T)$ and the total carbon flux into the surface at the field site is the sum of $A_X F_{C,X}$ over all the ice types X , weighted for their respective fractional areas A_X as determined in the quadrat surveys. Parameter values were as follows: $V_B = 250\text{mL}$, $R_B = 32\text{mm}$, $H_S \approx 30\text{mm}$, $\rho_S = 900\text{kgm}^{-3}$ (ice), $\rho_S = 300\text{kgm}^{-3}$ (snow), $\rho_W = 1000\text{kgm}^{-3}$. Quadrat surveys and BOD experiments did not always coincide, so
25 both data sets were linearly interpolated to daily resolution to allow calculation of daily values for the product $A_X F_{C,X}$ and for the net C flux.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2.3 Surface albedo

A surface broadband albedo product (AVHRR-SAL, CM-59), derived from data collected by the AVHRR instrument onboard the NOAA polar orbiting MetOp platforms, is available from the Climate Monitoring Satellite Application Facility (CM SAF) for the Arctic region. Monthly and weekly averaged products are available on a 15 km × 15 km grid, and an instantaneous product on a 1 km × 1 km grid. We use the latter product here because of its higher resolution. A summary of the product follows; for a full description see Karlsson et al. (2012). Albedo is calculated for each AVHRR-imaged pixel ($\sim 1.1 \text{ km}^2$), using satellite and sun viewing geometry and recorded top-of-atmosphere reflectances. In addition, land cover data, a cloud mask, surface pressure and water vapour from Deutsche Wetter Dienst are used. Cloud shadow effects are reduced by the removal of not only all cloud pixels, but also of all pixels neighbouring a cloud pixel. A topographical correction is applied to the observed reflectances and used to assign a reflectance to hidden slopes. Water vapour and surface pressure data from Deutsche Wetter Dienst are used under the assumption of constant ozone content and aerosol optical depth of the atmosphere, to convert top-of-atmosphere reflectances to surface reflectances. These surface reflectances are used to calculate black sky surface albedo, following Schaepman-Strub et al. (2006) for radiation of 0.25 to 2.5 μm – this is almost equivalent to full broadband surface albedo since shortwave infrared contributions to the full broadband albedo are generally very small, and we treat it here as equivalent. Distance-weighted interpolation was used to calculate the surface broadband albedo for the field site location from the 1 km × 1 km grid in which the CM SAF product is provided.

Only those images where all four closest pixels were assigned an albedo value were retained for further analysis. We also restricted the image acquisition time to within 6 h of local solar noon to avoid images acquired at high zenith angles. Finally, remaining images were grouped into daily bins before calculating the mean and standard deviation albedo value for each day.

TCD

8, 1337–1382, 2014

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



periods of sunny weather, before eventually joining the larger, visible drainage channels. The melt crust is formed by the penetration of solar radiation into the ice, which permits melting within a relatively thick surface layer and creates a wide variety of ice formations (Fig. 8). Exceptions to this pattern were periods of warm, windy weather, when the surface melt crust was quickly stripped away and replaced by a hard, slippery and impermeable surface; these times correspond to the higher (but still small) incidence of surface water in the quadrats in late July and early August.

The ice surface at the field site was initially quite smooth, mostly snow with a few ice patches and lumps of less than 50 cm in height (similar to the photo in Fig. 9a, taken on 1 June). As melt progressed, the surface became increasingly uneven with a mix of bumps up to approximately 150 cm high and melt channels up to 3 m deep (similar to the photo in Fig. 9b, taken on 31 July).

3.2 Productivity

In total, 17 incubation experiments were carried out between 4 June and 8 August (Fig. 7; Table 3). Results averaged over the regular weekly experiments in the light bottles showed changes in O_2 significantly different from zero ($p < 0.05$) in CI, DI and CS, and no significant change in CW, SN and OG. Significant inferred photosynthesis and respiration, as estimated from the combined light and dark bottle results (see Sect. 2.2), was found in DI, CS and OG (Table 3). In each of these three sample types, light bottles gained O_2 (stored C) while dark bottles lost O_2 (respired C). In contrast, CI samples significantly gained a very small but significant amount of O_2 in both the light and dark bottles, by amounts that were not significantly different from each other (Fig. 7a; $p > 0.05$). The greatest activity in the bottles was in OG, where mean photosynthesis and respiration rates were almost equal at $1.7 \text{ mgCL}^{-1} \text{ d}^{-1}$.

A better estimate of the variance within the main four sample types was provided by the additional incubation experiments with 24 bottles of the same type (12 light, 12 dark: Fig. 10). Although there was a wide spread of initial O_2 concentrations, especially in the debris-rich CS and DI samples, the variability of changes in dissolved O_2 after

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



24 h (which are the important results indicating the level of biological oxygen demand and inferred C storage) was much smaller. Specifically, standard deviations of ΔO_2 expressed as a fraction of the mean were 0.32, 0.27, 3.1 and 0.30 in light bottles containing CI, DI, CW and CS, respectively, and 0.49, 0.32, 1.5 and 0.13 in the dark bottles, respectively. Each of these means is significantly different from zero ($p < 0.05$) except for those for CW.

Seasonal variations in C storage within the bottles, as inferred from measured increases in dissolved O_2 , are shown in Fig. 7 for the four main surface types. There was no significant correlation between activity between any two pairs of surface types (Spearman's rank correlation; $p > 0.05$) and no clear seasonal pattern (Fig. 7), except for an apparently steady increase in the rates of photosynthesis and respiration in cryoconite hole (CS) samples from 11 June to 16 July. There was no significant correlation between incident SW radiation and C storage rate over the study period (Fig. 11); however, we found a strong link between incident SW intensity and productivity in the sub-24 h activity experiment on 4–5 August (Fig. 12), which is in contrast to the seasonal results. These short incubations showed a change from O_2 production (C storage) to O_2 storage (C release) during the darkest hours of the experiment (20:00–03:00 LT), but with an overall gain in dissolved O_2 (net C storage) after 24 h consistent with the standard experiments.

Variations in productivity with cryoconite hole size and location were investigated in an experiment on 6–7 August, when samples from 15 holes in 5 areas were incubated (Fig. 13). Spearman's correlation between hole size and oxygen demand showed weak negative correlation between hole size and net productivity in the light bottles ($r = -0.52$; $p = 0.027$) and stronger negative correlation between hole size and respiration rate in the dark bottles ($r = -0.65$; $p = 0.005$) (Fig. 13a). There was no apparent difference in productivity between the five locations sampled (Fig. 13b).

Biological oxygen demand in the bottles was converted to net inferred C flux for each surface type, using estimates of the sampled ice area in each bottle and the contribution of each surface type to the total surface area as measured in the quadrat

surveys. This yielded a seasonal pattern of C flux from each surface type (Fig. 14; Table 3) and an estimate of the total C flux at the field site during the summer study period of $3.4 \text{ mgC m}^{-2} \text{ d}^{-1}$. This mean was calculated from 66 daily values, each based on linearly interpolated quadrat and productivity data. The main contributions were $2.5 \text{ mgC m}^{-2} \text{ d}^{-1}$ from CS and $0.9 \text{ mgC m}^{-2} \text{ d}^{-1}$ from DI, while contributions from SN and CW were negligible. Note that CI was excluded from the total because the small gain in O_2 appeared to be an artefact of the method (see Sect. 4.3).

3.3 Albedo

Daily albedo data satisfying the quality control criteria (Sect. 2.3) were available for 69 out of the 153 days from 1 May to 30 September (Fig. 6g). Albedo (reported as mean $\pm 1\sigma$) from 1 to 25 May was $74 \pm 6\%$ with no significant trend. After a sharp drop to $57 \pm 3\%$ on 26 May, albedo fell unsteadily until early July, before stabilising at $39 \pm 2\%$ from 8 July to 9 August. Following fresh snow on 10 August, albedo was variable in the ca. 40–60% range before rising sharply on 1 September and remaining steady throughout that month at $73 \pm 3\%$.

4 Discussion

4.1 Surface characteristics

Results from the quadrat surveys, albedo retrieval and visual observations summarised above clearly demonstrate the great temporal and spatial variability in surface conditions at the field site. First, using observed patterns in surface classification and albedo we can divide the “summer” study period into four distinct periods, as follows.

Period 1. Pre-melt. Very little exposed ice before the winter snow has melted; cold, dry snow; no surface water; high albedo ($> 70\%$). In 2012, the field site was established on 26 May only a few hours before the onset of heavy rain marking the end of this period.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Period 2. Snow melt. 26–30 May. A rapid transition from dry snow to wet snow, slush, melt ponds and bare ice over just a few days, with a corresponding sharp drop in albedo. A similarly rapid transition was observed at a lower site 7 km from the margin (~ 600 m a.s.l.) in 2011. Period 2 is likely to be relatively short in this part of the ice sheet, due to the very low precipitation in comparison with other parts of the west coast (Burgess et al., 2010).

Period 3. Early summer ice surface. 30 May to 11 July. Superimposed ice and remaining snow patches gradually melt to reveal the dirty ice surface and cryoconite holes from the previous year. The area covered by cryoconite holes slowly increases to reach ~ 17%; albedo slowly decreases to a minimum (34 %).

Period 4. Late summer ice surface. 11 July to 31 August. Cryoconite hole cover remains steady while dirty ice coverage varies widely, depending on prevailing weather conditions (warm/cloudy or cold/sunny); albedo remains low but becomes more variable. This period ends with an unsteady transition back to Period 1 conditions as evidenced by the snowfall on 10–11 August and similar events observed in the albedo record as peaks after the field campaign had ended.

Once cryoconite coverage had stabilised (Period 4), the mean areal coverage of 16.7% (range 12% to 27% in individual quadrat surveys) was much greater than that measured by Cook et al. (2012) on their nearby ablation-zone transect of sites at ~ 67° N, where maximum coverage was reported as ~ 7% on 2–3 August 2010, but it is similar to coverage reported by Bøggild et al. (2010) along a transect of the very narrow ablation zone in northeast Greenland. Cook et al. (2012) analysed photographs of nine 10 cm × 10 cm quadrats to determine cryoconite hole extent at each site, therefore only sampling very small areas (noting the high spatial variability indicated by error bars in Fig. 6, and also that many holes exceed 10 cm diameter). We would argue that our strategy of five 2.7 m × 2.7 m quadrats (total 500 measurements) will have yielded a more spatially representative sample at our field site. Photographic analysis should be used with caution since many of the deep but narrow cryoconite holes quickly become obscured as the viewing angle deviates from vertical: for example,

a typical cryoconite hole 2 cm in diameter and 10 cm deep will be completely hidden in an image with a viewing angle of only $\tan^{-1}(2/10) = 11^\circ$ from vertical.

Despite the warm, cloudy conditions causing cryoconite holes to melt out and release their debris, resulting in the sharp increases in DI coverage following ablation peaks in July (Fig. 6), there was little corresponding reduction in CH coverage. This is probably because only the small, shallow holes melted out, and the debris contained in just a few holes is sufficient to create a much larger area of DI once it has been dispersed. The same conditions that lead to hole melt-out also caused the surface to become hard and impermeable, making the rapid dispersion of debris much easier than would have been the case over the permeable, rough surface typical of sunny conditions. Another possible process that could increase DI coverage is input of windblown dust – however, it is unclear why this dust would immediately settle in patches rather than being generally distributed over the ice surface, so do we do not believe this process made a significant contribution. Overall, the combination of an early-summer increase in CH and late-summer increase in DI caused a steady decrease in the area of CI from 98% on 5 June to a minimum of 41% on 8 August.

The increasing areal coverage of cryoconite holes through the melt season was mainly due to the re-emergence of the previous year's holes, which we believe became exposed by the melting of superimposed ice, rather than the formation of new holes. We infer this because (1) the increasing CH area did not occur at the expense of DI, (2) excavation of DI surfaces revealed no visible debris below the ice surface, (3) there was no visible evidence of significant input of wind blown dust (most clean ice remained clean), and (4) because of our observations of thick (> 10 cm) ice lids over newly emerging holes in June and July. These lids might have formed by refreezing of water in the hole over winter, by superimposed ice forming in snow above the hole, or both. In either case it seems that many holes can last for more than one season, becoming reactivated in the spring as has been observed elsewhere: for example, the Dry Valleys, Antarctica (Fountain et al., 2004). Some emerging holes had lids with a radial crystal structure characteristic of re-freezing processes, and were easy to spot in the

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rotting ice surface; others had little or no surface expression and were found only after the lid collapsed under foot pressure (these lids are more likely to consist of superimposed ice; their remains are shown in Fig. 8b). The large number of cryoconite holes apparently present beneath the surface early in the season shows that most holes persisted through the winter. At present it is uncertain how long the holes are active for in the spring before losing their winter ice lids, and this early season activity has yet to be measured.

Field data for this study were collected in 2012, which was notable for its unusually warm conditions across the whole ice sheet: in particular regarding the record-breaking melt extent and surface mass balance (Tedesco et al., 2012). Similarities between the seasonal changes in albedo observed in the present and previous studies (for example Knap and Oerlemans, 1996; Stroeve et al., 2001) suggest the pattern discussed above is persistent between years, including this extreme melt year. Extended visits to the same site in 2011 (25 June to 8 July; 31 July; 12 to 20 August) have also allowed qualitative visual comparison of the ice surface between the two summers. Many features were recognised in both seasons: the gradual exposure of cryoconite holes from under ice lids; patchiness of dirty ice coverage; increasing amplitude of the surface lumpiness as the melt season progresses. The only noted difference of significance was the frequent occurrence of warm, cloudy and windy conditions in July and August 2012, which created a hard and slippery ice surface and caused melt-out of some of the cryoconite holes – these events were not observed in 2011, when conditions were predominantly cooler and sunnier. This suggests a greater contribution of sensible or latent heat fluxes to surface melt in 2012.

4.2 Ice surface albedo

The mean albedo of 45% reported by Knap and Oerlemans (1996) for Zone II of the ablation zone on 25 July 1991 is similar to that retrieved from the 2012 AVHRR data for our field site once the albedo had stabilised early in Period 4 ($39 \pm 2\%$ from 8 July to 9 August). The seasonal changes in albedo (Fig. 6g) followed a similar pattern

TCO

8, 1337–1382, 2014

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to observations from this part of the ice sheet in previous remote sensing and AWS studies, in particular the gradual decrease in Period 3: for example at Site S6 on the K-Transect, located at the same altitude (1030 m) as the present field site (Knap and Oerlemans, 1996). However, while Knap and Oerlemans (1996) attributed albedo changes solely to variations in surface melt, the gradual decrease in albedo observed in our Period 3 occurred when there was an overall decrease in stored surface water and slush as the surface drainage system developed. Therefore, while the initial sharp drop at the end of May was due to a transition from loose, dry snow encountered on arrival at the site to water ponds and wet snow, the subsequent gradual decrease in albedo between 29 May and 10 July was more likely due to increasing exposure of cryoconite holes than to increasing melt water coverage (Fig. 6). This is consistent with the recent work of Wientjes et al. (2011). These two distinctly different responses demonstrate the ability of surface albedo retrievals to provide information on both surface melt and surface cryoconite hole coverage.

4.3 Surface productivity

Throughout the season, dirty ice (DI) and cryoconite water-plus-sediment (CS) incubations consistently showed O_2 loss in dark bottles and slight but significant O_2 gain in light bottles (Fig. 7b and c) corresponding to net C storage (see Sect. 3.2) indicative of active, weakly autotrophic biological communities as have been observed previously in cryoconite holes in Greenland and elsewhere (Anesio et al., 2009; Cook et al., 2012). In contrast, there was slight O_2 gain in both light and dark clean ice (CI) incubations (Fig. 7a), with O_2 gains found to be insignificantly different between the light and dark bottles. This O_2 gain was therefore a light-independent process and was most likely an artefact of the method, for example a small amount of O_2 may have been absorbed by the water when the oxygen probe was inserted to take the post-incubation reading. This problem would have had a greater effect in melted ice samples (CI, DI) than water samples (CW, CS) as melted ice samples were under-saturated with O_2 following melting: compare CI, DI starting O_2 levels with those of CW, CS in Fig. 10. Correcting

the DI results for this same artefact would reduce the total O₂ gain by about 30 %, from 1.1 down to 0.8 mgO₂L⁻¹d⁻¹. We also note that the measured DI productivity may not be a true representation of in-situ rates since we were using melted samples. Firstly, the solar radiation may have been weaker in the incubation environment than at the ice surface owing to the shading effect of the hole edges; secondly, the nutrient concentrations may have been altered during melting, for example with some chemical species becoming more diluted when water within the grain boundaries in the rotting surface ice was mixed with bulk melt from the surrounding crystals. In-situ measurements in ice would be extremely challenging, due to the open nature of the system (ice permeable to water and gases) and ice-water exchange of solutes during melting-refreezing processes. Nevertheless, our measurements represent the potential for activity in this surface environment.

In contrast to the clear seasonal changes in the ice surface characteristics, there was very little evidence for any seasonal signal in the results of productivity experiments, except for a general increase in O₂ consumption in cryoconite holes during Period 3 (Fig. 7c). This trend did not significantly alter the net productivity so was presumably balanced by increasing photosynthesis, suggesting a strong link between the rates of respiration and photosynthesis similarly to results reported by Hodson et al. (2010) and Telling et al. (2012a) but contrary to others (for example, net respiration: Stibal and Tranter, 2007). In our case, increasing respiration may reflect the increasing exposure time of holes to direct solar radiation following melting of the ice lids (i.e., increasing cell count), or a thicker accumulation of sediment. Unfortunately the sediment was always too thin (up to a few mm) to allow accurate thickness measurements. Interestingly, both the surface area and activity of cryoconite holes increased during Period 3, then remained steady in early Period 4, mirroring the surface albedo (decreasing in Period 3, steady in early Period 4). Total C flux into cryoconite holes increased during this time due to an increase in the area of exposed holes rather than changes in net productivity (Fig. 14).

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Combining results from the quadrat and BOD experiments yielded total C fixation over the course of the experiment (4 June–9 August) of 0.22 gC m^{-2} , of which DI and CS contributed 26 % and 74 %, respectively. The melt season continued beyond the end of the experiment, probably by ~ 3 weeks according to the albedo data, so the total flux for the complete season was most likely greater (simple extrapolation would yield 0.29 gC m^{-2}). However, the annual net C flux remains uncertain since the summer is relatively short and for much of the year there is either no daylight or the ice surface is covered by highly reflective snow which greatly reduces light intensity reaching the microbial communities beneath. Since high rates of respiration were recorded in dark bottles it is evident that any C stored during the summer might be subsequently released over the winter. The effect of this on annual fluxes will be dependent on the continuation of microbial activity in the extended periods without sunlight. Whilst much of this period is very cold, solute-rich water remains in the grain boundaries surrounding the ice crystals, which could allow microbial activity albeit at much reduced rates. This could have a significant effect on carbon fluxes and should clearly be addressed in future studies in Greenland.

The short term experiment with cryoconite hole sediment demonstrates the necessity of using incubations lasting for multiples of 24 h to minimise the effects of diurnal variations in activity driven by solar radiation intensity (Fig. 12). However, despite the clear dependency of radiation intensity on productivity at diurnal time scales, there was no clear relationship between activity and radiation intensity at seasonal time scales (Fig. 11), so we assume the incoming solar radiation intensity did not contribute significantly to the seasonal variability in productivity. Instead, other factors such as hole age, debris composition, debris thickness or water chemistry may have been important. Interestingly, Stibal et al. (2011) found slope angle was an important factor along their ablation zone transect at 67° N . However, their slopes were computed from GPS positions separated by many kilometres and are unlikely to reflect the local slope angle owing to the very rough ice sheet surface in the ablation zone at the horizontal scale of 1 to 100 m (compare local slopes in Figs. 2 and 9 with the calculated topographic slope

of less than 1°). The correlation reported by Stibal et al. (2011) is therefore unlikely to be a direct causal relationship.

In the same region of the ice sheet as our field site, Hodson et al. (2010) reported mean C storage rates in cryoconite holes of $-3.9 \mu\text{gCg}^{-1} \text{d}^{-1}$, which is equivalent to $-8.0 \text{mgCm}^{-2} \text{d}^{-1}$ using their value of 0.206gcm^{-2} of sediment in cryoconite holes. At Transect Site 7, representative of our field site, Stibal et al. (2011) measured NEP, photosynthesis, respiration C storage rates of 1.1 ± 0.5 , -4.9 ± 1.8 , $3.8 \pm 2.0 \text{mgCm}^{-2} \text{d}^{-1}$ (based on results reported in $\mu\text{gCm}^{-2} \text{d}^{-1}$ converted to $\text{mgCg}^{-1} \text{d}^{-1}$ using their measured surface debris cover of 180gcm^{-2}). By comparison, C fluxes from cryoconite holes in our study ranged from 5 to $33 \text{mgCm}^{-2} \text{d}^{-1}$ (mean $24 \text{mgCm}^{-2} \text{d}^{-1}$). Our data and those of previous studies highlight the great variability in measured C fluxes associated with microbial activity; even the direction of the flux (C storage or C release) varies between studies.

4.4 Linking albedo and productivity

Since there is a clear response of the AVHRR-derived albedo to the snow melt and increasing cryoconite hole exposure (Fig. 6; Periods P3–P4), it is likely that a relationship could be established between retrieved albedo and cryoconite hole coverage, which would enable a much more wide-ranging assessment of cryoconite hole coverage than has been possible in this or previous field studies. This could then be used to predict C fluxes given an appropriate estimate of cryoconite hole C fluxes, although the variability presented here shows there would be a large associated uncertainty. The albedo product used here appeared less sensitive to DI coverage, but it is worth noting that the sudden increases in DI during high melt events occurred exclusively in periods of cloudy weather, when albedo data are not available from satellite remote sensing, so it does not seem likely that AVHRR data will be useful for assessing DI coverage. Further, there was great spatial variability in DI cover (Fig. 6c), making it difficult to correlate albedo and DI cover with a relatively small data set. The contribution of DI to the mean C flux was approximately one third that of cryoconite holes, therefore an

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



alternative approach to measuring DI coverage remotely may be needed for estimates of ice sheet C fluxes.

5 Conclusions

A time-series of ice surface characteristics and productivity from 26 May to 10 August 2012 has provided a detailed insight into seasonal changes in ice surface processes during the melt season, at a site about half way between the margin and equilibrium line in west Greenland. Specific findings of interest are as follows:

1. Based on changes in ice surface characteristics observed in ground observations and satellite AVHRR imagery, the melt season was classified into four distinct periods. These are: winter snow cover (P1); snow melt (P2); melting of superimposed ice and exposure of cryoconite holes (P3); and the mid- to late-summer ice surface (P4).
2. There was a marked surface response to major melt events, most notably expressed as loss of the several centimetre-thick, permeable melt crust to expose the impermeable underlying ice, and the dispersion of debris from shallow cryoconite holes.
3. In-situ incubation experiments revealed active microbial communities, as have been reported in limited previous studies in Greenland. Rates of activity in the different ice surface environments were very variable both between environments and between incubations in the same environment. There was no clear seasonal signal, aside from an apparent correlation between albedo, cryoconite hole coverage and cryoconite hole productivity in melt season Periods 3 and 4. On average over the study period we found significant net O₂ production (inferred as net C storage) in dirty ice and cryoconite holes. There was no significant O₂ gain or loss in snow or water samples.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



While there is good evidence from this and previous studies that seasonal changes in ice surface characteristics are broadly persistent between years, the very variable productivity results obtained at this field site and elsewhere suggest that there remains great uncertainty in the magnitude of biologically mediated carbon fluxes into the ablation zone of the ice sheet. However, the link between cryoconite hole productivity, cryoconite hole areal coverage and ice surface albedo shows there is potential for the large-scale estimation of biological processes and associated carbon fluxes, which could aid in extrapolating point observations up to the ablation zone scale. However, the areas of dirty ice also contribute significantly to carbon fluxes but a similar correlation between dirty ice coverage and albedo was not apparent.

Acknowledgements. This work was funded by NERC grant NE/H023879/1. M. van den Broeke (IMAU) kindly provided the SW radiation data, and AVHRR imagery was supplied by the EU-METSAT Climate Monitoring Satellite Application Facility (CM SAF).

References

- Anesio, A. M., Hodson, A. J., Fritz, A., Psenner, R., and Sattler, B.: High microbial activity on glaciers: importance to the global carbon cycle, *Glob. Change Biol.*, 15, 955–960, doi:10.1111/j.1365-2486.2008.01758.x, 2009.
- Bayley, W. S.: Minerology and petrography, *Am. Natur.*, 25, 138–146, 1891.
- Bøggild, C. E., Brandt, R. E., Brown, K. J., and Warren, S. G.: The ablation zone in northeast Greenland: ice types, albedos and impurities, *J. Glaciol.*, 56, 101–113, 2010.
- Burgess, E. W., Forster, R. R., Box, J. E., Mosley-Thompson, E., Bromwich, D. H., Bales, R. C., and Smith, L. C.: A spatially calibrated model of annual accumulation rate on the Greenland Ice Sheet (1958–2007), *J. Geophys. Res.*, 115, F02004, doi:10.1029/2009JF001293, 2010.
- Cook, J. M., Hodson, A. J., Anesio, A. M., Hanna, E., Yallop, M., Stibal, M., Telling, J., and Huybrechts, P.: An improved estimate of microbially mediated carbon fluxes from the Greenland Ice Sheet, *J. Glaciol.*, 58, 212, doi:10.3189/2012JoG12J001, 2012.
- Duynkerke, P. G. and van den Broeke, M. R.: Surface energy balance and katabatic flow over glacier and tundra during GIMEX-91, *Global Planet. Change*, 9, 17–28, 1994.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fountain, A. G., Tranter, M., Nysten, T. H., Lewis, K. J., and Mueller, D. R.: Evolution of cryoconite holes and their contribution to runoff from glaciers in the McMurdo Dry Valleys, Antarctica, *J. Glaciol.*, 50, 35–45, 2004.

5 Gerdel, R. W. and Drouet, F.: The cryoconite of the Thule Area, Greenland, *T. Am. Microsc. Soc.*, 79, 256–272, 1960.

Greuell, W., Denby, B., Van der Wal, R. S. W., and Oerlemans, J.: 10 year of mass-balance measurements along a transect near Kangerlussuaq, central West Greenland, *J. Glaciol.*, 47, 157–158, 2010.

10 Hodson, A., Anesio, A. M., Ng, F., Watson, R., Quirk, J., Irvine-Fynn, T., Dye, A., Clark, C., McCloy, P., Kohler, J., and Sattler, B.: A glacier respire: Quantifying the distribution and respiration CO₂ flux of cryoconite across an entire Arctic supraglacial ecosystem, *J. Geophys. Res.*, 112, G04S36, doi:10.1029/2007JG000452, 2007.

15 Hodson, A., Bøggild, C., Hanna, E., Huybrechts, P., Langford, H., Cameron, K., and Houldsworth, A.: The cryoconite ecosystem on the Greenland Ice Sheet, *Ann. Glaciol.*, 51, 123–129, 2010.

Irvine-Fynn, T. D. L., Bridge, J. W., and Hodson, A.: In situ quantification of supraglacial cryoconite morphodynamics using time-lapse imaging: an example from Svalbard, *J. Glaciol.*, 57, 651–657, 2011.

20 Karlsson, K.-G., Riihelä, A., Müller, R., Meirink, J. F., Sedlar, J., Stengel, M., Lockhoff, M., Trentmann, J., Kaspar, F., Hollmann, R., and Wolters, E.: CLARA-A1: CM SAF Clouds, Albedo and Radiation dataset from AVHRR data – Edition 1 – Monthly Means/Daily Means/Pentad Means/Monthly Histograms, Satellite Application Facility on climate Monitoring, doi:10.5676/EUM_SAF_CM/CLARA_AVHRR/V001, 2012.

25 Kipp and Zonen: CNR1 Net radiometer instruction manual, available at: <http://www.kippzonen.com/Download-Center> (last access: 23 February 2014), 2002.

Knap, W. H. and Oerlemans, J.: The surface albedo of the Greenland ice sheet: satellite-derived and in situ measurements in the Søndre Strømfjord area during the 1991 melt season, *J. Glaciol.*, 42, 364–374, 1996.

30 Konzelmann, T. and Braithwaite, R. J.: Variations of ablation, albedo and energy balance at the margin of the Greenland ice sheet, Kronprins Christian Land, eastern north Greenland, *J. Glaciol.*, 41, 174–182, 1995.

Langford, H., Hodson, A., Banwart, S., and Bøggild, C.: The microstructure and biogeochemistry of Arctic cryoconite granules, *Ann. Glaciol.*, 51, 87–94, 2010.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nolin, A. W. and Payne, M. C.: Classification of glacier zones in western Greenland using albedo and surface roughness from the Multi-angle Imaging SpectroRadiometer (MISR), *Remote Sens. Environ.*, 107, 264–275, 2007.

Nolin, A. W., Fetterer, F. M., and Scambos, T. A.: Surface roughness characterizations of sea ice and ice sheets: case studies with MISR data, *IEEE T. Geosci. Remote*, 40, 1605–1615, 2002.

Oerlemans, J. and Vugts, H. F.: A meteorological experiment in the melting zone of the Greenland ice sheet, *B. Am. Meteorol. Soc.*, 74, 355–365, 1993.

PreSens: Fibox 3 Instruction Manual, available at: http://www.presens.de/fileadmin/user_upload/downloads/manuals/IM_Fibox_3_dv2.pdf (last access: 23 February 2014), 2006.

Schaepman-Strub, G., Schaepman, M. E., Painter, T. H., Dangel, S., and Martonchik, J. V.: Reflectance quantities in optical remote sensing – definitions and case studies, *Remote Sens. Environ.*, 103, 27–42, 2006.

Smeets, C. J. P. P. and van den Broeke, M. R.: Temporal and spatial variations in the aerodynamic roughness length in the ablation zone of the Greenland Ice Sheet, *Bound.-Lay. Meteorol.*, 128, 315–338, doi:10.1007/s10546-008-9291-0, 2008.

Steinböck, O.: Über Kryokonitlöcher und ihre biologische Bedeutung, *Z. Gletscherkd.*, 24, 1–21, 1936.

Stibal, M. and Tranter, M.: Laboratory investigation of inorganic carbon uptake by cryoconite debris from Werenskioldbreen, Svalbard, *J. Geophys. Res.*, 112, G04S33, doi:10.1029/2007JG000429, 2007.

Stibal, M., Telling, J., Cook, J., Mak, K. M., Hodson, A., and Anesio, A. M.: Environmental controls on microbial abundance and activity on the Greenland Ice Sheet: a multivariate analysis approach, *Microb. Ecol.*, 63, 74–84, doi:10.1007/s00248-011-9935-3, 2012.

Stroeve, J.: Assessment of Greenland albedo variability from the advanced very high resolution radiometer Polar Pathfinder data set, *J. Geophys. Res.*, 106, 33989–34006, 2001.

Takeuchi, N.: Optical characteristics of cryoconite (surface dust) on glaciers: the relationship between light absorbency and the property of organic matter contained in the cryoconite, *Ann. Glaciol.*, 34, 404–419, 2002.

Tedesco, M., Fettweis, X., Mote, T., Wahr, J., Alexander, P., Box, J. E., and Wouters, B.: Evidence and analysis of 2012 Greenland records from spaceborne observations, a regional climate model and reanalysis data, *The Cryosphere*, 7, 615–630, doi:10.5194/tc-7-615-2013, 2013.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Telling, J., Anesio, A. M., Hawkins, J., Tranter, M., Wadham, J. L., Hodson, A. J., Irvine-Fynn, T., and Yallop, M. L.: Measuring rates of gross photosynthesis and net community production in cryoconite holes: a comparison of field methods, *Ann. Glaciol.*, 51, 153–162, 2010.
- 5 Telling, J., Anesio, A. M., Tranter, M., Stibal, M., Hawkings, J., Irvine-Fynn, T., Hodson, A., Butler, C., Yallop, M., and Wadham, J. L.: Controls on the autochthonous production and respiration of organic matter in cryoconite holes on High Arctic glaciers, *JGR-Biogeosci.*, 117, G01017, doi:10.1029/2011JG001828, 2012a.
- 10 Telling, J., Stibal, M., Anesio, A. M., Tranter, M., Nias, I., Cook, J., Bellas, C., Lis, G., Wadham, J. L., Sole, A., Nienow, P., and Hodson, A.: Microbial nitrogen cycling on the Greenland Ice Sheet, *Biogeosciences*, 9, 2431–2442, doi:10.5194/bg-9-2431-2012, 2012b.
- Uetake, J., Naganuma, T., Hebsgaard, M. B., Kanda, H., and Koshima, S.: Communities of algae and cyanobacteria on glaciers in west Greenland, *Polar Science*, 4, 71–80, 2010.
- van de Wal, R. S. W. and Oerlemans, J.: An energy balance model for the Greenland Ice Sheet, *Global Planet. Change*, 9, 115–131, 1994.
- 15 Wientjes, I. G. M., Van de Wal, R. S. W., Reichert, G. J., Sluijs, A., and Oerlemans, J.: Dust from the dark region in the western ablation zone of the Greenland ice sheet, *The Cryosphere*, 5, 589–601, doi:10.5194/tc-5-589-2011, 2011.
- 20 Yallop, M. L., Anesio, A. M., Perkins, R. G., Cook, J., Telling, J., Fagan, D., MacFarlane, J., Stibal, M., Barker, G., Bellas, C., Hodson, A., Tranter, M., Wadham, J., and Roberts, N. W.: Photophysiology and albedo-changing potential of the ice algae community on the surface of the Greenland ice sheet, *ISME J.* 6, 2302–2313, 2012.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Cryoconite hole carbon flux measurements in Greenland.

Source	Method	Carbon storage rate ^a $\mu\text{gC}[\text{g or L}]^{-1}\text{d}^{-1}$	Notes
Anesio et al. (2009)	¹⁴ C	Water 54 ± 60 Sediments 115 ± 56	Frøya Glacier, east Greenland, 74° N; date not specified.
Hodson et al. (2010)	TDIC, O ₂	Sediments Mean -3.9^{b} <i>P</i> : 12–30 <i>R</i> : 4.8–29	South-west margin, 67° N; 2 km transect at margin (~ 2 % of melt zone); 23–30 Aug 2008
Stibal et al. (2012); Cook et al. (2012)	TDIC	Sediments Mean: 5.4^{c} Range from below detection limit to 22 ± 4.8	South-west margin, 67° N; 65 km transect with 9 stations from margin to slush zone; Aug 2010

^a Units $\mu\text{gCg}^{-1}\text{d}^{-1}$ or $\mu\text{gCL}^{-1}\text{d}^{-1}$ represent the carbon flux per gram of dry sediment per day or the carbon flux per litre of water per day.

^b Mean NEP calculated from O₂ and TDIC incubation experiments BI1 to BI5 reported in Table 2 of Hodson et al. (2010).

^c Mean NEP of 9 stations along the transect using results reported in their Table 3.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Table 2. List of surface types identified at the field site. All except Type F were present in the quadrat surveys.

Surface Type	Code	Description
Snow	S	Seasonal snow cover, including a coarse-grained granular material that is probably rotting superimposed ice (Fig. 4a).
Water	W	Surface water that is not covering cryoconite debris (Fig. 4c).
Clean ice	CI	Englacial ice with no visible debris (Fig. 4a–c).
Dirty ice	DI	Englacial ice with debris partially or completely covering the surface (Figs. 1a and b, 4b).
Cryoconite hole	CH	Water-filled holes with a layer of debris in the bottom (Figs. 1c, 4b and c).
Fractures, moulins	F	Open fractures, crevasses and moulins were all present close to the field site but were not recorded in the quadrat surveys.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Table 3. Summary of productivity data for the regular (~weekly) incubation experiments, presented as mean $\pm 1\sigma$ of carbon storage rates in n incubation experiments. P, R and net are results for inferred photosynthesis only, inferred respiration only, and both processes combined. C storage is the carbon storage rate in the bottle. C flux is the seasonal mean of C storage when converted to carbon flux, based on the surface area sampled to fill the bottle; C flux (total) is the C flux weighted by the fractional surface area of that type, with [5 %, 95 %] percentiles of 66 interpolated daily values. Observation periods for productivity were 4 June–8 August 2012 except snow (9 June only) and orange-green material (4 and 9 July only).

Surface type	n		C storage $\text{mgCL}^{-1}\text{d}^{-1}$	C flux $\text{mgCm}^{-2}\text{d}^{-1}$	C flux (total) $\text{mgCm}^{-2}\text{d}^{-1}$
Clean ice (CI)	10	Net	0.13 ± 0.08	5.7 ± 3.4	$2.4 [0.30, 4.1]$
		P	0.002 ± 0.048		
		R	0.12 ± 0.08		
Dirty ice (DI)	10	Net	0.40 ± 0.20	18 ± 9.0	$0.89 [0.07, 3.0]$
		P	0.64 ± 0.31		
		R	-0.24 ± 0.17		
Cryoconite hole water (CW)	10	Net	0.004 ± 0.010	0.5 ± 8.1	$-0.015 [-0.16, 0.001]$
		P	-0.010 ± 0.035		
		R	0.014 ± 0.16		
Cryoconite hole water & debris (CS)	8	Net	0.24 ± 0.11	30 ± 14	$2.5 [0, 4.7]$
		P	1.0 ± 0.33		
		R	-0.76 ± 0.47		
Snow (SN)	1	Net	-0.025 ± 0.027	-0.23 ± 0.24	$-0.010 [-0.060, 0]$
		P	-0.003 ± 0.035		
		R	-0.022 ± 0.022		
Orange-green material (OG)	2	Net	0.002 ± 0.064	0.3 ± 8.1	–
		P	1.7 ± 0.22		
		R	-1.7 ± 0.3		

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.




Fig. 1. Debris on the ice sheet surface at approximately 67.0° N, 049.3° W, 1030 m, in the 2012 melt season. **(a)** Dispersed debris. The foreground shows debris released by the melting-out of cryoconite holes while the more uniform cover in the background is typical “dirty ice”. The linear feature oriented right to left is a healed fracture. The image width at the surface is ~ 50 cm. **(b)** Thick deposits in a slowly draining lake bed, here reaching ~ 6 cm thick. **(c)** Cryoconite holes. The plastic pole is marked at 30 cm intervals. These are relatively small cryoconite holes; some reach over 1 m in diameter.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



Fig. 2. The ice sheet surface at 67.0° N, 050.0° W, 580 m (7 km from the margin), early in the melt season (10 June 2011). The foreground is a recently-drained pond with cryoconite debris and “spikes” of clean ice that have melted much more slowly due to their higher albedo. The “hills” in the background have heights of 3–5 m and very low debris cover.

Seasonal changes of
ice surface
characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Field site location (red triangle) in west Greenland, with the band of dark ice oriented north-south clearly visible in the ablation zone. The blue circle west of the field site marks Kangerlussuaq. The background image is a MODIS image acquired on 17 July 2012.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.




Fig. 4. (a) Q2, 1 June 2012, showing the quadrat ropes and pole. Surface cover is 55% snow (S) and 45% clean ice (CI). (b) Lower left corner of Q4, 13 July 2012, with clean ice (CI), dirty ice (DI) and cryoconite holes (CH). (c) Q4 on 29 July, with a very small melt water channel linking cryoconite holes under the mark on the pole. (d) Locations of quadrats on 1 June 2012, co-ordinates are relative to the portal from which melt water draining from the field site emerges at the ice margin.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Seasonal changes of
ice surface
characteristics**D. M. Chandler et al.




Fig. 5. (a) Sample collection for the cryoconite water plus sediment incubation (13 July) left 24 round spaces, each about the diameter of the BOD bottle (~ 65 mm). (b) Orange-green material sampled for the 4 and 9 July incubations. The lid diameter is 27 mm.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Seasonal changes of ice surface characteristics

D. M. Chandler et al.




Fig. 6. Percentage surface area (SA) covered by **(a)** snow, **(b)** clean ice, **(c)** dirty ice, **(d)** cryoconite holes and **(e)** water. Error bars are the range observed in 5 quadrats. **(f)** Daily ablation rate showing mean of 5 measurements (blue line); error bars are $\pm 1\sigma$. **(g)** Daily mean AVHRR-derived surface albedo $\pm 1\sigma$. Vertical red lines are *approximate* boundaries between periods P1 to P4 (see text).

Seasonal changes of ice surface characteristics

D. M. Chandler et al.




Fig. 7. Seasonal variation in productivity as measured by O_2 production or inferred C storage in the BOD incubation experiments for **(a)** clean ice; **(b)** dirty ice; **(c)** cryoconite holes (cryoconite debris plus water); and **(d)** cryoconite hole water. Symbols are respiration (R ; red squares) and net carbon storage (N ; blue circles), both measured directly, and inferred photosynthesis (P ; green triangles) calculated by $P = N - R$. Note the different scales on the vertical axes. Arrows mark the times of 24-bottle single-sample-type variance experiments (see Fig. 10) and red lines mark the approximate change from period P3 to P4 (see text).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal changes of ice surface characteristics

D. M. Chandler et al.




Fig. 8. Ice formations arising from radiative melt. **(a)** “Feathers” ca. 5–10 cm long, overlying a permeable melt crust a further several cm thick. These needed several days of cold, sunny weather to form. **(b)** “Mushrooms”, approximately 20 cm diameter and 15 cm tall, which we believe formed from remnants of the previous year’s cryoconite holes. **(c)** “Stacks” of cryoconite hole lids, formed when the water in the cryoconite hole freezes overnight. Each lid represents the water level that night, then the water level drops during the day as the surrounding ice surface ablates.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 9. Quadrat Q3 on **(a)** 1 June 2012, with the pole on Row 12 and **(b)** 31 July 2012, with the pole on Row 1. The photos were taken looking approximately northeast. Note the darker and rougher surface on 31 July compared with 1 June.

Seasonal changes of
ice surface
characteristics

D. M. Chandler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 10. Dissolved O₂ concentrations before and after incubation in experiments with 12 light (XX-L) and 12 dark bottles (XX-D) of each sample type. Samples plotting ABOVE the dashed line GAINED oxygen. Light samples reflect the balance between rates of respiration and photosynthesis, while dark samples reflect respiration rates only.




Fig. 11. Relationship between incident SW radiation intensity and inferred C storage rate by photosynthesis, for clean ice (CI, blue), dirty ice (DI, brown), cryoconite holes (CH, black) and cryoconite hole water (CW, purple).

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal changes of ice surface characteristics

D. M. Chandler et al.




Fig. 12. Changes in cryoconite hole C storage at sub-diurnal time scales. **(a)** Cumulative C storage as measured in bottles incubated for 3 to 24 h. **(b)** Equivalent C flux in the same bottles as part **(a)**, plotted alongside incident shortwave radiation intensity (3 h averages of 30 min data).

Seasonal changes of ice surface characteristics

D. M. Chandler et al.




Fig. 13. Variation in biological oxygen demand with **(a)** cryoconite hole surface area and **(b)** sampling location, 6–7 August 2012. To convert the measured O₂ demand to inferred C storage in mgCm⁻²d⁻¹, multiply the vertical axis values by 29.1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 14. Seasonal variation in C fluxes from the field site, showing individual contributions from DI and CS surfaces. Daily fluxes are plotted, which were calculated by linear interpolation of ~ weekly incubation measurements and quadrat surveys.

Seasonal changes of ice surface characteristics

D. M. Chandler et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

