

Abstract

We use a full-Stokes 2-D model (Elmer/Ice) to investigate the flow and calving dynamics of Store Gletscher, a fast flowing outlet glacier in West Greenland. Based on a new, subgrid-scale implementation of the crevasse depth calving criterion, we perform two sets of simulations; one to identify the primary forcing mechanisms and another to constrain future stability. We find that the mixture of icebergs and sea-ice, known as ice mélange or sikussak, is principally responsible for the observed seasonal advance of the ice front, whereas submarine melting plays a secondary role. Sensitivity analysis demonstrates that the glacier's calving dynamics are sensitive to seasonal perturbation, but are stable on interannual timescales due to the glacier's topographic setting. Our results shed light on the dynamics of calving glaciers while explaining why neighbouring glaciers do not necessarily respond synchronously to changes in atmospheric and oceanic forcing.

1 Introduction

Recent studies show accelerating net mass loss from the Greenland Ice Sheet (GrIS) (Khan et al., 2010; Rignot and Kanagaratnam, 2006; Howat et al., 2007), raising concerns about its future response to changing global climate and the impact this might have on global sea level. The two factors, which govern this loss, are (1) an overall negative surface mass balance stemming from intensified surface melting in the ice sheet's ablation zone (Hanna, 2005; van den Broeke et al., 2009; Enderlin et al., 2014), and (2) faster rates of ice discharge through calving glaciers which terminate in fjords (Luckman and Murray, 2005; Howat et al., 2005; Rignot and Kanagaratnam, 2006; Howat et al., 2007). The latter (dynamic) mechanism accounted for $\sim 67\%$ of the total net ice loss in 2005 (Rignot and Kanagaratnam, 2006), but less in recent years (Enderlin et al., 2014), highlighting the sensitivity of Greenland's marine-terminating glaciers to

TCD

8, 3525–3561, 2014

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the transient pulse of warm Atlantic water circumnavigating Greenland over the last decade (Holland et al., 2008; Straneo et al., 2010; Christoffersen et al., 2011).

Owing to the advancement of surface mass balance models over the last two decades (Hanna, 2005; Box et al., 2006; van den Broeke et al., 2009; Enderlin et al., 2014), this process is well represented in global sea level predictions (Stocker, 2013). The rapid dynamics associated with sudden increases in the discharge of ice into fjords by marine-terminating glaciers are, on the other hand, complex and poorly understood, and their relationship with climate remains elusive and is so far unconstrained (Stocker, 2013). The main processes involved in rapid dynamics are fast glacier flow and calving, i.e. the mechanism whereby pieces of ice and bergs break off glaciers terminating in water. These processes are complex because they interact with and respond to atmospheric as well as oceanic forcing effects. As such, calving and its associated dynamics comprise one of the most significant uncertainties in predictions of future ice sheet mass balance and sea level change.

While atmospheric processes were previously thought to be the main driver of rapid ice sheet dynamics (Zwally et al., 2002), recent studies point to warm water in coastal currents as the main forcing of mass loss by discharge (Holland et al., 2008). The rapid acceleration of Jakobshavn Isbræ from $\sim 4000 \text{ m a}^{-1}$ in 1995 to $\sim 17000 \text{ m a}^{-1}$ in 2012, is clearly linked to the continuing retreat of the calving ice front over this period (Joughin et al., 2012, 2014), and it has been hypothesised that submarine melting plays a crucial role in driving this retreat (Holland et al., 2008; Motyka et al., 2010). Spaceborne tracking of calving fronts also show that recent glacier retreat along the East Greenland coastline has been widespread and synchronous below 69° N , but largely absent at higher latitudes, where coastal water is much colder (Seale et al., 2011). This suggests that these glaciers are retreating in response to changes in the ocean system. Warmer fjord water increases the rate of submarine melting of the calving terminus. This effect is further amplified by atmospheric processes; buoyant pro-glacial plumes, driven by the delivery of surface meltwater to the terminus by the subglacial hydrological system, are capable of significantly increasing melt rates (Jenkins, 2011).

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Undercutting of calving ice fronts by submarine melting should, in addition, amplify calving rate due to the stress response (O’Leary and Christoffersen, 2013).

The formation of ice mélange, a rigid mixture of bergs and bergy bits, held together by sea-ice, henceforth referred to simply as mélange, may also play an important role with regard to rapid ice sheet dynamics (Sohn et al., 1998; Joughin et al., 2008). Data from Jakobshavn Isbræ indicate a complete cessation of calving when the glacier is buttressed by mélange, a response that explains why the glacier advances by up to 5 km in winter (Amundson et al., 2008) and why the glacier retreats suddenly when the mélange disintegrates (Joughin et al., 2008). A similar correspondence between mélange clearing date and increasing calving rate has been found for a number of glaciers, including those near Uummannaq in West Greenland (Howat et al., 2010). Walter et al. (2012) used changes in velocity observations and a force balance technique to infer a buttressing stress of 30–60 kPa exerted by mélange onto the terminus of Store Gletscher. This buttressing effect, and the effect of submarine melting (Xu et al., 2013), appear to be crucial for the calving dynamics of this glacier.

In this paper we present results from a numerical model developed using the open-source finite element (FEM) modelling package, Elmer/Ice, with newly implemented calving dynamics. Theoretical consideration of the calving process indicates the importance of the near-terminus stress field in controlling the propagation of crevasses and the detachment of icebergs (Nye, 1957; van der Veen, 1998a, b; Benn et al., 2007a, b). Linking calving to crevasse propagation and stress in this way provides a useful and physically-based framework for investigating calving in numerical models of glacier dynamics. Here, we implement a calving model based on the penetration of both surface and basal crevasses (Nick et al., 2009, 2010), and incorporate the full stress solution into the crevasse depth criterion, after Nye (1957). We use this model to investigate the seasonal dynamics of Store Gletscher, a fast flowing outlet glacier in the Uummannaq region of West Greenland. To examine the calving process, we focus on the calving front’s position and seasonal fluctuation, which have been stable and periodic, respectively, for at least four decades (Weidick, 1995, p.C41). We investigate the effects of

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



submarine melting, mélange buttressing and glacier geometry on calving, with the aim of identifying the role of each mechanism in driving the observed seasonal variability at the front. We find that mélange is likely to be the primary driver, and that submarine melting plays a secondary role. We also find that the topographic setting of Store Gletscher is responsible for its observed stability.

2 Store Gletscher

Store Gletscher, henceforth referred to as Store, is a fast-flowing marine terminating outlet glacier located in Ikerasak Fjord, near Uummannaq in West Greenland (Fig. 1). The glacier drains an area of 35 000 km² and is 5 km wide at the terminus, where surface velocity reaches $\sim 6600 \text{ m a}^{-1}$ (Joughin et al., 2011). The location of the terminus coincides with a bottleneck in fjord width (Fig. 1), as well as a pronounced basal pinning point (Fig. 2), suggesting that fjord topography may play an important role in calving dynamics.

In terms of climate, data from the Regional Atmospheric Climate Model (RACMO) suggest that $\sim 2 \text{ km}^3$ of meltwater forms on the surface of Store between June and August (Ettema et al., 2009). Recent modelling work (Xu et al., 2013) show that submarine melting at the terminus may occur at rates of 8 m d^{-1} in summer because a large proportion of runoff is discharged subglacially into Ikerasak Fjord. The latter is established from observations, which show upwelling of dirty, subglacially derived meltwater near the centre of the calving ice front during summer months (Chauché et al., 2013) (Fig. 1). The high melt rates are caused by entrainment of warm ambient fjord water into buoyant meltwater plumes, which rise rapidly in front of the glacier, from a depth of 490 m below sea level due to forced convection (Jenkins, 2011). The annual formation of mélange occurs in late January or early February and it tends to collapse around the end of May (Howat et al., 2010).

Store exhibits characteristic seasonal variabilities in terms of calving front position and velocity (Howat et al., 2010). Estimates of the terminus velocity of Store

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



terminus, we adopt a spatial resolution which varies from 250 m in the upper region of the glacier to 20 m near the terminus (Fig. 2b). The model evolves through time with a timestep of 1 day.

Temperature is an important factor in the stress–strain relationship of ice (Cuffey and Paterson, 2010). However, near the terminus, which is our region of interest, extensive crevassing makes the implementation of temperature difficult. The ability of subglacial meltwater to penetrate upwards through basal crevasses, as well as the effect of air circulation in surface crevasses, is likely to significantly affect the temperature profile of the ice. Due to these complications, and the lack of observations to constrain ice temperature, we assume for the sake of simplicity that the glacier is isothermal at -10°C .

Because basal friction exerts a critical control on the dynamics of fast flowing glaciers in general, we first use the adjoint inverse method (Gillet-Chaulet et al., 2012) to identify the basal friction profile which results in surface velocity as observed along the flowline. The result of the inverse method is a profile for the basal friction parameter (β^2) which is related to basal velocity (U_b) and basal shear stress (τ_b) by the relation (MacAyeal, 1992):

$$\tau_b = \beta^2 U_b \quad (1)$$

To integrate seasonal variation in ice flow in response to seasonal change in basal friction, we run the inverse model for both the summer and winter observed velocity profiles, thereby obtaining two basal friction profiles. A seasonal variability in ice flow, very similar to what is observed in reality, is imposed by varying the basal traction coefficient sinusoidally between summer and winter values.

3.2 Boundary conditions

Initial surface elevation along the modelled flowline is prescribed from the GIMP DEM product (Howat et al., 2014). The bed profile is obtained from airborne geophysical surveys carried out by the Greenland Outlet Glacier Geophysics (GrOGG) Project and

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NASA's Operation IceBridge (<https://espo.nasa.gov/missions/oib/>). We use a mass-conservation algorithm similar to that of McNabb et al. (2012) to constrain ice thickness and bed topography in the heavily crevassed region of fast flow near the terminus, where radar data are sparse.

Ice thickness evolves through time according to the mass continuity equation (Cuffey and Paterson, 2010), and we add and subtract mass according to RACMO surface mass balance data averaged between 1985 and 2008. The ice surface is treated as a stress free boundary, as we assume atmospheric pressure to be negligible. At the ice base, friction is prescribed through inverse methods as described above, except under the floating tongue which, when it exists, is a frictionless free surface. At the calving terminus, we apply an external pressure equal to the hydrostatic pressure from seawater (see Eq. 5 below). Above sea-level, atmospheric pressure is neglected.

We simulate the seasonal advance and retreat of Store Gletscher's floating tongue using an implementation of grounding line dynamics developed by Favier et al. (2012). The grounding line algorithm compares external water pressure and ice overburden pressure to detect where the glacier is floating, and modifies basal friction accordingly.

3.3 New scheme for implementation of flow convergence

Similar to most outlet glaciers, Store undergoes significant lateral narrowing as ice flows from catchment to coast. As such, it is important that dynamic effects from sidewall drag (Raymond, 1996) and ice convergence (Thomas et al., 2003) are accounted for.

Gagliardini et al. (2010) implemented a parameterization for sidewall friction in Elmer/Ice, and we use it here. The issue of ice convergence in full-Stokes 2-D models, however, has thus far received little attention from the glacier modelling community. Here, we developed a routine which adds flux sources to elements along the flow-line, corresponding to the downstream narrowing of the glacier. We derive a flux convergence term (see Supplement) and add it to the Stokes incompressibility equation

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where x is the direction of ice flow, and z the vertical. We multiply τ_e in Eq. (3) by the sign function of longitudinal deviatoric stress (τ_{xx}) to ensure crevasse opening is only predicted under longitudinal extension ($\tau_{xx} > 0$).

The second term on the right hand side of Eq. (3) represents creep closure, where ρ_i is the density of glacier ice, g is the force of gravity and d is depth through the ice.

The final term in Eq. (3) is water pressure (P_w) which acts to open crevasses when present. In basal crevasses, P_w is controlled by the subglacial hydrological system, and in surface crevasses, it is related to the depth of water in the crevasse.

Crevasses will exist wherever σ_n is positive, and ice remains intact elsewhere. Evaluating Eq. (3) for both surface and basal crevasses at every node in our model allows us to define “zero contours” which represent the base and top of surface and basal crevasse fields, respectively. The modified crevasse-penetration calving criterion (Nick et al., 2010) predicts that calving will occur where and when these zero contours meet. By calculating the net stress at every node, and interpolating the nodal values to find the zero contours (Fig. 3), we arrive at a calving implementation which accounts for changes in stress between surface and interior and which is independent of the model’s mesh resolution.

The magnitudes of the force components of Eq. (3) vary greatly between the surface and bed. Specifically, the rate of viscous creep closure will be much higher at the bed. However, when the terminus is near flotation, high basal water pressure will almost completely counteract this closing force. High basal water pressure is, thus, an essential condition for significant basal crevasse penetration (van der Veen, 1998a). Because our study focuses specifically on calving dynamics, we make the simplifying assumption that an efficient subglacial drainage system exists near the terminus and, thus, that there is negligible difference in basal water pressure within the region where calving may occur. With this assumption, basal water pressure is simply a function of sea level and bed elevation (van der Veen, 1998a):

$$P_w = -\rho_w g z \quad (5)$$

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where z is the z -coordinate which is negative below sea level.

Water pressure is essential for basal crevasse penetration, but it may also be significant in surface crevasses (Benn et al., 2007b). The process of “hydrofracturing” by water in surface crevasses is believed to have been a critical factor in the collapse of the Larsen B Ice Shelf (Scambos et al., 2003). However, while water in surface crevasses may be important, it is extremely difficult to quantify. The relationship between surface melt rate and crevasse water depth depends on the distribution, shape and depth of crevasses, melting and refreezing on crevasse walls, as well as potential drainage of water from crevasses into englacial, subglacial or proglacial water bodies. As such, it is currently impossible to estimate even an order of magnitude for crevasse water depth at Store Gletscher in summer. However, outside the 3 month summer melt season, surface crevasses must be assumed to be dry.

Modelling calving in a 2-D continuum model involves some implicit assumptions which may affect the accuracy of the calving criterion presented above. Firstly, the implementation of valley sidewall friction assumes that the calving terminus runs straight from one side of the valley to the other. However, Store’s terminus is usually arcuate in shape, with the centreline being further advanced in the fjord than the sidewalls. Thus, our implementation will overestimate lateral drag at the terminus. Secondly, by assuming a constant temperature of -10°C throughout the glacier, we neglect temperature dependent variations in viscosity and, thus, the stress field. Finally, Eq. (3) overestimates ice overburden pressure by assuming constant bulk density within the glacier. In fact, the presence of a crevasse field significantly reduces bulk density; this represents a positive feedback whereby the growth of a crevasse field reduces ice overburden pressure, leading to further crevasse deepening.

For the reasons outlined above, we expect our model to slightly underestimate the penetration of surface and basal crevasses near the present terminus position. As such, we apply a constant scaling factor of 1.075 to the effective stress term in Eq. (3). This scaling procedure is equivalent to the assumed presence of water in crevasses throughout the year in earlier work (Nick et al., 2010; Vieli and Nick, 2011). We note,

TCD

8, 3525–3561, 2014

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



meltwater discharge (van Angelen et al., 2012). Their results suggest an average melt rate across the entire face of 3.6 m d^{-1} in summer, with a local maximum at the base of the plume of 8 m d^{-1} . Because subglacial discharge is strongly influenced by surface runoff in summer months, we assume, for the sake of simplicity, that no submarine melting occurs in winter. If and when the floating tongue exists during the melt season, we apply a bottom melt rate of 1/10th of that applied on the vertical face, based on the “geometrical scale factor” proposed by Jenkins (2011).

In experiment 1, ice front melting is assumed to occur at a constant rate between June and August, both months included, as $> 90\%$ of all surface runoff in the Store catchment occurs over this period. In experiment 2, we investigate the effects of increasing summer melt rates by a factor of 1.5 and 2, and increasing its duration by 1 and 2 months, respectively.

3.5.2 Mélange backstress

We simulate the effect of mélange backstress by applying an external pressure on the calving terminus in addition to that exerted by the sea (Fig. 4). The applied pressure is similar to that found by Walter et al. (2012) from a force-balance study of Store, based on the observed speedup of the glacier following mélange collapse. Their result shows that the mélange yields a supporting pressure equivalent to a backstress of 30–60 kPa acting on the entire face of the terminus. In reality, this stress is applied only through the thickness of the mélange, a property not measured by Walter et al. (2012). To obtain a realistic forcing scenario at the calving front of our model, we convert Walter et al.’s backstress (σ_{fb}) into an equivalent mélange-glacier contact pressure:

$$\sigma_{sik} = \sigma_{fb} \frac{H_{term}}{H_{sik}} \quad (6)$$

where H_{term} and H_{sik} are the thicknesses of the glacier terminus and the mélange respectively.

4.2 Experiment 1

To attain a realistic “present day” simulation, we start by adding submarine melting, as described above, with rates up to 8 m d^{-1} from June to August. This forcing slightly increases the frequency and reduces the magnitude of calving events, though the overall terminus position varies only negligibly (Fig. 5a). Terminus velocity during the melt season is slightly suppressed compared with the melt-free simulation (Fig. 5b). This experiment suggests that neither seasonal variability in basal dynamics nor submarine melting explain the seasonal calving dynamics observed at Store. Only when the stabilizing effect of mélange buttressing is included does our model respond by significant frontal advance and retreat. Figure 6 shows the evolution of calving terminus position through time for each of the two seasonal forcings as well as the combined effect.

In our model, the formation of the mélange triggers the immediate formation of a floating ice tongue, which advances into the fjord. The terminus advances by 1300 m between February and May, while the mélange is present, and begins to retreat rapidly when the mélange collapses, irrespective of whether or not submarine melting is applied (Fig. 6). Figure 7 shows the evolution of the floating tongue through the mélange season. As the floating tongue advances, both the surface and basal crevasse fields are suppressed near the terminus. Note that the surface elevation rises as the floating tongue extends into the fjord, indicating that the dynamic regime near the grounding line leads to super-buoyancy which is only overcome once the floating tongue is long enough to exert a sufficient upward bending moment on the grounding line. Once significant upward bending is exerted, this is manifested as a suppression of surface crevasse field, clearly visible in Fig. 7.

When the mélange effect is combined with submarine melting, the collapse of the floating tongue is followed by further 250 m retreat beyond the stable terminus position at 113 km. After this retreat, the terminus slowly readvances through the melt season to 113 km, where it remains, calving periodically, until the mélange forms during the following winter.

Calving dynamics of Store

J. Todd and
P. Christoffersen

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

When mélange strength is reduced from 45 kPa to 33.75 kPa, the floating tongue does not begin to form until halfway through the mélange season. As a result, the maximum length of the tongue is reduced from 1.3 to 0.7 km (Fig. 9a). When mélange strength is further reduced to 22.5 kPa, no floating tongue forms in spring, though there remains a clear change in calving dynamics throughout the mélange season. These results suggest that any future climate related reduction in the strength of mélange may significantly affect the seasonality of Store.

Reducing the length of the mélange season from 0.32 to 0.21 years limits the length of the floating tongue to 0.8 km for the 45 kPa case (Fig. 9a). However, further reduction to 0.11 years has no effect on calving dynamics, and the floating tongue continues to advance for a month following mélange breakup. This is a surprising result, which suggests that the floating tongue is at least temporarily self-stabilising. In the 33.75 kPa case, when season length is reduced from 0.32 to 0.21, the floating tongue begins to advance slightly sooner and so the final length is slightly higher. However, in the 0.11 year, 33.75 kPa case, no floating tongue forms.

An increase in the duration of submarine melting, from 0.25 years to 0.33 and 0.41 years respectively (Fig. 9b), leads to more rapid collapse of the floating tongue, though in no case does the tongue collapse while rigid mélange is still present. As in experiment 1 (Fig. 6), submarine melting has an appreciable effect on the calving dynamics of the grounded terminus in late summer. As such, a longer submarine melt season leads to a longer period of larger, less frequent calving events and a retreat in average terminus position (Fig. 9b). The response of the modelled terminus to increasing melt magnitude appears somewhat stochastic. In some cases, the average terminus position during the melt season is further inland, but in other cases the reverse is true. Broadly speaking, however, the calving dynamics appear unaffected by increasing melt magnitude. In even the most severe “warming climate” scenario, with melt rate double present-day values and duration increased from 3 to 5 months, the modelled terminus remains stable.

5 Discussion

The results of our modelling experiments shed new light on marine terminating glacier dynamics and the calving mechanism. The calving dynamics of the modelled glacier vary significantly through the year (experiment 1, Fig. 6), from high frequency (8.7 days), low magnitude (~ 80 m) calving events when no seasonal forcing is applied, to complete cessation of calving during the *mélange* season, with rapid retreat following *mélange* collapse, and seemingly stochastic calving behaviour during the melt season. This behaviour is in good overall agreement with year-round observation of Store (N. Chauché, personal communication, 2014). Our model captures two important aspects of Store's behaviour. Seasonally, Store's terminus position is highly sensitive to external perturbation. However, on interannual timescales, Store's calving dynamics are stable, and the terminus position remains fairly constant (Howat et al., 2010).

In our model, the seasonal advance and retreat is specifically related to a floating tongue, which forms during winter in response to the buttressing effect of rigid *mélange* (Fig. 5b) and breaks apart once the buttressing effect of the *mélange* disappears. This finding provides theoretical understanding for the observed temporal correlation between *mélange* break up and frontal retreat at Store and other glaciers in the Uummannaq region (Howat et al., 2010), as well as Jakobshavn Isbræ (Amundson et al., 2010) farther south, and glaciers such as Kangerdlugssuaq and Daugaard-Jensen on the east coast (Seale et al., 2011). Our results from experiment 2 suggest that the estimate of Walter et al. (2012) of a *mélange* strength of 30–60 kPa is most likely correct, and that any future climate driven reduction in *mélange* strength or thickness could significantly impact the seasonal dynamics of Store (Fig. 9).

When we isolated the effect of submarine melting of the ice front (experiment 1, Fig. 5), we found a slight increase in calving frequency, an associated decrease in calving event size, and a slight dampening of the glacier's velocity response to calving events. However, the overall effect of submarine melting alone was minimal. Only when combined with *mélange* forcing was submarine melting capable of significantly affecting

TCD

8, 3525–3561, 2014

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



calving dynamics (Fig. 6). This suggests that some process during the mélange season preconditions the glacier for slight instability later in the season. Potentially, the upward bending associated with the formation of the floating tongue (Fig. 7) changes the glacier geometry near the grounding line such that it is more susceptible to the effect of undercutting by submarine melting.

Despite doubling melt rates and increasing melt duration by 66 % in experiment 2 (Fig. 9), the terminus of Store remained stable at 113 km, suggesting that there is no direct link between submarine undercutting and longer-term calving stability of the grounded terminus at present. This result contradicts previous work suggesting that undercutting of the terminus promotes calving (Motyka et al., 2003; Rignot et al., 2010) by intensifying extensional stresses near the terminus (O’Leary and Christoffersen, 2013). We propose, however, that this apparent contradiction is a feature specific to Store, due to the strong stabilising influence of topography.

The location of the terminus of Store coincides with a significant basal pinning point (Fig. 2), as well as a “bottleneck” in the fjord width (Fig. 1). The combined effect of these topographical features is to significantly affect the stress field and crevasse depth (Fig. 4). The suppression of crevasse penetration depth at the stoss side of the basal pinning point at the terminus exceeds the deepening of crevasses in response to undercutting of the ice front by submarine melting. As such, the latter alone cannot cause the front to retreat in this case. This suggests that so long as the melt rate is less than the rate of ice delivery to the front, the terminus position of Store will be relatively insensitive to the rate of ice front melting. Thus, the rate of iceberg production will be solely controlled by the velocity at the terminus. The topographic setting of Store explains why this glacier remained stable during a period when others in the same region experienced rapid retreat (Howat et al., 2010) and, more generally, why neighbouring glaciers are often observed to respond asynchronously to similar climate forcing (Moon et al., 2012).

Inland of Store’s stable frontal pinning point is a 28 km long overdeepening reaching 950 m below sea level (Fig. 2), which could make Store susceptible to sudden retreat,

Calving dynamics of Store

J. Todd and
P. Christoffersen

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

i.e. if the terminus becomes ungrounded from its current pinning point at 113 km. However, none of our climate forcing scenarios were able to trigger such a retreat which suggests that the current configuration of Store is stable and will most likely remain so in the near future.

As laid out above, our model is capable of reproducing the flow and seasonal calving dynamics of Store simply by perturbing the backstress exerted by mélange and the rate of submarine melting. Our model excludes the effect of water in surface crevasses, which may conceivably affect calving due to hydrofracture if water levels are high (Benn et al., 2007a). Although recent work included this effect (Nick et al., 2010), we ignore it because high resolution images captured in repeat surveys of Store with an unmanned aerial vehicle in July 2013 detected water in only a small number of surface crevasses near the terminus (Ryan et al., 2014). Although we cannot exclude the possibility that undetected water is contributing to crevasse penetration, it is not necessary to invoke this process to explain the observed behaviour of Store. This exclusion of hydrofracturing is a useful model simplification, as it is difficult and potentially impossible to accurately estimate the depth of water in crevasses. The latter would require knowledge of surface meltwater production as well as the number and size of surface crevasses, which is infeasible with the type of model used here.

Although our model captures the flow and seasonal calving dynamics of Store in a realistic manner, it is important to note that the outcome of our study is specifically limited to this glacier and that multiyear dynamics remain to be fully investigated. We use inverse methods to determine basal traction, rather than a hydrological model; this ensures that the flow field matches observations, allowing us to focus on processes at the terminus. However, prescribing basal traction means we are unable to investigate its interannual evolution in response to dynamic thinning, rising sea level or hydrological processes. The difficulty of implementing realistic hydrological routing in a flowline model suggests that only a 3-D model will be fully capable of representing these processes.

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



It is useful, at this point, to compare the development of time-evolving models for calving with recent developments in the implementation of grounding line dynamics. The lack of consistency of grounding line treatment in ice flow models was raised by Vieli (2005), and this issue has since received a great deal of attention from the ice-sheet modelling community. A comprehensive intercomparison study, MISIMIP (Pattyn et al., 2012), compared the ability of various 2-D ice flow models to simulate grounding line dynamics, before MISIMIP3d (Pattyn et al., 2013), did the same for 3-D models. Similarly, we hope that the 2-D model presented here will guide the future development of full 3-D time-evolving models for calving.

Finally, we note that in terms of accounting for the feedback between crevasse formation and bulk density and flow characteristics, a damage mechanics approach may prove useful. A counterpart study to this one by Krug et al. (2014) attempts to couple a damage model with a calving model for a synthetic glacier geometry.

6 Conclusions

Here we have presented results from a seasonally transient but interannually stable calving model of Store Gletscher in West Greenland. The calving numerics in our model differ from previous implementations of the crevasse depth criterion (Nick et al., 2010; Vieli and Nick, 2011; Cook et al., 2014) in that the balance of crevasse opening and closing forces is calculated through the entire thickness, not just at the boundaries, meaning that changes through depth are taken into account. In agreement with recent related work (Nick et al., 2010), we find that the inclusion of basal crevasses in the calving criterion is important. We propose the addition of a new divergence term to the Stokes equations, which is not only practical, but most likely essential for accurate simulation of glaciers in 2-D flowline models. We also find that the frequently assumed presence of water in surface crevasses is not necessary for seasonal calving dynamics.

We find that basal traction varies very little between winter and summer; basal lubrication by surface meltwater is therefore unlikely to play an important role in the

seasonal advance and retreat of the ice front. This does not imply, however, that calving and flow dynamics are not strongly coupled. Our results indicate a strong correlation between terminus position and velocity (Figs. 5 and 6). The deceleration which results from advance of the floating tongue is transmitted up to 30 km inland (Fig. 8). This finding supports previous studies which found that dynamic change at Helheim Glacier (Nick et al., 2009) and Jakobshavn Isbræ (Joughin et al., 2012) were triggered at the terminus.

A key outcome from this study is that the buttressing pressure from rigid mélange is principally responsible for observed seasonal advance. However, sensitivity analysis revealed that, in a warming climate, reduction in mélange strength or duration could prevent Store from advancing a floating tongue in winter. The model also indicates that submarine melting has only a limited effect on calving dynamics and that even large changes to melt rates in the future are unlikely to destabilize the terminus of Store. We propose that Store's highly stable terminus configuration is due to its topographic setting, being located at both a basal pinning point and a "bottleneck" in fjord width. We also find, however, that behind this basal pinning point, Store flows across a very large trough, reaching 950 m below sea level and extending 28 km inland from the current grounding line. This suggests that, were the terminus to be forced to retreat from its current pinning point, further retreat may be rapid and sudden, of a similar magnitude to that experienced by Jakobshavn Isbræ which resulted in a sustained increase of ice flux and contribution to sea level rise (Joughin et al., 2012).

The Supplement related to this article is available online at doi:10.5194/tcd-8-3525-2014-supplement.

Acknowledgements. This study was funded by the Natural Environment Research Council through a Ph.D. studentship (grant no. NE/K500884/1) to J. Todd and research grant (NE/K005871/1) to P. Christoffersen. We thank T. Zwinger, P. Råback and O. Gagliardini for

TCD

8, 3525–3561, 2014

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



help with the Elmer/Ice model, M. van den Broeke for providing RACMO climate data, A. Hubbard and J. Box for useful discussions related to Store Gletscher.

References

- Amundson, J. M., Truffer, M., Lüthi, M. P., Fahnestock, M., West, M., and Motyka, R. J.: Glacier, fjord, and seismic response to recent large calving events, Jakobshavn Isbræ, Greenland, *Geophys. Res. Lett.*, 35, L22501, doi:10.1029/2008GL035281, 2008. 3528
- Amundson, J. M., Fahnestock, M., Truffer, M., Brown, J., Lüthi, M. P., and Motyka, R. J.: Ice mélange dynamics and implications for terminus stability, Jakobshavn Isbræ, Greenland, *J. Geophys. Res.*, 115, F01005, doi:10.1029/2009JF001405, 2010. 3542
- Benn, D. I., Hulton, N. R. J., and Mottram, R. H.: “Calving laws”, “sliding laws” and the stability of tidewater glaciers, *Ann. Glaciol.*, 46, 123–130, 2007a. 3528, 3533, 3544
- Benn, D. I., Warren, C. R., and Mottram, R. H.: Calving processes and the dynamics of calving glaciers, *Earth-Sci. Rev.*, 82, 143–179, 2007b. 3528, 3533, 3535
- Box, J. E., Bromwich, D. H., Veenhuis, B. A., Bai, L., Stroeve, J. C., Rogers, J. C., Steffen, K., Haran, T., and Wang, S. H.: Greenland ice sheet surface mass balance variability (1988–2004) from calibrated polar MM5 output, *J. Climate*, 19, 2783–2800, 2006. 3527
- Chauché, N., Hubbard, A., Gascard, J.-C., Box, J. E., Bates, R., Koppes, M., Sole, A., and Patton, H.: Ocean properties, ice–ocean interactions, and calving front morphology at two major west Greenland glaciers, *The Cryosphere Discuss.*, 7, 5579–5611, doi:10.5194/tcd-7-5579-2013, 2013. 3529, 3536
- Christoffersen, P., Mugford, R. I., Heywood, K. J., Joughin, I., Dowdeswell, J. A., Syvitski, J. P. M., Luckman, A., and Benham, T. J.: Warming of waters in an East Greenland fjord prior to glacier retreat: mechanisms and connection to large-scale atmospheric conditions, *The Cryosphere*, 5, 701–714, doi:10.5194/tc-5-701-2011, 2011. 3527
- Cook, S., Rutt, I. C., Murray, T., Luckman, A., Zwinger, T., Selmes, N., Goldsack, A., and James, T. D.: Modelling environmental influences on calving at Helheim Glacier in eastern Greenland, *The Cryosphere*, 8, 827–841, doi:10.5194/tc-8-827-2014, 2014. 3533, 3545
- Cuffey, K. and Paterson, W.: *The Physics of Glaciers*, Butterworth-Heinemann, Oxford, UK, 4th Edn., 2010. 3531, 3532, 3533

TCO

8, 3525–3561, 2014

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Calving dynamics of Store

J. Todd and
P. Christoffersen

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Enderlin, E. M., Howat, I. M., Jeong, S., Noh, M.-J., van Angelen, J. H., and van den Broeke, M. R.: An improved mass budget for the Greenland Ice Sheet, *Geophys. Res. Lett.*, 41, 866–872, doi:10.1002/2013GL059010, 2014. 3526, 3527
- 5 Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Bamber, J. L., Box, J. E., and Bales, R. C.: Higher surface mass balance of the Greenland ice sheet revealed by high-resolution climate modeling, *Geophys. Res. Lett.*, 36, L12501, doi:10.1029/2009GL038110, 2009. 3529
- Favier, L., Gagliardini, O., Durand, G., and Zwinger, T.: A three-dimensional full Stokes model of the grounding line dynamics: effect of a pinning point beneath the ice shelf, *The Cryosphere*, 6, 101–112, doi:10.5194/tc-6-101-2012, 2012. 3532
- 10 Gagliardini, O., Durand, G., Zwinger, T., Hindmarsh, R. C. A., and Le Meur, E.: Coupling of ice-shelf melting and buttressing is a key process in ice-sheets dynamics, *Geophys. Res. Lett.*, 37, L14501, doi:10.1029/2010GL043334, 2010. 3532
- Gagliardini, O., Zwinger, T., Gillet-Chaulet, F., Durand, G., Favier, L., de Fleurian, B., Greve, R., Malinen, M., Martín, C., Råback, P., Ruokolainen, J., Sacchetti, M., Schäfer, M., Seddik, H., and Thies, J.: Capabilities and performance of Elmer/Ice, a new-generation ice sheet model, *Geosci. Model Dev.*, 6, 1299–1318, doi:10.5194/gmd-6-1299-2013, 2013. 3530
- 15 Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice sheet contribution to sea-level rise from a new-generation ice-sheet model, *The Cryosphere*, 6, 1561–1576, doi:10.5194/tc-6-1561-2012, 2012. 3531
- Gladstone, R. M., Lee, V., Rougier, J., Payne, A. J., Hellmer, H., Le Brocq, A., Shepherd, A., Edwards, T. L., Gregory, J., and Cornford, S. L.: Calibrated prediction of Pine Island Glacier retreat during the 21st and 22nd centuries with a coupled flowline model, *Earth Planet. Sc. Lett.*, 333–334, 191–199, doi:10.1016/j.epsl.2012.04.022, 2012. 3533
- 25 Hanna, E.: Runoff and mass balance of the Greenland ice sheet: 1958–2003, *J. Geophys. Res.*, 110, D13108, doi:10.1029/2004JD005641, 2005. 3526, 3527
- Holland, D. M., Thomas, R. H., de Young, B., Ribergaard, M. H., and Lyberth, B.: Acceleration of Jakobshavn Isbræ triggered by warm subsurface ocean waters, *Nat. Geosci.*, 1, 659–664, 2008. 3527
- 30 Howat, I., Joughin, I., Tulaczyk, S., and Gogineni, S.: Rapid retreat and acceleration of Helheim Glacier, east Greenland, *Geophys. Res. Lett.*, 32, L22502, doi:10.1029/2005GL024737, 2005. 3526

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Howat, I., Joughin, I., and Scambos, T.: Rapid changes in ice discharge from Greenland outlet glaciers, *Science*, 315, 1559–1561, doi:10.1126/science.1138478, 2007. 3526
- Howat, I. M., Box, J. E., Ahn, Y., Herrington, A., and McFadden, E. M.: Seasonal variability in the dynamics of marine-terminating outlet glaciers in Greenland, *J. Glaciol.*, 56, 601–613, 2010. 3528, 3529, 3530, 3538, 3540, 3542, 3543
- Howat, I. M., Negrete, A., and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface elevation datasets, *The Cryosphere Discuss.*, 8, 453–478, doi:10.5194/tcd-8-453-2014, 2014. 3531
- Jenkins, A.: Convection-driven melting near the grounding lines of ice shelves and tidewater glaciers, *J. Phys. Oceanogr.*, 41, 2279–2294, doi:10.1175/JPO-D-11-03.1, 2011. 3527, 3529, 3537
- Joughin, I., Howat, I. M., Fahnestock, M., Smith, B., Krabill, W., Alley, R. B., Stern, H., and Truffer, M.: Continued evolution of Jakobshavn Isbrae following its rapid speedup, *J. Geophys. Res.*, 113, F04006, doi:10.1029/2008JF001023, 2008. 3528
- Joughin, I., Smith, B., Howat, I., and Scambos, T.: MEaSURES Greenland Ice Velocity: Selected Glacier Site Velocity Maps from InSA R. Boulder, Colorado, USA: NASA DAAC at the National Snow and Ice Data Center, doi:10.5067/MEASURES/CRYOSPHERE/nsidc-0481.001, 2011. 3529, 3530, 3553
- Joughin, I., Smith, B. E., Howat, I. M., Floricioiu, D., Alley, R. B., Truffer, M., and Fahnestock, M.: Seasonal to decadal scale variations in the surface velocity of Jakobshavn Isbrae, Greenland: Observation and model-based analysis, *J. Geophys. Res.*, 117, F02030, doi:10.1029/2011JF002110, 2012. 3527, 3546
- Joughin, I., Smith, B. E., Shean, D. E., and Floricioiu, D.: Brief Communication: Further summer speedup of Jakobshavn Isbræ, *The Cryosphere*, 8, 209–214, doi:10.5194/tc-8-209-2014, 2014. 3527
- Khan, S. A., Wahr, J., Bevis, M., Velicogna, I., and Kendrick, E.: Spread of ice mass loss into northwest Greenland observed by GRACE and GPS, *Geophys. Res. Lett.*, 37, L06501, doi:10.1029/2010GL042460, 2010. 3526
- Krug, J., Weiss, J., Gagliardini, O., and Durand, G.: Combining damage and fracture mechanics to model calving, *The Cryosphere Discuss.*, 8, 1631–1671, doi:10.5194/tcd-8-1631-2014, 2014. 3545
- Luckman, A. and Murray, T.: Seasonal variation in velocity before retreat of Jakobshavn Isbræ, Greenland, *Geophys. Res. Lett.*, 32, L08501, doi:10.1029/2005GL022519, 2005. 3526

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- MacAyeal, D. R.: The basal stress distribution of Ice Stream E, Antarctica, inferred by control methods, *J. Geophys. Res.*, 97, 595–603, doi:10.1029/91JB02454, 1992. 3531
- McNabb, R. W., Hock, R., O’Neel, S., Rasmussen, L. A., Ahn, Y., Braun, M., Conway, H., Herreid, S., Joughin, I., Pfeffer, W. T., Smith, B. E., and Truffer, M.: Using surface velocities to calculate ice thickness and bed topography: a case study at Columbia Glacier, Alaska, USA, *J. Glaciol.*, 58, 1151–1164, 2012. 3532
- Moon, T., Joughin, I., Smith, B., and Howat, I.: 21st-century evolution of Greenland outlet glacier velocities, *Science*, 336, 576–578, doi:10.1126/science.1219985, 2012. 3543
- Motyka, R., Fahnestock, M., and Truffer, M.: Volume change of Jakobshavn Isbræ, West Greenland: 1985–1997–2007, *J. Glaciol.*, 56, 635–646, 2010. 3527
- Motyka, R. J., Hunter, L., Echelmeyer, K. A., and Connor, C.: Submarine melting at the terminus of a temperate tidewater glacier, LeConte Glacier, Alaska, USA, *Ann. Glaciol.*, 36, 57–65, doi:10.3189/172756403781816374, 2003. 3543
- Nick, F. M., Vieli, A., Howat, I. M., and Joughin, I.: Large-scale changes in Greenland outlet glacier dynamics triggered at the terminus, *Nat. Geosci.*, 2, 110–114, doi:10.1038/ngeo394, 2009. 3528, 3546
- Nick, F. M., Van der Veen, C. J., Vieli, A., and Benn, D. I.: A physically based calving model applied to marine outlet glaciers and implications for the glacier dynamics, *J. Glaciol.*, 56, 781–794, 2010. 3528, 3533, 3534, 3535, 3544, 3545
- Nye, J. F.: The distribution of stress and velocity in glaciers and ice-sheets, *P. Roy. Soc. A-Math. Phys.*, 239, 113–133, doi:10.1098/rspa.1957.0026, 1957. 3528, 3533
- O’Leary, M. and Christoffersen, P.: Calving on tidewater glaciers amplified by submarine frontal melting, *The Cryosphere*, 7, 119–128, doi:10.5194/tc-7-119-2013, 2013. 3528, 3543
- Otero, J., Navarro, F. J., Martin, C., Cuadrado, M. L., and Corcuera, M. I.: A three-dimensional calving model: numerical experiments on Johnsons Glacier, Livingston Island, Antarctica, *J. Glaciol.*, 56, 200–214, doi:10.3189/002214310791968539, 2010. 3533
- Pattyn, F., Schoof, C., Perichon, L., Hindmarsh, R. C. A., Bueller, E., de Fleurian, B., Durand, G., Gagliardini, O., Gladstone, R., Goldberg, D., Gudmundsson, G. H., Huybrechts, P., Lee, V., Nick, F. M., Payne, A. J., Pollard, D., Rybak, O., Saito, F., and Vieli, A.: Results of the Marine Ice Sheet Model Intercomparison Project, MISMP, *The Cryosphere*, 6, 573–588, doi:10.5194/tc-6-573-2012, 2012. 3545
- Pattyn, F., Perichon, L., Durand, G., Favier, L., Gagliardini, O., Hindmarsh, R. C., Zwinger, T., Albrecht, T., Cornford, S., Docquier, D., Fürst, J. J., Goldberg, D., Gudmundsson, G. H.,

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Humbert, A., Hütten, M., Huybrechts, P., Jouvét, G., Kleiner, T., Larour, E., Martin, D., Morlighem, M., Payne, A. J., Pollard, D., Rückamp, M., Rybak, O., Seroussi, H., Thoma, M., and Wilkens, N.: Grounding-line migration in plan-view marine ice-sheet models: results of the ice2sea MISMP3d intercomparison, *J. Glaciol.*, 59, 410–422, 2013. 3545

5 Raymond, C.: Shear margins in glaciers and ice sheets, *J. Glaciol.*, 42, 90–102, 1996. 3532

Rignot, E. and Kanagaratnam, P.: Changes in the velocity structure of the Greenland Ice Sheet, *Science*, 311, 986–990, doi:10.1126/science.1121381, 2006. 3526

Rignot, E., Koppes, M., and Velicogna, I.: Rapid submarine melting of the calving faces of West Greenland glaciers, *Nat. Geosci.*, 3, 187–191, doi:10.1038/ngeo765, 2010. 3536, 3543

10 Ryan, J. C., Hubbard, A. L., Todd, J., Carr, J. R., Box, J. E., Christoffersen, P., Holt, T. O., and Snooke, N.: Repeat UAV photogrammetry to assess calving front dynamics at a large outlet glacier draining the Greenland Ice Sheet, *The Cryosphere Discuss.*, 8, 2243–2275, doi:10.5194/tcd-8-2243-2014, 2014. 3544

Scambos, T., Hulbe, C., and Fahnestock, M.: Climate-induced ice shelf disintegration in the Antarctic Peninsula, *Antarct. Res. Ser.*, 79, 79–92, 2003. 3535

15 Seale, A., Christoffersen, P., Mugford, R. I., and O’Leary, M.: Ocean forcing of the Greenland Ice Sheet: Calving fronts and patterns of retreat identified by automatic satellite monitoring of eastern outlet glaciers, *J. Geophys. Res.*, 116, F03013, doi:10.1029/2010JF001847, 2011. 3527, 3542

20 Sohn, H.-G., Jezek, K. C., and van der Veen, C. J.: Jakobshavn Glacier, west Greenland: 30 years of spaceborne observations, *Geophys. Res. Lett.*, 25, 2699–2702, doi:10.1029/98GL01973, 1998. 3528

Stocker, D.: Climate change 2013: the physical science basis, Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Summary for Policymakers, IPCC, 2013. 3527

25 Straneo, F., Hamilton, G. S., Sutherland, D. A., Stearns, L. A., Davidson, F., Hammill, M. O., Stenson, G. B., and Rosing-Asvid, A.: Rapid circulation of warm subtropical waters in a major glacial fjord in East Greenland, *Nat. Geosci.*, 3, 182–186, doi:10.1038/ngeo764, 2010. 3527

30 Thomas, R. H., Abdalati, W., Frederick, E., Krabill, W. B., Manizade, S., and Steffen, K.: Investigation of surface melting and dynamic thinning on Jakobshavn Isbrae, Greenland, *J. Glaciol.*, 49, 231–239, 2003. 3532

van Angelen, J. H., Lenaerts, J. T. M., Lhermitte, S., Fettweis, X., Kuipers Munneke, P., van den Broeke, M. R., van Meijaard, E., and Smeets, C. J. P. P.: Sensitivity of Greenland

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice Sheet surface mass balance to surface albedo parameterization: a study with a regional climate model, *The Cryosphere*, 6, 1175–1186, doi:10.5194/tc-6-1175-2012, 2012. 3537

van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E., van de Berg, W. J., van Meijgaard, E., Velicogna, I., and Wouters, B.: Partitioning recent Greenland mass loss., *Science*, 326, 984–986, doi:10.1126/science.1178176, 2009. 3526, 3527

van der Veen, C.: Fracture mechanics approach to penetration of bottom crevasses on glaciers, *Cold Reg. Sci. Technol.*, 27, 213–223, 1998a. 3528, 3533, 3534

van der Veen, C.: Fracture mechanics approach to penetration of surface crevasses on glaciers, *Cold Reg. Sci. Technol.*, 27, 31–47, 1998b. 3528, 3533

Vieli, A.: Assessing the ability of numerical ice sheet models to simulate grounding line migration, *J. Geophys. Res.*, 110, F01003, doi:10.1029/2004JF000202, 2005. 3545

Vieli, A. and Nick, F.: Understanding and modelling rapid dynamic changes of tidewater outlet glaciers: issues and implications, *Surv. Geophys.*, 32, 437–458, 2011. 3535, 3545

Walter, J. I., Jason, E., Tulaczyk, S., Brodsky, E. E., Howat, I. M., Yushin, A. H. N., and Brown, A.: Oceanic mechanical forcing of a marine-terminating Greenland glacier, *Ann. Glaciol.*, 53, 181–192, 2012. 3528, 3537, 3538, 3540, 3542

Weidick, A.: Greenland, with a section on Landsat images of Greenland, edited by: Williams Jr., R. S. and Ferrigno, J. G., *Satellite image atlas of glaciers of the world*, U.S. Geological Survey Professional Paper 1386-C, 141 pp., ISBN 0-607-71454-9, available at: <http://pubs.usgs.gov/prof/p1386c> (last access: 15 May 2014), 1995. 3528

Xu, Y., Rignot, E., Fenty, I., Menemenlis, D., and Flexas, M. M.: Subaqueous melting of Store Glacier, west Greenland from three-dimensional, high-resolution numerical modeling and ocean observations, *Geophys. Res. Lett.*, 40, 4648–4653, doi:10.1002/grl.50825, 2013. 3528, 3529, 3536

Zwally, H. J., Abdalati, W., Herring, T., Larson, K., Saba, J., and Steffen, K.: Surface melt-induced acceleration of Greenland ice-sheet flow, *Science*, 297, 218–22, doi:10.1126/science.1072708, 2002. 3527




Figure 1. Store Gletscher in Ikerasak Fjord, Greenland. Colour scale shows summer velocity (m a^{-1}) from the MEaSUREs program (Joughin et al., 2011). Yellow line indicates the flowline used in this study, and the green star indicates the location of the main proglacial plume forming where subglacial water is discharged into the fjord.

Calving dynamics of Store

J. Todd and P. Christoffersen

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 3. The terminus of the flowline mesh of Store Gletscher. White line indicates the net stress (σ_n) zero contour for both surface and basal crevasses. Blue line indicates sea level. Net stress (MPa) is > 0 where crevasses exist and < 0 in solid and unfractured ice. Calving occurs in the model when surface and basal crevasses meet.

Calving dynamics of Store

J. Todd and P. Christoffersen

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Calving dynamics of Store


J. Todd and
P. Christoffersen

Figure 4. Proximal and distal processes affecting calving. **(a)** Varying basal friction (τ_b) affects the stress field in the glacier. **(b)** Changing fjord water temperatures and, more importantly, subglacial water flux affect the rate of submarine melting of the calving face and floating tongue (when present). **(c)** The seasonal formation of mélangé provides a buttressing force which suppresses surface crevasse depth and, thus, calving. **(d)** Surface melt water in crevasses causes hydrofracturing, which acts to deepen surface crevasses. **(e)** Glacier geometry exerts a strong influence on crevasse field depth. Compressional forces on the stoss side of Store's pinning point suppress the depth of crevasses, while rapid loss of basal traction in the lee side deepen them.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 5. Plots showing variations in terminus position **(a)** and velocity **(b)**, over the course of a year for baseline model run (blue line) and run with submarine melting applied (red line). Red shading indicates melt season. The saw-toothed pattern in both panels is a result of calving.

Calving dynamics of Store

J. Todd and
P. Christoffersen

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Calving dynamics of Store


J. Todd and
P. Christoffersen

Figure 6. Plots showing changes in calving terminus position **(a)** and velocity **(b)** during a five year period within a 40 year long stable simulation, with coloured solid lines illustrating the effect of four different combinations of melting and ice mélange perturbation. Blue and red shade indicates melté and melt season respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Calving dynamics of Store


J. Todd and
P. Christoffersen

Figure 7. Sequential profiles of Store Gletscher during advance of its calving terminus due to mélange backstress. As the floating tongue advances from the grounding line (marked GL), it rises upwards due to buoyant forces, which also act to close surface crevasses near the grounding line. This indicates that flow dynamics at the grounding line are forcing the terminus below flotation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Calving dynamics of Store


J. Todd and
P. Christoffersen

Figure 9. Plots showing terminus position through one year for varying mélange season length **(a)** and melt season length **(b)**. Durations of mélange and melt season are indicated by blue and red shading (and panel title), respectively, varying from left to right, with colour indicating the magnitude of the perturbation. Moving from left to right panels, and moving from blue, through green, to red line colours, indicate progressively more severe “warming climate” scenarios.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

