

Improved Arctic sea ice thickness
projections using
bias corrected CMIP5
simulations

N. Melia et al.

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia¹, K. Haines², and E. Hawkins³

¹Department of Meteorology, University of Reading, Reading, UK

²National Centre for Earth Observation, Department of Meteorology, University of Reading, Reading, UK

³NCAS-Climate, Department of Meteorology, University of Reading, Reading, UK

Received: 26 June 2015 – Accepted: 29 June 2015 – Published: 22 July 2015

Correspondence to: N. Melia (n.melia@pgr.reading.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Projections of Arctic sea ice thickness (SIT) have the potential to inform stakeholders about accessibility to the region, but are currently rather uncertain. The latest suite of CMIP5 Global Climate Models (GCMs) produce a wide range of simulated SIT in the historical period (1979–2014) and exhibit various spatial and temporal biases when compared with the Pan-Arctic Ice Ocean Modelling and Assimilation System (PIOMAS) sea ice reanalysis. We present a new method to constrain such GCM simulations of SIT to narrow projection uncertainty via a statistical bias correction technique. The bias correction successfully constrains the spatial SIT distribution and temporal variability in the CMIP5 projections whilst retaining the climatic fluctuations from individual ensemble members. The bias correction acts to reduce the uncertainty in projections of SIT and reveals the significant contributions of sea ice internal variability in the first half of the century and of scenario uncertainty from mid-century onwards. The projected date of ice-free conditions in the Arctic under the RCP8.5 high emission scenario occurs in the 2050s, which is a decade earlier than without the bias correction, with potentially significant implications for stakeholders in the Arctic such as the shipping industry. The bias correction methodology developed could be similarly applied to other variables to narrow uncertainty in climate projections more generally.

1 Introduction

Global Climate Models (GCMs) are the primary tool for making climate predictions on seasonal to decadal time scales, and climate projections over the next century (Flato et al., 2013). In a warming climate, changes to sea ice thickness (SIT) are expected to lead to significant implications for polar regions and beyond. A reduction in SIT will likely open up the Arctic Ocean to economic diversification including new marine shipping routes (Smith and Stephenson, 2013) and extraction of natural resources, as well as changes to the Arctic ecosystem and potential links to mid-latitude weather

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Francis and Vavrus, 2012). Many of these economic opportunities may rely on SIT, but current projections have considerable uncertainty.

The GCMs from the Coupled Model Intercomparison Project, phase 5 (CMIP5) (Taylor et al., 2012) exhibit a large range in sea ice volume (SIV), spatial SIT distribution, and temporal SIT variability under present day forcing conditions (e.g. Blanchard-Wrigglesworth and Bitz, 2014). For September sea ice extent, Swart et al. (2015) showed the uncertainty in CMIP5 projections over the next few decades is dominated by these differences between models, termed model uncertainty by Hawkins and Sutton (2009, 2011). Uncertainty in climate projections arises from three distinct sources: (1) model uncertainty, (2) internal variability, and (3) scenario uncertainty, as discussed by Hawkins and Sutton (2009, 2011) for temperature and precipitation respectively. In contrast to projections of temperature where the anomalies are often used, the absolute value of SIT is important – for example, ships have critical SIT thresholds above which their use is not possible (Stephenson et al., 2013). Since absolute values are used, model bias makes a contribution to model uncertainty.

Bias correction (BC) of GCM simulations has the potential to reduce the model uncertainty and hence increase confidence in near term climate projections. The importance of BC in impact based climate change studies was described in a special report of the IPCC (Seneviratne et al., 2012), but BC has not previously been applied to projections of sea ice. There are many different types of proposed BC techniques, (e.g. Christensen et al., 2008; Ho et al., 2011; Vrac and Friederichs, 2014; Watanabe et al., 2012, and references therein), which have mainly been applied to temperature and precipitation. However, these existing methods need refining for sea ice as SIT is a particularly challenging variable. This is due to its positive semi-definite nature, and the spatial and temporal occurrence of zeros, in observations and projections of SIT.

This study addresses the development of a new BC technique that constrains both the mean and variance of SIT in GCMs to an estimate of the observed statistics. It is important to correct the mean as this corrects the spatial SIT distribution. Variability in SIT also has a significant impact on the range of regional ice-free dates, something of

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



great interest to stakeholders, and the CMIP5 GCMs exhibit a wide range in their SIT variability. The study also uses multiple ensemble members from the same model when performing the BC, something that is often not utilised in other studies. This is important as it enables an assessment of the role of internal variability in future projections to be made. The techniques described in this paper are not limited to SIT, and would work for many climate variables. The exact implementation used in this study should also be calibrated to the user's needs based on factors such as the length of reliable observations and number of ensemble members.

In this paper we use the Pan-Arctic Ice Ocean Modelling and Assimilation System (PIOMAS) (Zhang and Rothrock, 2003) as an observationally based estimate of recent SIT, along with climate projections from a subset of six GCMs from the CMIP5 archive (Sect. 2). We first test the performance of increasingly complex BC approaches in a toy model environment (Sect. 3) and then apply our favoured method to the subset of CMIP5 GCMs in Sect. 4. We test the BC method by splitting the historical PIOMAS data, and then explore how the uncertainty in SIT projections is reduced using these techniques (Sect. 4) and summarise and discuss the results in Sect. 5.

2 Climate simulations and observations

2.1 PIOMAS

For an observationally based estimate of SIT, we use the PIOMAS reanalysis. PIOMAS is a coupled ice–ocean model that is forced with the National Centers for Environmental Prediction (NCEP) atmospheric reanalysis, and assimilates satellite observed sea ice concentration (Lindsay and Zhang, 2006) and sea surface temperature (Schweiger et al., 2011). It does not however assimilate sea ice thickness (SIT), although this has been attempted using the NASA Operation IceBridge and SIZONet campaigns of 2012 (Lindsay et al., 2012).

TCD

9, 3821–3857, 2015

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We choose PIOMAS to represent observations of SIT as satellite observations are limited in their spatial and temporal range. For example, data from ICESat are only available between October and March 2003–2008 (Kwok et al., 2009). More recently Cryosat-2 (Laxon et al., 2013) has started producing real-time SIT datasets but only for the non-summer months. This is also not ideal as it is the summer months when the ice is thinnest that are most relevant for potential economic activity. The spatial consistency, temporal length and completeness of the data are important considerations when computing climatological means and variances as the longest time series possible is needed to validate the statistics. It is for this reason primarily that PIOMAS has been chosen to represent observations in this study. Several studies (e.g. Laxon et al., 2013; Schweiger et al., 2011; Lindsay and Zhang, 2006; Stroeve et al., 2014) have compared PIOMAS to satellite and in situ observations and models and find it a suitable estimate of observed SIT. PIOMAS is also deemed realistic enough to initialise numerical models for seasonal forecasts e.g., the Sea Ice Outlook (Blanchard-Wrigglesworth and Bitz, 2014) where the accuracy of the initial conditions is vital.

Figure 1 shows the mean September SIT and temporal standard deviation (SD) after linear detrending for PIOMAS over the satellite era (1979–2014). The thickest ice is located north of the Canadian archipelago and Greenland. In the heart of the Canadian archipelago, ice thickness is up to 1.5 m, in the central Arctic it is about two meters, and it is between zero and one meter along the north Russian coast. The SIT is most variable around the edge of the ice pack and especially near land. An effective BC should ensure that the simulations replicate these patterns of mean SIT and SD over this recent period.

2.2 Global climate models

This paper utilises a subset of six GCMs from CMIP5. Since a large part of this work assesses SIT variability, it is necessary for each GCM to have multiple ensemble simulations in the historical period and for each of the representative concentration pathways (RCPs) 2.6, 4.5 and 8.5 for future scenarios (Van Vuuren et al., 2011). In addition, the

GCM mean spring thickness must fall within the 10th and 90th percentile of PIOMAS (Stroeve et al., 2014), have a reasonable spatial resolution, and at least one ocean channel through the Canadian archipelago. The six GCMs that comprise this CMIP5 subset are listed in Table 1.

For the CMIP5 subset the historical simulations are used for the period 1979–2005. In most of the analysis for the period post-2005 the RCP8.5 scenario is used, which ramps up the amount of greenhouse gases to have a cumulative effect of increasing the direct radiative forcing by 8.5 W m^{-2} (approximately 1370 ppm CO_2 equivalent) by 2100 (Van Vuuren et al., 2011). The impact of other scenarios is assessed later in the analysis. Figure 2 shows the 1979–2014 ensemble-mean September SIT for the CMIP5 subset, highlighting the considerable differences between the model simulations, and indicating that model bias is likely to be the dominant uncertainty in near-term projections.

The aim of the SIT BC outlined in this paper is to correct the mean and variance in the CMIP5 subset shown in Fig. 2 to the observed statistics. Although this should improve short-term predictions, a caveat to this approach is that there is only one realisation of the past, and we have to assume that the relatively short period over which we have observations (36 years) captures a representative sample of the behaviour we expect from the climate system. In the short term, this is probably a reasonable assumption, as the GCMs will not have evolved far from their corrected state of the recent past; this assumption is explored further in Sect. 4.

3 Bias correction methodology

Bias correction methods effectively aim to reduce model uncertainty by constraining GCMs to observations. There are two components to model uncertainty: the overall mean difference (or bias), and differences in the amplitude of response to specified forcings. We have deliberately chosen not to try and correct the simulated ice loss trend to that which is currently observed. Our reasoning is to keep this as prescribed by the

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



different models because the response of the SIT to future warming is unknown and GCMs are designed to give an estimate of this. It is also doubtful how well the current trend can be determined from 36 years of data given the high noise to signal ratio for trends, especially on grid point scales. It is also unclear how much of the recent ice loss seen in the observations can be attributed to changes in external forcing as opposed to internal variability (e.g. Day et al., 2012; Kay et al., 2011; Swart et al., 2015).

To test the performance of different possible BC methods a “toy model” was used as proxy ensemble timeseries (representing SIT at a single grid point for the same month each year for the period 1979–2100). The timeseries are shown in Fig. 3a for a high mean–high variance model (blue) and a low mean–low variance model (red), where the black line shows the “truth” observations with one realisation over the historical period only. The time series were all produced using a first order auto-regressive (AR1) model imposed on a declining linear trend, with five separate model ensemble members (thin coloured lines) and the thick lines representing the ensemble means. The statistics in all the legends are calculated over the observation window (1979–2014). A successful BC method should transform the individual ensemble members (thin red and blue lines) to match the mean and variance of the observations (black line), producing matched statistics. We test various approaches for such a bias correction. The mathematical notation for the following equations is in Table 2.

3.1 Additive correction

A basic additive correction, which has previously been used for temperature projections, is shown in Fig. 3b. This approach simply corrects the mean by subtracting the difference between the historical model ensemble-mean time-mean, $\langle \overline{M_h} \rangle$, and observation time mean, $\overline{O_h}$, from each of the model ensemble members, M .

$$\text{Additive corrected thickness} = M - \left(\langle \overline{M_h} \rangle - \overline{O_h} \right) \quad (1)$$

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



However, as the low ice model is adjusted up by the addition of a constant, it equilibrates at a positive value in the future rather than zero. Likewise the high ice model equilibrates at negative values. Neither of these properties are sensible.

This study makes use of multiple ensemble members from the same model, raising the question of how to treat ensemble member statistics when calculating a particular GCM's bias. For calculating the mean SIT, each GCM's ensemble mean is used because it is the GCM's mean bias that we wish to correct. This is important because a particular ensemble member's deviation from the ensemble mean is retained; it allows an individual ensemble member's time mean to be different to the observations over the historical period, but not the ensemble mean. The treatment of ensemble members for the SD calculation is described in Sect. 3.4.

3.2 Multiplicative correction

If a multiplicative correction is used (Fig. 3c), where the ratio of the observed time mean and model ensemble-mean time-mean, $\overline{O_h}/\langle\overline{M_h}\rangle$, is multiplied as a factor to the model ensemble members, M , then the corrected thickness is:

$$\text{Multiplicative corrected thickness} = M \frac{\overline{O_h}}{\langle\overline{M_h}\rangle} \quad (2)$$

Multiplicative methods effectively preserve the future zero ice year, which is potentially an important value for a wide range of stakeholders. However, when applied as above this approach has the undesired effect of distorting the variances by the same factor as the mean correction, as visible in Fig. 3c.

3.3 Mean multiplicative correction

To avoid altering the variances, the mean multiplicative correction can be introduced (Fig. 3d), where the multiplicative mean correction, $\overline{O_h}/\langle\overline{M_h}\rangle$, is applied only to the

11 year-centred running-mean ensemble-mean, $\langle \tilde{M} \rangle$. This corrects the model mean evolution without corrupting the sub-decadal variance as $\langle \tilde{M} \rangle$ is smoothed. The model anomalies for each ensemble member, $M - \langle \tilde{M} \rangle$, are then added back to the corrected mean evolution:

$$5 \quad \text{Mean multiplicative corrected thickness} = \left(M - \langle \tilde{M} \rangle \right) + \langle \tilde{M} \rangle \frac{\overline{O_h}}{\langle \tilde{M}_h \rangle} \quad (3)$$

This works to correct the mean SIT and does not suffer from any peculiarities of the previous two methods. The model variance now remains unchanged but the approach opens up the possibility of correcting the variance towards that observed in the historical period. Note that by using the ensemble mean, $\langle \tilde{M}_h \rangle$, for all these corrections we ensure that each ensemble member is corrected in the same way, thus preserving certain ensemble properties into the future.

3.4 Mean and variance correction

The GCMs from CMIP5 show a large range in sea ice variance, and the magnitude of these variations is a significant factor determining when regions of the Arctic may first become accessible (when one ensemble member may first become ice free). Therefore a variance correction is incorporated into Eq. (3) by taking the ratio of the temporal standard deviation of the detrended observations, $\sigma_{\hat{O}_h}$, to the square root of the ensemble mean of the variance of the detrended model ensembles, $\langle \sigma_{\tilde{M}_h} \rangle$ (detrended mean ensemble SD), over the historical period. The detrending in the models is calculated using each model's ensemble mean linear trend. This has some similarities to the approach of Ho et al. (2011) in application to temperature projections for Europe. Also see Appendix A for some further discussion of the choices made.

To incorporate the variance correction, the mean multiplicative correction (Eq. 3) is first de-trended, the variance correction applied, and the trend re-applied. This creates

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the Mean And Variance Correction (MAVRIC), shown in Eq. (4):

$$\text{MAVRIC} = (M - \langle \tilde{M} \rangle) \frac{\sigma_{\tilde{O}_h}}{\langle \sigma_{\tilde{M}_h} \rangle} + \langle \tilde{M} \rangle \frac{\overline{O}_h}{\langle \overline{M}_h \rangle} \quad (4)$$

Figure 3e shows the MAVRIC does a near perfect job of correcting both the mean and variance to the observed statistics while still retaining the individual ensemble members' own climate fluctuations, but fractionally scaled by the variance ratio.

4 Bias corrected sea ice thickness projections

Figure 3e illustrates that the MAVRIC successfully corrects the mean and variance in a toy model environment. Before proceeding to investigate the impact of the MAVRIC on SIT projections it is prudent to test whether the MAVRIC can improve GCM performance by validating with real observations. We use CSIRO-Mk3.6.0 (CSIRO) as the GCM to test. CSIRO generally has too much ice and too little variability and is a CMIP5 outlier model with regards to SIT (Stroeve et al., 2014). However, CSIRO benefits from having 10 ensemble members, increasing the robustness of the statistics. For these two reasons, it is considered a thorough test of the MAVRIC's performance within a real GCM.

The test uses a data denial method where we train the MAVRIC on a subset of PI-OMAS observations, 1979–1999, termed the calibration window. From this we examine how the MAVRIC predicts the observations for 2000–2014, termed the validation window. A limitation with this method is the length of observations: the period over which the MAVRIC calibration takes place must be long enough to capture a robust measure of the observed statistics. The validation period must also be long enough to be able to draw robust conclusions. It is not clear whether either the 21 year calibration or the 15 year validation windows are long enough for robust method calibration and results verification, but we are limited by the data available.

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Figure 4 shows the performance of the MAVRIC at three grid points for September. The raw CSIRO ensembles (grey) are bias corrected via the MAVRIC using the PIOMAS observations (black) over the calibration window, producing the MAVRIC corrected ensembles (green) for the validation window. If the MAVRIC can produce plausible predictions, the characteristics of PIOMAS should be indistinguishable from individual corrected ensemble members in the validation window. It is clear from the validation beanplots (right), that the distribution from the corrected ensembles resembles PIOMAS much more closely than the raw distribution, e.g. non-zero probability of zero ice. We do not expect the distribution from PIOMAS to match the corrected distribution perfectly as PIOMAS only has one realisation (15 data points) while CSIRO has 10 realisations. We can tentatively accept that this test demonstrates the validity of the MAVRIC approach.

In the following sections the MAVRIC is applied to the CMIP5 subset of six GCMs used in this study (Table 1). PIOMAS estimates of Arctic SIT are available from 1979–2014. This 36 year window is the period over which statistics are calculated in the observations, and in the CMIP5 subset (using historical runs for 1979–2005 and RCP8.5 for 2006–2014). Each model, month, and grid point has its own specific correction which is applied to all years (1979–2100). However, separate ensemble members from the same GCM are treated with the same correction, as we wish to correct the model bias and retain the ensemble spread. Results are shown for September, initially only for CSIRO and later for all six models combined to form the “CMIP5 subset” used for this study.

4.1 Temporal perspective example

Figure 5 shows the impact of the MAVRIC in September in CSIRO at the same three grid points as Fig. 4 but for the entire calibration window (1979–2014). The East Siberian Sea in CSIRO has about double the SIT and half the SD of PIOMAS (Fig. 5a). The correction therefore reduces the mean SIT whilst increasing the variance. This brings forward the range of first year ice-free conditions (the first occurrence in each

ensemble member of a SIT below 0.15 m) from after 2100 to 1981–2032. Ice age (and hence strength) correlates well to ice thickness (Maslanik et al., 2007), and values below 0.15 m correspond to young and grey ice categories, and operations in this ice regime require no specific ice strengthening of vessels (Transport Canada, 1998). Similarly in the Beaufort Sea (Fig. 5b) the SD needs to be almost tripled, and the correction results in the first ice-free year coming over 100 years earlier. In the Fram Strait (Fig. 5c) CSIRO and PIOMAS have similar SIT requiring only a small mean adjustment, however CSIRO requires a big increase in variance. The MAVRIC moves the first possible ice-free date about 30 years earlier and increases the ensemble uncertainty range from 32 to 63 years. It is worth noting that the dominant cause of this shift to earlier ice free date at this location is due to the variance correction term in the MAVRIC rather than the mean correction term. This highlights the importance of correcting the variance in addition to the mean. Figure 5 demonstrates that the MAVRIC can lead to projections that look significantly more like reality in the historical period and have an impact on regional ice-free projections.

4.2 Historical spatial perspective

In addition to examining the MAVRIC in a temporal sense, it is important to evaluate the results spatially to see where the MAVRIC is having the most effect and if it works at all locations. Figures 2 and 6 show that the mean September SIT distribution is very different in HadGEM2-ES and CSIRO. After the MAVRIC has been applied, the mean SIT fields are almost identical for the historical period (Fig. 6). It is important to note there are still differences when considering individual years and ensemble members i.e. the year-to-year variability and ensemble spread is preserved (although adjusted by the MAVRIC).

Figure 6 also shows the SD before and after the MAVRIC. The SD shown is the detrended mean ensemble SD as before. CSIRO has too low variability in the majority of locations although correctly places the maximum SD near the edges of the ice pack similarly to PIOMAS. HadGEM2-ES exhibits about the same magnitude of variability as

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the observations but the variability is too high in the centre of the ice pack and too low at the edges. After the correction the SD fields in both GCMs now look more similar to each other with the highest variability located at the edge of the ice pack and at coastal locations. They are now also both similar to the estimate from PIOMAS (Fig. 1).

4.3 CMIP5 subset multi-model sea ice thickness projections

The bias corrected SIT from each GCM can be brought together to form the multi-model mean CMIP5 subset, computed using three ensemble members (the maximum available across all models) from each of the six GCMs for the historical and future decadal periods (Fig. 7). It is remarkable how the raw multi-model mean product for the historical period is not too different from PIOMAS in Fig. 1, showing that the location and magnitude of model biases cancel out to a considerable degree, at least with this subset of models. Given this result it is not so surprising that the raw and corrected fields are fairly similar for the future projections also.

Nevertheless, even in this multi-model multi-ensemble framework the MAVRIC is still making some discernible differences. These differences are most apparent in the Canadian archipelago and the Russian Arctic seas, where the correction leads to a reduction in SIT of approximately 1 m in both regions. Both the raw and bias corrected fields predict a SIT loss of about 0.25 m per decade.

The fact that the MAVRIC is still making a significant difference on the regional scale is critical, e.g. for ship route availability. Currently studies that assess the future opening of Arctic shipping routes, which critically depend on the absolute value of SIT, do not yet account for such factors and will need to be reassessed.

4.4 Sources of uncertainty in projections of sea ice thickness

The uncertainty in climate projections can be partitioned into three distinct sources: (1) model uncertainty: for the same radiative forcing different models simulate different mean distributions and temporal changes. (2) Internal variability: the natural fluctua-

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tions of the climate present with or without any anthropogenic induced changes to radiative forcing. (3) Scenario uncertainty: uncertainty in future radiative forcing resulting from unknown future emissions. Hawkins and Sutton (2009, 2011) assessed these sources of uncertainty in global and regional temperature and precipitation projections, and here we quantify the sources of uncertainty in SIT, utilising the CMIP5 subset multi-model ensemble. Crucially we use the absolute values of SIT rather than considering anomalies as is often done for other climate variables. The methodology for partitioning these sources of uncertainty is detailed in Appendix B.

The MAVRIC method outlined in this study acts to eliminate the model bias (and hence reduce the uncertainty) in the MAVRIC calibration period (1979–2014). After this period the model uncertainty grows due to the GCM's differing responses to changes in external forcing. The sources of uncertainty for SIT for the decade 2015–2024, immediately following the MAVRIC calibration period, are shown in Fig. 8. The total uncertainty in the corrected CMIP5 subset is strikingly lower than in the raw CMIP5 subset. Closer analysis reveals that this is due to the substantial reduction in model uncertainty owing to the MAVRIC. The other sources of uncertainty do not change as much.

The temporal evolution of these sources of uncertainty is shown in Fig. 9a by taking the median variance from each of the panels in Fig. 8 for this and other periods. There are three competing factors for how the uncertainty will change with time. First, the SIT is decreasing, and this will reduce the uncertainty as the range of values of which the SIT can occupy shrinks. Second, the separate GCM's simulated SIT responses due to external forcing will differ from each other, causing GCMs to drift apart over time. Thirdly, sea ice at the grid point scale becomes more mobile and vulnerable to external factors as it thins. This will increase variability, initially at least (Sou and Flato, 2009). All of these factors are involved in the evolution of the uncertainties.

The raw CMIP5 subset exhibits a decrease in total uncertainty with time (dashed black in Fig. 9a). This is primarily due to the reduction in model uncertainty (dashed blue), likely because the mean SIT is reducing. The corrected total uncertainty is lower than the raw uncertainty until at least the end of the century. This means that the

MAVRIC can reduce uncertainty and increase confidence in climate projections of SIT throughout this period. The corrected model uncertainty increases for the first three decades, as the models start from a similar state and subsequently diverge because of differing responses to the changes in external forcing. Later the corrected model uncertainty reduces as the mean SIT decreases towards zero.

The total uncertainty is the sum of model uncertainty, internal variability, and scenario uncertainty (see Appendix B for more details). The other panels in Fig. 9 illustrate the relative importance of these sources of uncertainty in terms of the percentage total variance explained, for the raw data, and after the MAVRIC.

Figure 9b illustrates that in the raw projections, model uncertainty remains the dominant (> 50%) source of uncertainty until at least 2100, whereas it only becomes dominant for a few decades mid-century after the MAVRIC (Fig. 9c). The absolute magnitude of internal variability, and its contribution to the total uncertainty, decreases with time because SIT also decreases with time. In the corrected projections, the internal variability is the major contributor to the total uncertainty for the first 25 years, compared to a maximum contribution of only 26% in the raw projections. This highlights the importance of correcting the variance to realistic magnitudes and also the key role of natural variations in predicting the near future evolution of sea ice. The scenario uncertainty accounts for less than 10% of the total uncertainty for the first 50+ years.

4.5 Reduced uncertainty in timing of ice-free conditions

By reducing the model uncertainty, confidence in SIT projections is improved as the range of possible outcomes has been reduced. Figure 10 shows the raw and corrected CMIP5 subset SIV* projections until 2100 using the 18 multi-model ensemble members in each scenario as before. The SIV* calculated here does not consider sea ice concentration (SIC) as it is not bias corrected. Instead, 100% SIC is assumed throughout. It is worth noting that SIV is heavily influenced by the thicker ice to the north of the Canadian archipelago where the true SIC is near 100%, so this assumption should only have a relatively small effect.

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The thick coloured lines are the multi-model scenario mean and the coloured regions represent the 16–84 percentiles (equivalent to 1σ around the mean of a Gaussian distribution) of the ensemble members. To account for the large range in SIT at any particular time in the CMIP5 subset, we use a method similar to that of Massonnet et al. (2012) to calculate first ice-free conditions. We postulate that SIV for ice-free conditions is $2 \times 10^3 \text{ km}^3$, which is in agreement with previous studies calculating first ice-free dates (e.g. Massonnet et al., 2012; Overland and Wang 2013), and is equivalent to two meter thick ice for an ice extent of 10^6 km^2 .

The MAVRIC reduces the total SIV, but the absolute magnitude of this reduction decreases as SIV declines. The 16–84 % range has also been vastly reduced, particularly for the near future. For example, in 2025 the MAVRIC has reduced the 16–84 % range from 12×10^3 to $5 \times 10^3 \text{ km}^3$. It is this reduction in the plausible range of SIV that leads to increased confidence in projections of SIT and SIV. To assess when the Arctic will first display ice-free conditions, we focus on RCP8.5, the most realistic scenario from the last 10 years (Fuss et al., 2014). The cumulative number of ensemble members having satisfied the ice-free criterion as a function of time is shown in Fig. 10c. If uncertainty in this parameter has reduced, this will be shown by the gradient of the line increasing after MAVRIC, and this is clearly seen. Figure 10d further illustrates the uncertainty reduction with boxplots, where the line represents the median (9th) ensemble member to go ice-free. This occurs in 2052 with the MAVRIC, nine years earlier than before. The box represents 16–84 % of the ensemble members, this range has been reduced by about 20 years; dates after 2085 can now be eliminated.

Corrected results from the other emission scenarios show similar features but with later ice-free dates, as expected for lower emissions, and some ensemble members fail to go ice-free by 2100. For RCP4.5 the MAVRIC makes a profound difference with the median ice-free date occurring 35 years earlier in 2060. For RCP2.6 there is uncertainty reduction mid-century but the CMIP5 subset before and after the MAVRIC are in good agreement by the end of the century, with projected ice-free dates around 2090.

5 Summary and discussion

5.1 Summary

This study has developed a bias correction methodology for simulations of sea ice thickness (SIT). By constraining CMIP5 simulations with the PIOMAS reanalysis we have demonstrated that:

- GCMs simulate a wide range of SIT in the historical period and exhibit various spatial and temporal biases when compared with the PIOMAS reanalysis. This model uncertainty is the dominant source of uncertainty in CMIP5 future climate projections of SIT.
- The Mean And VaRIance Correction (MAVRIC) technique outlined in this paper significantly reduces the total uncertainty in future projections of SIT out to 2100 by reducing model uncertainty. Correcting both mean and variance of models is found to be critical for improving the robustness of the projections.
- The MAVRIC results in internal variability being the dominant source of uncertainty until 2022, and model uncertainty is dominant thereafter. From mid-century onwards, scenario uncertainty becomes increasingly important and as influential as model uncertainty by 2100.
- The MAVRIC results in projected September ice-free conditions in the Arctic under RCP8.5 occurring up to 10 years earlier (2050s) than without the correction, and with a much narrower uncertainty range, e.g. excluding post 2085 dates.

5.2 Discussion

Without the MAVRIC, the true magnitude of the internal variability and scenario uncertainty in projections of SIT is concealed by the dominant model uncertainty. This

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



demonstrates that time invested in running many ensemble members to sample internal variability in SIT may be more beneficial than running many future emission scenarios for near term projections. However, for projections at the end of the century, the scenarios become more important.

5 The MAVRIC bias correction technique developed in this study results in a significant improvement in model simulations of SIT with respect to observations. In future projections, the MAVRIC results in a substantial reduction in uncertainty of SIT, leading to increased confidence in climate projections. As absolute values of SIT are utilised, this reduction in uncertainty potentially has important implications for stakeholder sectors operating in Arctic waters such as shipping. The application of the bias correction results in a 60 % reduction in the likely range (16–84 percentiles) of sea ice volume in September 2025.

15 There are a number of caveats to these findings. No attempt is made to constrain the trend in the GCMs. This would be difficult because of the short time scale over which observations are available, raising serious questions about the robustness of calculated historical trends. However future studies could consider this further and assess the feasibility of a trend correction to GCMs. In addition, it is important to recognise that PIOMAS, used here as observations, will also have errors. It would be possible to reduce the multiplicative weightings in Eq. (4) to reflect some uncertainty in the historical data constraint. Other temporally and spatially complete sea ice reanalyses could also be used in future to address this issue.

25 The simulations tend to show an increase in variance as the sea ice thins, before subsequently declining as the thickness approaches zero (Goosse et al., 2009). Blanchard-Wrigglesworth and Bitz (2014) assessed the relationship of this mean state dependant variance in 19 GCMs, including five of the six used in this study, in addition to PIOMAS. They find a relationship between mean thickness variability and mean thickness in models, i.e. models with thicker SIT depict more variable SIT. In the 19 GCMs assessed, PIOMAS sits on the trend line for the correlation between mean thickness variability and mean thickness. However, in the developed MAVRIC, the change in vari-

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ance is decoupled from the applied change to the mean state. This aspect could be further developed, but only by making additional assumptions about future changes in SIT variability.

Studies should make use of this MAVRIC in assessing the impact on potential stakeholders sensitive to SIT and a paper utilising the MAVRIC to investigate the opening of the Arctic sea routes is in preparation. We also intend to make the bias corrected SIT fields freely available online for further investigations.

Appendix A: Supplementary bias correction methodology details

The BC shown in Eq. (4) contains two terms for the representation of the variance in both observations $\sigma_{\hat{\sigma}_h}$ and models $\langle \sigma_{\hat{M}_h} \rangle$. Over the 36 year period of observations the magnitude of the ice loss trend is significant. To accurately calculate variances this externally forced trend should first be removed to leave the variance due to internal variability. Here a choice needs to be made about how best to remove the externally forced trend. For the PIOMAS observations we choose to linearly detrend the monthly data. A smoothed detrending was considered, however this might remove longer time scale variability which is undesirable. Using similar reasoning it is possible that the linear detrending is removing some variability on the multi-decadal timescale. This is assumed to be significantly less than variability on smaller timescales, and much of the trend is attributed to be externally forced over the 36 years, hence should not be included as internal variability. The performance of a smoothed detrend was tested in a theoretical framework and resulted in a 10% loss of accuracy in the standard deviation correction due to describing variance as trend.

The calculation of variance in the models is more complicated due to the fact that there is more than one realisation. It is obvious that the required variance should be calculated from the individual ensemble members rather than the ensemble mean. The variance should be calculated in each ensemble member and then the mean taken. There is another choice to make, i.e. whether each ensemble member should be de-

trended with its own trend, or should the ensemble mean trend be used? We propose that the ensemble mean trend should be used as this is the models response to the changes in forcings. The model detrended ensemble mean standard deviation, $\langle \sigma_{\tilde{M}_h} \rangle$, was calculated by calculating the detrended ensemble variances, then taking the square root of their mean.

The running mean for the future model correction term $\langle \tilde{M} \rangle$ is calculated over an 11 year period of the ensemble mean, this window hence starts at 1975 for the historical calculations. The chosen period must be long enough to adequately smooth the time series, whilst still being able to capture variations in the sea ice decline trend. This was also tested and found to outperform a 21 year period.

Appendix B: Partitioning sources of uncertainty

The sources of uncertainty in Sect. 4.4, Figs. 8 and 9 are calculated for each decadal period (2005–2014, 2015–2024, etc.) separately as follows. Three ensemble members from each of the six GCMs are utilised for three different emission scenarios (RCP2.6, 4.5, and 8.5). This results in each decade having 6 (GCMs) \times 3 (ensemble members) \times 3 (scenarios) \times 10 (years) = 540 (fields).

- The total uncertainty is the variance calculated across all 540 fields.
- The internal variability is calculated similarly to the total variability except instead of the absolute values the anomalies from the models' decadal-mean ensemble-mean for each scenario are used.
- To calculate the model uncertainty, each of the six models' decadal-mean ensemble-mean is calculated, resulting in six fields. The variance is then calculated across these six fields, and repeated for all three scenarios separately (to eliminate differential model dependent responses to the different emission scenarios). The model uncertainty is the square root of the mean of these three fields.

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- The scenario uncertainty is calculated in a similar way. For each model, each of the three scenarios decadal-mean ensemble-means are calculated resulting in three (scenario-dependant) decadal-mean ensemble-means for each of the six models. The variance is then calculated through these three scenario mean fields for each of the six models, resulting in six fields of the variance in each model. The square root of the mean of the six models scenario uncertainty is the scenario uncertainty.

To create Fig. 8b and c it is assumed that the total variance (total uncertainty, T^2) is the sum of the variance due to model uncertainty (M^2), internal variability (I^2), and scenario uncertainty (S^2), formally:

$$T^2 = M^2 + I^2 + S^2 \quad (\text{B1})$$

We note that the variances calculated above do not always sum exactly in this way due to small interaction terms which we ignore.

Author contributions. N. Melia, K. Haines, and E. Hawkins designed the methodology and experiments.

N. Melia developed the code, and performed the experiments.

N. Melia, K. Haines, and E. Hawkins wrote the manuscript.

Acknowledgements. We thank Steffen Tietsche for the conversion of the PIOMAS data, Daniel Feltham and Ellie Highwood for comments on a pre-submission draft and analysis.

N. Melia and E. Hawkins are funded by the APPOSITE project (grant NE/I029447/1), funded by the UK Natural Environment Research Council as part of the Arctic Research Programme. E. Hawkins is also funded by a NERC fellowship. K. Haines is partly funded by the National Centre for Earth Observation NCEO.

References

Blanchard-Wrigglesworth, E. and Bitz, C. M.: Characteristics of Arctic sea-ice thickness variability in GCMs, *J. Climate*, 27, 8244–8258, doi:10.1175/jcli-d-14-00345.1, 2014.

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Christensen, J. H., Boberg, F., Christensen, O. B., and Lucas-Picher, P.: On the need for bias correction of regional climate change projections of temperature and precipitation, *Geophys. Res. Lett.*, 35, L20709 doi:10.1029/2008gl035694, 2008.

Day, J. J., Hargreaves, J. C., Annan, J. D., and Abe-Ouchi, A.: Sources of multi-decadal variability in Arctic sea ice extent, *Environ. Res. Lett.*, 7, 034011, doi:10.1088/1748-9326/7/3/034011, 2012.

Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W., Cox, P., Driouech, F., Emori, S., Eyring, V., Forest, C., Gleckler, P., Guilyardi, E., Jakob, C., Kattsov, V., Reason, C., and Rummukainen, M.: Evaluation of climate models, in: *Climate Change 2013: The Physical Science Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 741–866, 2013.

Francis, J. A. and Vavrus, S. J.: Evidence linking Arctic amplification to extreme weather in mid-latitudes, *Geophys. Res. Lett.*, 39, L06801 doi:10.1029/2012gl051000, 2012.

Fuss, S., Canadell, J. G., Peters, G. P., Tavoni, M., Andrew, R. M., Ciais, P., Jackson, R. B., Jones, C. D., Kraxner, F., Nakicenovic, N., Le Quere, C., Raupach, M. R., Sharifi, A., Smith, P., and Yamagata, Y.: Betting on negative emissions, *Nature Clim. Change*, 4, 850–853, doi:10.1038/nclimate23922014, 2014.

Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., and Vertenstein, M.: The community climate system model version 4, *J. Climate*, 24, 4973–4991, 2011.

Goosse, H., Arzel, O., Bitz, C. M., de Montety, A., and Vancoppenolle, M.: Increased variability of the Arctic summer ice extent in a warmer climate, *Geophys. Res. Lett.*, 36, L23702, doi:10.1029/2009gl0405462009, 2009.

Hawkins, E. and Sutton, R.: The potential to narrow uncertainty in regional climate predictions, *B. Am. Meteorol. Soc.*, 90, 1095–1107, doi:10.1175/2009bams2607.1, 2009.

Hawkins, E. and Sutton, R.: The potential to narrow uncertainty in projections of regional precipitation change, *Clim. Dynam.*, 37, 407–418, doi:10.1007/s00382-010-0810-6, 2011.

Ho, C. K., Stephenson, D. B., Collins, M., Ferro, C. A. T., and Brown, S. J.: Calibration strategies: a source of additional uncertainty in climate change projections, *B. Am. Meteorol. Soc.*, 93, 21–26, doi:10.1175/2011bams3110.1, 2011.

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Jungclaus, J., Keenlyside, N., Botzet, M., Haak, H., Luo, J.-J., Latif, M., Marotzke, J., Mikolajewicz, U., and Roeckner, E.: Ocean circulation and tropical variability in the coupled model ECHAM5/MPI-OM, *J. Climate*, 19, 3952–3972, 2006.

Kay, J. E., Holland, M. M., and Jahn, A.: Inter-annual to multi-decadal Arctic sea ice extent trends in a warming world, *Geophys. Res. Lett.*, 38, L15708, doi:10.1029/2011gl048008, 2011.

Kwok, R., Cunningham, G. F., Wensnahan, M., Rigor, I., Zwally, H. J., and Yi, D.: Thinning and volume loss of the Arctic Ocean sea ice cover: 2003–2008, *J. Geophys. Res.-Oceans*, 114, C07005, doi:10.1029/2009jc005312, 2009.

Laxon, S. W., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., Hendricks, S., Krishfield, R., Kurtz, N., Farrell, S., and Davidson, M.: CryoSat-2 estimates of Arctic sea ice thickness and volume, *Geophys. Res. Lett.*, 40, 732–737, doi:10.1002/grl.50193, 2013.

Lindsay, R. W. and Zhang, J.: Assimilation of ice concentration in an ice–ocean model, *J. Atmos. Ocean. Tech.*, 23, 742–749, 2006.

Lindsay, R. W., Haas, C., Hendricks, S., Hunkeler, P., Kurtz, N., Paden, J., Panzer, B., Sonntag, J., Yungel, J., and Zhang, J.: Seasonal forecasts of Arctic sea ice initialized with observations of ice thickness, *Geophys. Res. Lett.*, 39, L21502, doi:10.1029/2012gl053576, 2012.

Maslanik, J. A., Fowler, C., Stroeve, J., Drobot, S., Zwally, J., Yi, D., and Emery, W.: A younger, thinner Arctic ice cover: increased potential for rapid, extensive sea-ice loss, *Geophys. Res. Lett.*, 34, L24501, doi:10.1029/2007gl032043, 2007.

Massonnet, F., Fichet, T., Goosse, H., Bitz, C. M., Philippon-Berthier, G., Holland, M. M., and Barriat, P.-Y.: Constraining projections of summer Arctic sea ice, *The Cryosphere*, 6, 1383–1394, doi:10.5194/tc-6-1383-2012, 2012.

Meehl, G. A., Washington, W. M., Arblaster, J. M., Hu, A., Teng, H., Kay, J. E., Gettelman, A., Lawrence, D. M., Sanderson, B. M., and Strand, W. G.: Climate change projections in CESM1 (CAM5) compared to CCSM4, *J. Climate*, 26, 6287–6308, 2013.

Overland, J. E. and Wang, M.: When will the summer Arctic be nearly sea ice free?, *Geophys. Res. Lett.*, 40, 2097–2101, doi:10.1002/grl.50316, 2013.

Rotstain, L. D., Jeffrey, S. J., Collier, M. A., Dravitzki, S. M., Hirst, A. C., Syktus, J. I., and Wong, K. K.: Aerosol- and greenhouse gas-induced changes in summer rainfall and circulation in the

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Johns, T. C., Johnson, C. E., Jones, A., Jones, C. P., Joshi, M. M., Keen, A. B., Liddicoat, S., Lock, A. P., Maidens, A. V., Manners, J. C., Milton, S. F., Rae, J. G. L., Ridley, J. K., Sellar, A., Senior, C. A., Totterdell, I. J., Verhoef, A., Vidale, P. L., and Wiltshire, A.: The HadGEM2 family of Met Office Unified Model climate configurations, *Geosci. Model Dev.*, 4, 723–757, doi:10.5194/gmd-4-723-2011, 2011.

Transport Canada: Arctic Ice Regime Shipping System (AIRSS), Transport Canada, Ottawa, 1998.

Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., and Lamarque, J.-F.: The representative concentration pathways: an overview, *Climatic Change*, 109, 5–31, 2011.

Vrac, M. and Friederichs, P.: Multivariate–intervariable, spatial, and temporal–bias correction, *J. Climate*, 28, 218–237, doi:10.1175/jcli-d-14-00059.1, 2014.

Watanabe, M., Suzuki, T., O’ishi, R., Komuro, Y., Watanabe, S., Emori, S., Takemura, T., Chikira, M., Ogura, T., and Sekiguchi, M.: Improved climate simulation by MIROC5: mean states, variability, and climate sensitivity, *J. Climate*, 23, 6312–6335, 2010.

Watanabe, S., Kanae, S., Seto, S., Yeh, P. J. F., Hirabayashi, Y., and Oki, T.: Intercomparison of bias-correction methods for monthly temperature and precipitation simulated by multiple climate models, *J. Geophys. Res.-Atmos.*, 117, D23114, doi:10.1029/2012jd018192, 2012.

Zhang, J. and Rothrock, D.: Modeling global sea ice with a thickness and enthalpy distribution model in generalized curvilinear coordinates, *Mon. Weather Rev.*, 131, 845–861, 2003.

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Table 1. List of models used: the CMIP5 subset and observations.

Institution	Model name	Ensemble members ^a
Commonwealth Scientific and Industrial Research Organisation (CSIRO) Met Office Hadley Centre	CSIRO Mark version 3.6.0: CSIRO-Mk3.6.0 (Rotstayn et al., 2012)	10
	Hadley Centre Global Environment Model version 2-Earth System: HadGEM2-ES (Martin et al., 2011)	4
National Center for Atmospheric Research National Center for Atmospheric Research	Community Climate System Model, version 4: CCSM4 (Gent et al., 2011)	6
	Community Earth System Model, Community Atmosphere Model, version 5: CESM1-CAM5 (Meehl et al., 2013)	3
Model for Interdisciplinary Research on Climate (MIROC) Max Plank Institute for Meteorology (MPI)	MIROC version 5: MIROC5 (Watanabe et al., 2010)	3
	MPI Earth System Model, low resolution: MPI-ESM-LR (Jungclaus et al., 2006)	3
Applied Physics Laboratory (University of Washington)	Pan-Arctic Ice Ocean Modelling and Assimilation System: PIOMAS ^b (Zhang and Rothrock, 2003)	1

^a Multi-model statistics are calculated (Sect. 4.3 onwards) using the first 3 ensemble members.

^b Used as observations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Notation key.

Notation	Description
M	Model
O_h	Observations
x_h	x over the historical period (1979–2014)
\bar{x}	Time mean of x over historical period
$\langle x \rangle$	Ensemble mean of x
\tilde{x}	Running time mean (11 years) of x
\hat{x}	Temporally detrended x over the historical period
σ	Standard deviation

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.




Figure 1. September 1979–2014 mean SIT and standard deviation (SD) from the PIOMAS reanalysis. SD is calculated after removing the linear trend.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.




Figure 2. Mean September SIT for each of the six GCMs considered, averaged over the period 1979–2014.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.




Figure 3. Performance of different SIT BCs for one particular month at a hypothetical grid point in a toy model. Mean, SD (detrended) and trend legend statistics are calculated over the observation period (1979–2014). “Ice-free” is defined as the first occurrence of any ensemble member below 0.15 m. The black line represents “observations”, the blue and red lines represent high and low ice models respectively. The thin coloured lines represent ensemble members, and the thick lines are the ensemble mean.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.




Figure 4. September SIT at three grid point locations in the Arctic, from PIOMAS (black) and CSIRO-Mk3.6.0 historical (1979–2005) and RCP8.5 (2006–2014) raw output (grey) and post MAVRIC (green). The raw CSIRO ensembles (grey) are bias corrected via the MAVRIC using the PIOMAS observations (black) over the calibration window, producing the MAVRIC ensembles (green) for the validation window. Beanplots (right) show the distribution of the SIT for the validation period. Small horizontal lines show every SIT value, the frequency of which is illustrated by the width of the shaded region. Thick horizontal line is the mean.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.




Figure 5. September SIT at three grid point locations in the Arctic, from PIOMAS (black) and CSIRO-Mk3.6.0 historical (1979–2005) and RCP8.5 (2006–2100) raw output (grey) and post MAVRIC (green). Thin lines are individual ensemble members, thick lines are the ensemble means. Mean, SD and trend legend statistics calculated over the period of observations (1979–2014). The SD is the detrended mean ensemble SD. Ice-free is the range of the first occurrence of the first and last ensemble member below 0.15 m.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.




Figure 6. CSIRO-Mk3.6.0 and HadGEM2-ES, September 1979–2014 ensemble mean SIT and SD (detrended). The raw columns are the model solutions as found in the CMIP5 archive. The corrected columns show the distribution after the MAVRIC has been applied.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.




Figure 7. September mean SIT from the CMIP5 subset, using the raw data (top row) and after MAVRIC (bottom row). The multi-model ensemble mean (three members from each model) is shown.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.




Figure 8. September 2015–2024 sources of SIT uncertainty from the CMIP5 subset. The multi-model ensemble mean (three members from each) is shown when comparing raw (top row) and after MAVRIC (bottom row).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 9. The evolution of the sources of September SIT uncertainty in the CMIP5 sub-set with lead time. Year zero is the MAVRIC window mid-point (1997) and the emission scenarios (RCPs) start in 2006. Panel (a) shows the change in magnitude of the different sources of uncertainty. The uncertainty shown is the median SIT variance and hence the lines scale additively. The dashed lines are for the raw model output and solid lines are for post MAVRIC. Contributions of model uncertainty, internal variability and scenario uncertainty as a fraction of total uncertainty are shown for the raw output (b) and post MAVRIC (c).

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved Arctic sea ice thickness projections using bias corrected CMIP5 simulations

N. Melia et al.




Figure 10. CMIP5 subset sea ice volume (SIV*) projections and first ice-free conditions. Panels (a, b) show the projected SIV* from all six models (18 ensemble members total) in both the raw and corrected GCMs (11 year running mean), and shaded regions are the 16–84th percentiles. Panel (c) shows the number of ensemble members having passed the ice-free threshold. Panel (d) shows the statistics of (c), with the whiskers representing the range (1st and 18th ensemble member ice-free), the box capturing the 16–84th percentiles, and the bold line showing the median (9th ensemble member). Ice-free is defined as the first year the pan-Arctic SIV* dips below $2 \times 10^3 \text{ km}^3$ for a particular ensemble member. *Volume (SIV*) is calculated using 100% sea ice concentration (SIC) throughout.