

Linking
catchment-scale
subglacial discharge
to glacially modified
waters

L. A. Stevens et al.

Linking catchment-scale subglacial discharge to subsurface glacially modified waters near the front of a marine terminating outlet glacier using an autonomous underwater vehicle

L. A. Stevens¹, F. Straneo², S. B. Das³, A. J. Plueddemann², A. L. Kukulya⁴, and M. Morlighem⁵

¹Massachusetts Institute of Technology, Woods Hole Oceanographic Institution Joint Program in Oceanography, Applied Ocean Science and Engineering, Woods Hole, MA 02543, USA

²Department of Physical Oceanography, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

³Department of Geology and Geophysics, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

⁴Department of Applied Ocean Physics and Engineering, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

⁵Department of Earth System Science, University of California, Irvine, Croul Hall, Irvine, CA 92697, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 29 July 2015 – Accepted: 15 August 2015 – Published: 1 September 2015

Correspondence to: L. A. Stevens (stevensl@mit.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

TCD

9, 4583–4624, 2015

**Linking
catchment-scale
subglacial discharge
to glacially modified
waters**

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Measurements of near-ice (<200 m) hydrography and near-terminus subglacial hydrology are lacking due in large part to the difficulty in working at the margin of calving glaciers. Here we pair detailed hydrographic and bathymetric measurements collected with an Autonomous Underwater Vehicle as close as 150 m from the ice/ocean interface of the Sarqardliup sermia/Sarqardleq Fjord system, West Greenland, with modeled and observed subglacial discharge locations and magnitudes. We find evidence of two main types of subsurface glacially modified water localized in space and with distinct properties that are consistent with runoff discharged at two locations along the grounded margin. These locations, in turn, correspond with two prominent subglacial subcatchments beneath Sarqardliup sermia. Thus, near-ice observations and subglacial discharge routing indicate that subglacial discharge from this glacier occurs at only two primary locations and gives rise to two distinct glacially modified waters. Furthermore, we show that the location with the largest discharge flux is associated with the lighter, fresher glacially modified watermass. This is qualitatively consistent with results from an idealized plume model.

1 Introduction

Greenland Ice Sheet mass loss quadrupled over the last two decades, contributing roughly 7.4 mm to global sea level rise from 1992–2011 (Shepherd et al., 2012), and increasing freshwater inputs into the North Atlantic (Bamber et al., 2012). Ice sheet mass loss occurs through run-off of surface melt, ice discharge through iceberg calving, and submarine melt at marine-terminating outlet glacier margins (van den Broeke et al., 2009; Enderlin et al., 2014). The synchronous retreat and speedup of marine-terminating glaciers in southeast Greenland in the early 2000s was likely initiated by a dynamic change at marine termini (van den Broeke et al., 2009; Rignot and Kanagaratnam, 2006; Thomas et al., 2009), and points towards common external forcings from

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the warming atmosphere (Box et al., 2009) and/or ocean around Greenland (Straneo and Heimbach, 2013), though the exact forcing mechanisms and relative magnitudes remain unclear (Joughin et al., 2012; Straneo et al., 2013).

Increased submarine melt rates at outlet glacier marine termini may be a leading cause of Greenland Ice Sheet outlet glacier speed up and retreat (Holland et al., 2008; Joughin et al., 2012; Motyka et al., 2013; Post et al., 2011). The heat to drive submarine melting is supplied by waters of the subpolar North Atlantic and Arctic Seas, whose circulation inside the fjords is a result of processes across a range of spatiotemporal scales (Jackson et al., 2014; Straneo et al., 2010). Ultimately, melt rates are affected by ocean properties (temperature and stratification) and circulation in near-ice waters (< 200 m) (Jenkins et al., 2010). Submarine melting is thought to be enhanced in summer as a result of meltwater runoff along the ice sheet bed entering the fjord across the grounding line as subglacial discharge, which provides an additional buoyancy source alongside submarine melt for initiating buoyant plumes along the terminus face (Jenkins, 1999, 2011; Sciascia et al., 2013; Xu et al., 2013). Relatively fresh waters rising in the core of these plumes become denser as they entrain salty ambient fjord waters, and serve as a mechanism for transporting ambient fjord waters to the glacier face (Jenkins, 1999, 2011; Sciascia et al., 2013; Xu et al., 2013).

Plume theory and models combined with melt rate parameterizations suggest that higher subglacial discharge rates drive higher entrainment and submarine melt rates (Jenkins, 1999, 2011; Sciascia et al., 2013; Xu et al., 2013), however ocean property and plume measurements needed to inform and validate model simulations and theory are lacking due to difficulty in working at the margin of calving glaciers (Straneo and Cenedese, 2015). As a result, current modeling-sourced estimates of submarine melt rates at tidewater glaciers and their sensitivity to external forcings of the near-ice environment are highly uncertain, and based on unconstrained models of plume dynamics using ice/ocean boundary parameterizations forced by far field (> 1 km) ocean property measurements and largely unknown subglacial discharge fluxes (Jenkins, 2011; Kimura et al., 2014; Sciascia et al., 2013; Slater et al., 2015; Xu et al., 2012, 2013).

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

For example, in a recent numerical study the spatial distribution of subglacial discharge along the grounding line was found to have a large effect on both the total submarine melt rate and its distribution along marine termini (Slater et al., 2015). With a lack of observations of both the near-ice environment and subglacial discharge configurations, we are unable to define likely subglacial discharge scenarios and their associated influence on ice/ocean interactions, resulting in an inadequate and untested understanding of how tidewater glaciers respond to oceanic forcing now and in the future (Straneo and Cenedese, 2015). Specifically, ocean measurements collected at distances > 1 km from the glacier terminus provide limited information on the near-ice processes because the signals of glacial modification have, by that time, largely been smeared by lateral mixing processes. Indeed, the picture that emerges from such far-field measurements is of a horizontally invariant overturning cell(s) (Chauché et al., 2014; Inall et al., 2014; Johnson et al., 2011; Mortensen et al., 2011; Straneo et al., 2011; Sutherland et al., 2014).

In this study, we present fjord hydrography and bathymetry measurements from the near-ice environment of a tidewater glacier in west Greenland (Fig. 1) that allow us to reconstruct the distribution of subglacial discharge and provide key details on the ice-ocean exchanges. We do this by identifying the distribution of Glacially Modified Waters (GMW) – a product of ambient fjord waters mixing with subglacial discharge and glacial melt (Jenkins, 2011; Straneo et al., 2011) – within a few 100 m of the glacier face, and by delineating the subglacial catchments that route subglacial meltwater to discharge locations along the grounded terminus. These hydrographic measurements were obtained primarily in July 2012, using a REMUS-100 (Remote Environmental Measuring UnitS) Autonomous Underwater Vehicle (AUV) (Fig. 2a) to observe the temperature, salinity, and turbidity of waters in Sarqardleq Fjord (SF) from ~ 2 km away to within a couple hundred meters of Sarqardliup sermia (SS), a medium-sized tidewater glacier in West Greenland (68.90° N, 50.32° W) (Fig. 1). This novel, high-risk field campaign was successful in obtaining multiple vertical sections of fjord water properties as

the SS outlet glacier during the 2007–2009 winters were on order 125–175 m yr⁻¹, with the center third of the SS terminus reaching speeds of 200 m yr⁻¹ (Joughin et al., 2013).

The Sarqardleq-Tasiussaq fjord system is the southern side fjord off the larger, deeper Jakobshavn Isbræ (JI) fjord, which connects the largest and fastest Greenland ice stream (JI) to Disko Bugt (Fig. 1a). From the SS terminus, the shallower Sarqardleq-Tasiussaq Fjord system extends roughly 30 km to the northwest before reaching JI fjord. SF meets Tasiussaq Fjord over a previously unknown 70 m-deep sill, 15 km from the SS terminus (Figs. 1 and 7a). Tasiussaq Fjord meets JI fjord over an at most 125 m-deep sill (Gladish et al., 2015a) 30 km from the SS terminus (Fig. 1). Waters along the SS terminus range from 20–150 m-depth, and are deepest in two troughs near the center of the glacier (Fig. 2, Table 3). Both SS lateral terminus regions are grounded in relatively shallow lagoons (< 20 m) (Fig. 3). A 40 m-deep seamount is located 2.5 km from the vertical SS calving face (Fig. 3).

3.2 Subglacial catchment and runoff

To first order, subglacial catchments are delineated by ice sheet surface and bed topography, which governs subglacial hydraulic potential at the bed (Cuffey and Patterson, 2010). It is well known that gradients in subglacial hydraulic potential at the ice-sheet bed do not completely dictate subglacial meltwater pathways due to the constantly evolving subglacial hydraulic system over the summer melt season (Andrews et al., 2014; Chandler et al., 2013; Hewitt et al., 2012; Schoof, 2010). However, recent modeling studies of supra- and subglacial meltwater routing find that topographic channeling of surface routing and channelized subglacial drainage systems occurs due to surface and bed gradients in both alpine valley glacier (Werder et al., 2013) and Greenland Ice Sheet outlet glacier (Banwell et al., 2013; Palmer et al., 2011) systems. Here we use a simple approach for calculating subglacial meltwater routing involving subglacial hydraulic potential gradients from two recent Greenland bedmaps (Bamber et al., 2013; Morlighem et al., 2014) and runoff values from the Regional Atmospheric Cli-

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lower than 400 m a.s.l. were prescribed the same runoff values as the RACMO2.3 grid point within the catchment at 432 m a.s.l. (68.82° N, 50.19° W) (Fig. 7a), as there are no RACMO2.3 grid points at lower elevations within the catchment. We assume that the ice-sheet bed is impermeable (does not store water) over the timescales considered here, and that all surface runoff is transferred immediately to the bed directly beneath the location of runoff formation at the ice sheet surface.

4 Results

4.1 Glacially Modified Water (GMW) temperature, salinity, and turbidity properties in Sarqardleq Fjord

We find the summer fjord waters are characterized by a ~ 10–20 m fresh and relatively warm surface layer overlying a thick layer of weakly stratified, relatively salty ($S = 30.5$ – 32.5) and cold ($\theta \approx 1^\circ\text{C}$) waters (Table 2, Fig. 5a and b). The summer fjord waters are the same as the Surface Waters (SW) and Ilulissat Icefjord Waters (IIW) observed by recent hydrographic surveys throughout Ilulissat Icefjord (Gladish et al., 2015a, b). SW are a mixture of IIW and fresher, warmer waters originating from local freshwater sources and warmed by summer atmospheric forcing. IIW originates from Arctic Waters observed in Disko and Baffin Bays (Gladish et al., 2015b) that enter SF after crossing sills at the mouth of JI fjord (Schumann et al., 2012), the confluence of JI fjord and Tasiussaq fjord (Gladish et al., 2015a), and the mouth of SF (Fig. 1). These summer fjord waters are observed in the outer SF by a set of far-field CTD profiles taken near the fjord mouth more than 10 km from the SS terminus (triangles in Fig. 7a). We define ambient fjord waters as the average of these far-field CTD profiles (red profile in Figs. 5 and 6).

Near the glacier we observe a range of water masses not found in the outer fjord. These waters are generally colder, fresher, and more turbid than waters near the mouth of the fjord (Fig. 5a and b). The REMUS sections reveal two distinct GMW, which we

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

refer to as GMW1 and GMW2 (Fig. 4, Table 2). GMW1 and GMW2 are cold anomalies with a high turbidity signal that are most evident at two distinct locations (Fig. 4). GMW1 is observed in the southwestern ends of R1–R5 at ~ 40 m depth, while GMW2 is observed in the northeastern ends of R1–R5 at ~ 60 m depth (Fig. 4). Both GMW1's and GMW2's temperature and turbidity anomalies are most pronounced close to the glacier (Fig. 4a–c), and decrease as these waters spread away from the glacier (Fig. 4g–i). For example, the high turbidity associated with GMW1 spreads laterally beneath the pycnocline at R1 (Fig. 4i). Turbidity does not consistently map onto regions of local temperature minima; there are regions in the REMUS sections with high turbidity but with temperatures above 0.9°C (northeastern R1 below 80 m depth; Fig. 4i). High turbidity in these regions may be due to other sources including suspended sediment sourced from proglacial streams that enter SF as surface runoff near the northeastern end of R1 (Fig. 3) or iceberg discharge.

CTD casts 1–3 were taken closer to the SS face than the R5 transect during the same July 2012 field campaign (Fig. 3), and provide additional θ/S characteristics below the 100 m REMUS depth limit (Fig. 6a–c). These casts record deeper cold anomalies at the bottom of SF, as well as cold excursions from ~ 40 to 80 m depth, similar to REMUS measurements (Fig. 6a–c). Overall the CTD profiles align well with REMUS measurements where coincident (above 100 m).

Further insight into the origins of GMW1 and GMW2 is found in θ/S space, where GMW1 and GMW2 stand out as cold anomalies as compared to waters near the mouth of the fjord (Figs. 5d, 6a and b). GMW1 and GMW2 are clustered at two distinct densities (Fig. 6a and b). At a density of $\sigma_\theta \approx 24.8 \text{ kg m}^{-3}$, where σ_θ is potential density less 1000 kg m^{-3} , GMW1 is lighter than GMW2 ($\sigma_\theta \approx 25.5 \text{ kg m}^{-3}$) (Table 2, Fig. 6a and b). In general, GMW is fresher and more turbid compared to ambient waters, consistent with fjord waters mixing with submarine melt and subglacial discharge. If we assume that both GMW1 and GMW2 are driven by subglacial discharge plumes that emerged at the grounding line, then we can assume that the bulk of the entrainment was of deeper waters at densities of $\sigma_\theta = 26\text{--}26.5 \text{ kg m}^{-3}$ (Fig. 6a and b). In θ/S space,

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Fig. 7b). Disregarding the possibility for periods of subglacial water storage during the en- and subglacial transport of runoff to the SS terminus, daily discharge rates across the terminus during the field expedition are $146 \text{ m}^3 \text{ s}^{-1}$ (MBM2014 estimate) (Table 3). An additional though likely minor amount of surface meltwater runoff enters the fjord through proglacial streams, which discharge at land-terminating margins abutting SS (Fig. 2). Daily runoff discharges for C1 and C2 scale primarily with area differences and are 115.78 and $20.62 \text{ m}^3 \text{ s}^{-1}$, respectively (MBM2014) (Table 3). While error estimates for the RACMO2.3 runoff rates are not calculated, we take the standard deviation of July 2012 daily discharge rates as a measure of the potential variation observed during the field expedition (Table 3).

4.3 Buoyant plume model for the SS/SF system

As described above, we have found evidence for three main subglacial catchments discharging runoff into SF at three locations along the terminus. The two prominent discharge locations, D1 and D2, coincide with GMW1 and GMW2 observations. The picture that emerges is that different properties of GMW1 and GMW2 are attributable to differences in subglacial discharge magnitude at that location. Here, we use a buoyant plume model to investigate the extent to which the two plumes' predicted characteristics compare with the GMW1 and GMW2 observations. Buoyant plume theory states that the growth of a plume is dictated by the plume's buoyancy forcing, which can be due to subglacial discharge at the grounding line and/or submarine melting along the terminus (Morton et al., 1956; Turner, 1979). The buoyancy forcing of the plume determines the plume's vertical velocity and entrainment of ambient fjord waters (Morton et al., 1956; Turner, 1979). A class of simple, one-dimensional buoyant plume models has been used to investigate plume dynamics and terminus melt rates near glaciers (Hellmer and Olbers, 1989; Jenkins, 1991, 2011). Solutions to these models estimate plume temperature, salinity, vertical velocity, and intrusion depth, the depth at which the plume becomes neutrally buoyant and changes from flowing vertically up the terminus

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



have yet to be well observed at submerged glacier termini, we take the channel mouth geometry to be semi-circular, following observations of channel mouth geometries at land terminating glaciers (Fountain and Walder, 1998). The Jenkins (2011) model is uniform in the across-flow direction, thus the subglacial flux, q_{sg} , used for model simulations is the flux through a 1 m vertical slice at the center of the channel, such that $q_{sg} = (Q_{sg} r) S^{-1}$, where r is the radius of the channel (Table 4).

Given the observed ocean stratification and the modeled subglacial discharge, the plume model confirms that GMW1 should be notably fresher and lighter than GMW2 (Fig. 5c, Table 4). This supports the conclusion that GMW1 and GMW2 are the result of two distinct discharge locations with different subglacial discharge magnitudes. In both scenarios, the plume model predicts plume properties that are much lighter and fresher than the observed GMW (Fig. 5c, Tables 2 and 4). The predicted D1 and D2 plumes would reach the pycnocline at neutral buoyancy depths of ~ 7 and ~ 10 m, respectively (Table 4). With a minimum amount of overshoot, we might expect the plumes to reach the surface or depths close enough to the surface to be visible during field observations. In reality, the plumes were not observed to reach the surface, and GMW was only observed beneath the pycnocline (Fig. 4). There are several possible reasons for this discrepancy. First, the plume model may have an incorrect entrainment and/or submarine melt parameterization. Second, the prescribed subglacial flux, a function of estimated subglacial discharge and channel surface area, could be incorrect. In addition, after detaching from the terminus at the plume's intrusion depth, GMW spreads an additional 150 m away from the SS face before being observed at R5. Over this time, we would expect lateral mixing to further dilute the GMW properties. The plume model does not describe lateral mixing, as the model ends when the plume reaches intrusion depth.

5 Discussion

5.1 Subglacial catchments, discharge, and GMW observations

Our analysis of the ocean data and subglacial catchments both suggest that there are two primary subglacial discharge locations along the ice/ocean interface. On the outlet glacier catchment side of the interface, the primary subcatchments, C1 and C2 (Fig. 7a), route substantial (> 90%) of the total SS meltwater runoff (Table 3) into the fjord across the grounding line at discharge locations D1 and D2, respectively (Fig. 3). On the ocean side of the interface, GMW1 and GMW2 are located near D1 and D2, respectively, and show fresher, colder waters with high turbidity as compared to ambient fjord waters (Fig. 5a and b). The properties of these waters, in particular, are consistent with glacial modification due to significant injection of runoff at depth as is expected from a localized discharge of meltwater at D1 and D2. Finally, between D1 and D2, there is a 2 km stretch of the terminus where GMW show cold excursions with low to high turbidity along R4 and R5 (Fig. 6c). The formation of this GMW is less clear, though in this region between subglacial discharge locations, GMW properties are more indicative of submarine melt and limited subglacial discharge and/or lateral mixing of GMW1 and GMW2.

Although we lack observations within the plumes themselves in 2012, the ocean observations of GMW suggest that these waters are produced by ambient fjord waters interacting with a limited number of discrete plumes along the terminus. Our observations of GMW beneath the pycnocline at a distance of ~150 m from the terminus suggest that the two plumes reach neutral buoyancy beneath the fjord surface. Visual observations during the 2012 field campaign confirm that the plumes did not reach the fjord surface during this time. In contrast, during the July 2013 field campaign at SF, a vigorous, turbulent plume was observed to break through at the fjord surface at D1 (Mankoff et al., 2014).

Differences in subglacial discharge magnitude entering the fjord at D1 and D2 is both observed and predicted to result in water mass differences between GMW1 and

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

GMW2. Fed by subglacial discharge from the largest subglacial subcatchment, GMW1 is fresher and lighter than GMW2 (Table 3, Figs. 5a–d and 6a, b). D2 receives roughly 20% of the subglacial discharge magnitude at D1 (Table 3). This smaller subglacial discharge results in a relatively saltier and heavier GMW2 in comparison to GMW1 (Figs. 5a–d and 6a, b). While a greater volume of subglacial discharge leads to a fresher water mass, the strength of the resultant buoyant plume also plays a role in near-ice water mass transformation. Plume theory predicts that a plume fed by a greater amount of subglacial discharge will have a stronger buoyancy forcing, leading to faster entrainment of ambient waters and a decrease in the fraction of subglacial discharge in the plume (Jenkins, 2011). In this fjord, the entrainment of ambient waters into a plume results in GMW with temperatures and salinities that are warmer and saltier than the subglacial discharge entering the fjord ($\theta = 0^\circ\text{C}$, $S = 0$ PSU). The differences between GMW1 and GMW2 temperature are consistent with plume theory, as GMW1 temperatures are closer to ambient fjord waters than the colder GMW2 (Fig. 6a and b). Greater subglacial discharge at D1 drives a more vigorous plume and greater mixing with ambient fjord waters, which results in GMW that is closer in θ and S to IIW (Fig. 6a). Smaller subglacial discharge at D2 drives a less vigorous plume with less mixing with ambient fjord waters, resulting in GMW that retains the cold signature of subglacial discharge and submarine melting.

Consistent with the ocean data, the plume model predicts end plume conditions at D1 are fresher and lighter than those at D2 as they contain a greater amount of subglacial discharge (Fig. 5d, Table 4). However, the end plume conditions from the Jenkins (2011) model for both D1 and D2 scenarios are lighter than the GMW we observe and even portions of the surface waters of SF (Fig. 5c, Tables 2 and 4). In addition to errors in the plume model and subglacial flux estimates, lateral mixing within ~ 150 m of the terminus is an additional consideration for comparing the plume model results and observed GMW. Large amounts of mixing with ambient waters likely occur once the plume detaches from the terminus and GMW is exported away from the ice/ocean interface. This lateral mixing has been observed in other marine terminating

outlet glacier systems in Greenland, where GMW from an inferred localized subglacial discharge location was found uniformly across the fjord in profiles taken ~ 200 m from the terminus (Chauché et al., 2014).

5.2 Observing the heterogeneous near-ice environment

5 The coupling of near-ice observations and subglacial discharge routing is necessary for understanding ice-ocean interactions at marine terminating outlet glaciers. While multiple recent studies have observed GMW in fjords (Chauché et al., 2014; Inall et al., 2014; Johnson et al., 2011; Mortensen et al., 2011; Straneo et al., 2011; Sutherland et al., 2014) and others have measured and modeled runoff based on surface catchment
10 area (Mernild et al., 2015), no studies have directly linked the two sides of this interface or considered the role of basal routing on catchment area. For this study, we pair near-ice observations and subglacial discharge routing to show for the first time that the observed GMW characteristics align with the subglacial discharge magnitudes from outlet glacier subcatchments.

15 Our results highlight the necessity of subsurface observations within the near-ice zone for accurately characterizing the heterogeneous processes at the ice/ocean interface. We observe heterogeneous, subsurface GMW as high turbidity, cold excursions in across-fjord sections as far as 1.5 km from the SS terminus (Fig. 4). Further away from the terminus, only the cold excursion at the density of GMW1 remains in the far-field profiles (Fig. 5d). Thus, while in the near-ice zone there are multiple subglacial discharge locations across the SS grounding line and different types of GMW observed, only a modified GMW1 is identifiable in far-field profiles. Noble gas observations of GMW in neighboring Greenland fjords observe a dilution of GMW as you move away from the terminus, suggesting that GMW is highly diluted outside of the near-ice zone
20 (Beaird et al., 2015). Thus, the fact that only a modified GMW1 is detectable in the far-field profiles is likely due to the larger volume flux of GMW1 entering the fjord as compared to GMW2 (Table 4). Sill depth may be an additional factor impeding the export of GMW2; GMW2 is observed at or barely above the 70 m sill depth, while GMW1

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is observed at shallower depths (Figs. 1 and 3, Table 2). The implication is that far-field measurements only provide a partial representation of processes along the ice/ocean interface.

Similar to the single cold excursion observed in the ambient SF waters, many studies have observed evidence of subsurface GMW uniformly distributed across fjord width outside of the near-ice zone (Johnson et al., 2011; Mortensen et al., 2011; Straneo et al., 2011; Chauché et al., 2014; Inall et al., 2014; Sutherland et al., 2014). Observations at Store and Rink glaciers as close as ~200 m to termini identify one to a couple surface and subsurface plumes along each glacier termini (Chauché et al., 2014). However, the GMW observed 200 m from the termini is uniform across the fjord (Chauché et al., 2014). While our observations of subglacial discharge locations in SF are consistent with the low number of subglacial discharge locations found at Store and Rink glaciers (Chauché et al., 2014), we are able to further differentiate and map types of GMW to outlet glacier subcatchments.

The subsurface nature of the plumes and resultant GMW we observed is consistent with multiple studies that have also observed subsurface GMW (Chauché et al., 2014; Inall et al., 2014; Johnson et al., 2011; Mortensen et al., 2011; Straneo et al., 2011; Sutherland et al., 2014). Together these findings drive home the point that plumes and other processes at the ice/ocean interface actively driving submarine melt can and often do operate without creating an expression on the fjord surface. Surface expressions of plumes have been detected at many Greenland tidewater glaciers and invoked as evidence for runoff release from the ice sheet into fjords and proglacial streams (Chu et al., 2009; Tedstone and Arnold, 2012), and have even been proposed as a potentially useful remote measure of runoff variability (Chu et al., 2012). However, our observations of plumes and GMW that reach neutral buoyancy beneath the pycnocline suggest in many cases this relationship does not hold true. The magnitude of subglacial discharge entering a fjord, fjord stratification, and fjord depth have all been shown to affect whether a plume reaches the surface (Sciascia et al., 2013). The absence of plume surface expression does not negate the presence of subglacial discharge plumes that

Linking
catchment-scale
subglacial discharge
to glacially modified
waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



may be driving significant submarine melt and circulation along a tidewater terminus. Thus, across-fjord subsurface observations within the near-ice zone provide the most comprehensive characterization of ice/ocean interactions in Greenland fjords.

5.3 Observational constraints for modeling the heterogeneous near-ice environment

While spatial distribution of subglacial discharge is a critical component for estimating submarine melt rates at marine terminating outlet glaciers in numerical models (Slater et al., 2015), we have few observations to constrain subglacial discharge scenarios. Model configurations of subglacial discharge for major Greenland outlet glaciers range from a distributed subglacial system where equal amount of subglacial discharge emerge across the entire grounding line width (Jenkins, 2011; Sciascia et al., 2013), to distributing subglacial discharge between a number of equally-spaced plumes along the terminus (Kimura et al., 2014; Slater et al., 2015), to routing all subglacial discharge through one, central plume (Slater et al., 2015; Xu et al., 2013). All these models, which share the same melt parameterization, agree that submarine melt rates increase with increasing subglacial discharge (Jenkins, 2011; Kimura et al., 2014; Sciascia et al., 2013; Slater et al., 2015; Xu et al., 2012, 2013), the amount and distribution of the increased melting depends on the largely unknown pattern of subglacial discharge (Straneo and Cenedese, 2015). Most recently, Slater et al. (2015) concluded that a distributed system yields as much as 5 times more submarine melting than a channelized system consisting of a few plumes along the terminus (Slater et al., 2015). Thus, spatial distribution of subglacial melt is critically important for estimating submarine melt rates in a numerical model (Slater et al., 2015; Straneo and Cenedese, 2015).

For this system, we observe two, localized areas of subglacial discharge separated by wide areas of the terminus with little to no subglacial discharge. A simple subglacial meltwater routing model using MBM2014, the GIMP ice sheet surface digital elevation model, and RACMO2.3 runoff estimates was able to predict the number, approximate location, and relative magnitude of subglacial discharge locations. And while this sub-

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



glacial catchment delineation method should be supplemented with ocean measurements and field observations where possible, in many cases it may prove a useful first order approximation of the spatial distribution of subglacial discharge at marine terminating outlet glaciers where fjord observations are lacking or difficult to obtain.

6 Conclusions

Hydrographic surveys completed by an AUV in Sarqardleq Fjord provide several new observational insights to the characteristics and distribution of near-ice GMW in a shallow-silled, moderate-sized west Greenland fjord. Overcoming navigation difficulties in the acoustically noisy, iceberg-filled fjord, the AUV covered a large portion of the near-ice waters along the terminus. AUV observations provide the most comprehensive and spatiotemporally detailed snapshots of across-fjord hydrography in the near-ice zone to date. From these measurements we identified two types of GMW that map onto two plumes based on θ/S /turbidity near-ice properties and subcatchment runoff estimates. The two plumes are, notably, not observed to reach the surface in the fjords, but attain neutral buoyancy beneath the pycnocline of the strongly stratified summer fjord conditions.

Our observations detail how mixing processes at the ice/ocean interface driven by either submarine melting and/or plumes fed by subglacial discharge can produce GMW that is colder, fresher, and at times more turbid than ambient fjord waters. An idealized plume model for plumes fed by a range of RACMO2.3-derived subglacial fluxes appropriate for the two plumes observed in this fjord is qualitatively consistent with the largest discharge flux being associated with the lighter, fresher glacially modified watermass. The characterization of GMW and subglacial catchments for this outlet glacier system provides critical observational constraints on the widely varying subglacial discharge scenarios employed by the current set of submarine melt modeling studies. Results supply near-ice observations abutting one Greenland Ice Sheet outlet glacier, though the continued investigation of other Greenland outlet glaciers is much needed to ul-

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



timately move towards an accurate representation of oceanic forcing at outlet glacier termini and an improved understanding of the ice sheet's outlet glacier dynamics.

Author contributions. F. Straneo, S. B. Das, and A. J. Plueddemann conceived the study. F. Straneo, S. B. Das, and A. L. Kukulya performed the fieldwork. A. J. Plueddemann, A. L. Kukulya, and L. A. Stevens processed the REMUS data. F. Straneo, S. B. Das, A. J. Plueddemann, and L. A. Stevens analyzed the REMUS and CTD data. L. A. Stevens created the bathymetry map. M. Morlighem provided the reprocessed bedmap. L. A. Stevens, F. Straneo, S. B. Das, and A. J. Plueddemann interpreted the results. L. A. Stevens wrote the paper. All authors commented on the paper.

Acknowledgements. Support was provided by the National Science Foundation's Office of Polar Programs (NSF-OPP) through PLR-1418256 to F. Straneo, S. B. Das and A. J. Plueddemann, PLR-1023364 to S. B. Das, and through the Woods Hole Oceanographic Institution Ocean and Climate Change Institute Arctic Research Initiative to F. Straneo, S. B. Das, and A. J. Plueddemann. L. A. Stevens was also supported by a National Science Foundation Graduate Research Fellowship. S. B. Das was also supported by the Woods Hole Oceanographic Institution James E. and Barbara V. Moltz Research Fellowship. M. Morlighem was supported by the National Aeronautics and Space Administration's (NASA) Cryospheric Sciences Program through NNX15AD55G. The authors would like to acknowledge Michiel van den Broeke for providing RACMO output; Jeff Pietro and John Kemp of the WHOI Mooring Operations, Engineering and Field Support Group and Rebecca H. Jackson and Ove Villadsen for their help during the 2012 field operations; Clark Richards for guidance using the Barnes Objective Analysis; Kenneth D. Mankoff for conversations on plumes in Sarqardleq Fjord; Nicholas L. Beard for conversations on glacially modified waters in Greenland fjords; and Adrian Jenkins for the plume model code.

The authors declare no competing financial interests.

TCD

9, 4583–4624, 2015

Linking
catchment-scale
subglacial discharge
to glacially modified
waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Andrews, L. C., Catania, G. A., Hoffman, M. J., Gullely, J. D., Lüthi, M. P., Ryser, C., Hawley, R. L., and Neumann, T. A.: Direct observations of evolving subglacial drainage beneath the Greenland Ice Sheet, *Nature*, 514, 80–83, doi:10.1038/nature13796, 2014.
- 5 Bamber, J., van den Broeke, M., Ettema, J., Lenaerts, J., and Rignot, E.: Recent large increases in freshwater fluxes from Greenland into the North Atlantic, *Geophys. Res. Lett.*, 39, 1–4, doi:10.1029/2012GL052552, 2012.
- Bamber, J. L., Griggs, J. A., Hurkmans, R. T. W. L., Dowdeswell, J. A., Gogineni, S. P., Howat, I., Mouginot, J., Paden, J., Palmer, S., Rignot, E. and Steinhage, D.: A new bed elevation dataset for Greenland, *The Cryosphere*, 7, 499–510, doi:10.5194/tc-7-499-2013, 2013.
- 10 Banwell, A. F., Willis, I. C., and Arnold, N. S.: Modeling subglacial water routing at Paakitsoq, W Greenland, *J. Geophys. Res.-Earth*, 118, 1282–1295, doi:10.1002/jgrf.20093, 2013.
- Barnes, S. L.: Applications of the Barnes Objective Analysis Scheme, Part I: Effects of Under-sampling, Wave Position, and Station Randomness, *J. Atmos. Ocean. Tech.*, 11, 1433–1448, 15 1994.
- Beard, N., Straneo, F., and Jenkins, W.: Noble Gases Trace Greenland's Surface and Submarine Melt in the Ocean, *Geophys. Res. Lett.*, submitted, 2015.
- Box, J. E., Yang, L., Bromwich, D. H., and Bai, L.-S.: Greenland Ice Sheet Surface Air Temperature Variability: 1840–2007, *J. Climate*, 22, 4029–4049, doi:10.1175/2009JCLI2816.1, 20 2009.
- Chandler, D. M., Wadham, J. L., Lis, G. P., Cowton, T., Sole, A., Bartholomew, I., Telling, J., Nienow, P., Bagshaw, E. B., Mair, D., Vinen, S., and Hubbard, A.: Evolution of the subglacial drainage system beneath the Greenland Ice Sheet revealed by tracers, *Nat. Geosci.*, 6, 1–4, 2013.
- 25 Chauché, N., Hubbard, A., Gascard, J. C., Box, J. E., Bates, R., Koppes, M., Sole, A., Christoffersen, P., and Patton, H.: Ice–ocean interaction and calving front morphology at two west Greenland tidewater outlet glaciers, *The Cryosphere*, 8, 1457–1468, doi:10.5194/tc-8-1457-2014, 2014.
- Chu, V. W., Smith, L. C., Rennermalm, A. K., Forster, R. R., Box, J. E., and Reehy, N.: Sediment plume response to surface melting and supraglacial lake drainages on the Greenland ice sheet, *J. Glaciol.*, 55, 1072–1082, doi:10.3189/002214309790794904, 2009.
- 30

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Linking
catchment-scale
subglacial discharge
to glacially modified
waters**

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Chu, V. W., Smith, L. C., Rennermalm, A. K., Forster, R. R., and Box, J. E.: Hydrologic controls on coastal suspended sediment plumes around the Greenland Ice Sheet, *The Cryosphere*, 6, 1–19, doi:10.5194/tc-6-1-2012, 2012.
- Cuffey, K. M. and Patterson, W. S. B.: *The Physics of Glaciers*, 4th Edn., Elsevier, Oxford, 2010.
- 5 Enderlin, E., Howat, I. M., and Jeong, S.: An improved mass budget for the Greenland ice sheet, *Geophys. Res. Lett.*, 41, 866–872, doi:10.1002/2013GL059010, 2014.
- Fountain, A. G. and Walder, J. S.: Water flow through temperate glaciers, *Rev. Geophys.*, 36, 299–328, doi:10.1029/97RG03579, 1998.
- Gladish, C. V., Holland, D. M., Rosing-Asvid, A., Behrens, J. W., and Boje, J.: Oceanic Boundary Conditions for Jakobshavn Glacier, Part I: Variability and Renewal of Ilulissat Icefjord Waters, 2001–14, *J. Phys. Oceanogr.*, 45, 3–32, doi:10.1175/JPO-D-14-0044.1, 2015a.
- 10 Gladish, C. V., Holland, D. M., and Lee, C. M.: Oceanic Boundary Conditions for Jakobshavn Glacier, Part II: Provenance and Sources of Variability of Disko Bay and Ilulissat Icefjord Waters, 1990–2011, *J. Phys. Oceanogr.*, 45, 33–63, doi:10.1175/JPO-D-14-0045.1, 2015b.
- 15 Hellmer, H. H. and Oibers, D. J.: A two-dimensional model for the thermohaline circulation under an ice shelf, *Antarct. Sci.*, 1, 325–336, doi:10.1017/S0954102089000490, 1989.
- Hewitt, I. J., Schoof, C., and Werder, M. A.: Flotation and free surface flow in a model for subglacial drainage, Part 2. Channel flow, *J. Fluid Mech.*, 702, 157–187, 2012.
- Holland, D. M., Thomas, R. H., de Young, B., Ribergaard, M. H., and Lyberth, B.: Acceleration of Jakobshavn Isbræ triggered by warm subsurface ocean waters, *Nat. Geosci.*, 1, 659–664, doi:10.1038/ngeo316, 2008.
- 20 Howat, I. M., Negrete, A., and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets, *The Cryosphere*, 8, 1509–1518, doi:10.5194/tc-8-1509-2014, 2014.
- 25 Inall, M. E., Murray, T., Cottier, F. R., Scharrer, K., and Boyd, T. J.: Oceanic heat delivery via Kangerdlugssuaq Fjord to the south-east Greenland ice sheet, *J. Geophys. Res.-Oceans*, 119, 631–645, doi:10.1002/2013JC009295, 2014.
- Jackson, R. H., Straneo, F., and Sutherland, D. A.: Externally forced fluctuations in ocean temperature at Greenland glaciers in non-summer months, *Nat. Geosci.*, 7, 503–508, doi:10.1038/ngeo2186, 2014.
- 30 Jenkins, A.: A one-dimensional model of ice shelf-ocean interaction, *J. Geophys. Res.*, 96, 671–677, 1991.

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Moon, T. and Joughin, I.: Changes in ice front position on Greenland's outlet glaciers from 1992 to 2007, *J. Geophys. Res.*, 113, F02022, doi:10.1029/2007JF000927, 2008.
- Morlighem, M., Rignot, E., Mouginot, J., Seroussi, H., and Larour, E.: Deeply incised submarine glacial valleys beneath the Greenland ice sheet, *Nat. Geosci.*, 7, 18–22, doi:10.1038/ngeo2167, 2014.
- 5 Mortensen, J., Lennert, K., Bendtsen, J., and Rysgaard, S.: Heat sources for glacial melt in a sub-Arctic fjord (Godthåbsfjord) in contact with the Greenland Ice Sheet, *J. Geophys. Res.*, 116, C01013, doi:10.1029/2010JC006528, 2011.
- Morton, B. R., Taylor, G., and Turner, J. S.: Turbulent Gravitational Convection from Maintained and Instantaneous Sources, *P. Roy. Soc. A*, 234, 1–23, doi:10.1098/rspa.1956.0011, 1956.
- 10 Motyka, R. J., Dryer, W. P., Amundson, J., Truffer, M., and Fahnestock, M.: Rapid submarine melting driven by subglacial discharge, *LeConte Glacier, Alaska*, *Geophys. Res. Lett.*, 40, 5153–5158, doi:10.1002/grl.51011, 2013.
- Nye, J. F.: Water flow in glaciers: jökulhlaups, tunnels and veins, *J. Glaciol.*, 17, 181–207, 1976.
- 15 Palmer, S., Shepherd, A., Nienow, P., and Joughin, I.: Seasonal speedup of the Greenland Ice Sheet linked to routing of surface water, *Earth Planet. Sc. Lett.*, 302, 423–428, doi:10.1016/j.epsl.2010.12.037, 2011.
- Plueddemann, A. J., Kukulya, A. L., Stokoy, R., and Freitag, L.: Autonomous Underwater Vehicle Operations Beneath Coastal Sea Ice, *IEEE/ASME T. Mechatron.*, 17, 54–64, 2012.
- 20 Post, A., O'Neel, S., Motyka, R., and Streveler, G.: A Complex Relationship Between Calving Glaciers and Climate, *EOS Trans.*, 92, 305–312, 2011.
- Rignot, E. and Kanagaratnam, P.: Changes in the velocity structure of the Greenland Ice Sheet, *Science*, 311, 986–990, doi:10.1126/science.1121381, 2006.
- Röthlisberger, H.: Water pressure in intra- and subglacial channels, *J. Glaciol.*, 11, 177–203, 1972.
- 25 Schoof, C.: Ice-sheet acceleration driven by melt supply variability, *Nature*, 468, 803–806, 2010.
- Schumann, K., Völker, D., and Weinrebe, W. R.: Acoustic mapping of the Ilulissat Ice Fjord mouth, West Greenland, *Quaternary Sci. Rev.*, 40, 78–88, doi:10.1016/j.quascirev.2012.02.016, 2012.
- 30 Sciacchia, R., Straneo, F., Cenedese, C., and Heimbach, P.: Seasonal variability of submarine melt rate and circulation in an East Greenland fjord, *J. Geophys. Res.-Oceans*, 118, 2492–2506, doi:10.1002/jgrc.20142, 2013.

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Shepherd, A., Ivins, E. R. A. G., Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H., Bromwich, D. H., Forsberg, R., Galin, N., Horwath, M., Jacobs, S., Joughin, I., King, M. A., Lenaerts, J. T. M., Li, J., Ligtenberg, S. R. M., Luckman, A., Luthcke, S. B., McMillan, M., Meister, R., Milne, G., Mougnot, J., Muir, A., Nicolas, J. P., Paden, J., Payne, A. J., Pritchard, H., Rignot, E., Rott, H., Sorensen, L. S., Scambos, T. A., Scheuchl, B., Schrama, E. J. O., Smith, B., Sundal, A. V., van Angelen, J. H., van de Berg, W. J., van den Broeke, M. R., Vaughan, D. G., Velicogna, I., Wahr, J., Whitehouse, P. L., Wingham, D. J., Yi, D., Young, D., and Zwally, H. J.: A Reconciled Estimate of Ice-Sheet Mass Balance, *Science*, 338, 1183–1189, 2012.

Shreve, R. L.: Movement of water in glaciers, *J. Glaciol.*, 11, 205–214, 1972.

Slater, D. A., Nienow, P. W., Cowton, T. R., Goldberg, D. N., and Sole, A. J.: Effect of near-terminus subglacial hydrology on tidewater, *Geophys. Res. Lett.*, 42, 1–8, doi:10.1002/2014GL062494, 2015.

Straneo, F. and Cenedese, C.: The Dynamics of Greenland's Glacial Fjords and Their Role in Climate, *Ann. Rev. Mar. Sci.*, 7, 89–112, doi:10.1146/annurev-marine-010213-135133, 2015.

Straneo, F. and Heimbach, P.: North Atlantic warming and the retreat of Greenland's outlet glaciers, *Nature*, 504, 36–43, doi:10.1038/nature12854, 2013.

Straneo, F., Hamilton, G. S., Sutherland, D. A., Stearns, L. A., Davidson, F., Hammill, M. O., Stenson, G. B., and Rosing-Asvid, A.: Rapid circulation of warm subtropical waters in a major glacial fjord in East Greenland, *Nat. Geosci.*, 3, 182–186, doi:10.1038/ngeo764, 2010.

Straneo, F., Curry, R. G., Sutherland, D. A., Hamilton, G. S., Cenedese, C., Våge, K., and Stearns, L. A.: Impact of fjord dynamics and glacial runoff on the circulation near Helheim Glacier, *Nat. Geosci.*, 4, 322–327, doi:10.1038/ngeo1109, 2011.

Straneo, F., Heimbach, P., Sergienko, O., Hamilton, G., Catania, G., Griffies, S., Hallberg, R., Jenkins, A., Joughin, I., Motyka, R., Pfeffer, W. T., Price, S. F., Rignot, E., Scambos, T., Truffer, M., and Vieli, A.: Challenges to Understanding the Dynamic Response of Greenland's Marine Terminating Glaciers to Oceanic and Atmospheric Forcing, *B. Am. Meteorol. Soc.*, 94, 1131–1144, doi:10.1175/BAMS-D-12-00100.1, 2013.

Sutherland, D. A., Straneo, F., and Pickart, R. S.: Characteristics and dynamics of two major Greenland glacial fjords, *J. Geophys. Res.-Oceans*, 119, 3767–3791, doi:10.1002/jgrc.20224, 2014.

**Linking
catchment-scale
subglacial discharge
to glacially modified
waters**

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Tedstone, A. J. and Arnold, N. S.: Automated remote sensing of sediment plumes for identification of runoff from the Greenland ice sheet, *J. Glaciol.*, 58, 699–712, doi:10.3189/2012JoG11J204, 2012.

Thomas, R., Frederick, E., Krabill, W., Manizade, S., and Martin, C.: Recent changes on greenland outlet glaciers, *J. Glaciol.*, 55, 147–162, doi:10.3189/002214309788608958, 2009.

Turner, J. S.: Buoyancy effects in fluids, Cambridge University Press, 1979.

Van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E., van de Berg, W. J., van Meijgaard, E., Velicogna, I., and Wouters, B.: Partitioning recent Greenland mass loss, *Science*, 326, 984–986, doi:10.1126/science.1178176, 2009.

Werder, M. A., Hewitt, I. J., Schoof, C. G., and Flowers, G. E.: Modeling channelized and distributed subglacial drainage in two dimensions, *J. Geophys. Res.-Earth*, 118, 2140–2158, doi:10.1002/jgrf.20146, 2013.

Xu, Y., Rignot, E., Menemenlis, D., and Koppes, M.: Numerical experiments on subaqueous melting of Greenland tidewater glaciers in response to ocean warming and enhanced subglacial discharge, *Ann. Glaciol.*, 53, 229–234, doi:10.3189/2012/AoG60A139, 2012.

Xu, Y., Rignot, E., Fenty, I., Menemenlis, D., and Flexas, M. M.: Subaqueous melting of Store Glacier, west Greenland from three-dimensional, high-resolution numerical modeling and ocean observations, *Geophys. Res. Lett.*, 40, 4648–4653, doi:10.1002/grl.50825, 2013.

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
Table 1. REMUS missions in Sarqardleq Fjord.

| Mission | Date | Local time at Mission start | Duration (h:mm) | Transect sampling path (m depth) | Distance traveled (km) |
|---------|--------|-----------------------------|-----------------|--|------------------------|
| R1 | 18 Jul | 21:10 | 1:28 | Yo-Yo = 5–90 | 9.00 |
| R2 | 21 Jul | 15:37 | 3:41 | Yo-Yo = 5–50; fixed depth = 50, 70; altitude = 10 m off bottom | 23.11 |
| R3 | 22 Jul | 14:58 | 6:25 | Yo-Yo = 5–55; fixed depth = 60, 70; altitude = 10 m above bottom | 41.36 |
| R4 | 23 Jul | 14:37 | 5:05 | Yo-Yo = 5–50; fixed depth = 60, 70; altitude = 10 m above bottom | 30.93 |
| R5 | 24 Jul | 18:12 | 5:26 | Yo-Yo = 5–60; fixed depth = 40, 55, 70; altitude = 10 m above bottom | 34.91 |

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Table 2. Water mass properties in Sarqardleq Fjord.

| Water mass | Surface Water (SW) | Ilulissat Icefjord Waters (IIW) | Glacially Modified Water 1 (GMW1) | Glacially Modified Water 2 (GMW2) |
|--|--------------------|---------------------------------|-----------------------------------|-----------------------------------|
| Depth range (m) | 0–20 | 20–SF bottom | 35–60 | 50–70 |
| S (PSU) | 21–30.5 | 32.5–33.5 | 30.8–31.5 | 31.1–32.3 |
| θ ($^{\circ}\text{C}$) | 1.5–10 | 0.8–1.5 | 0.75–0.85 | 0.59–0.75 |
| σ_{θ} (ρ_{θ} –1000 kg m $^{-3}$) | 16.0–24.3 | 25.9–26.7 | 24.6–25.1 | 24.8–25.8 |
| Turbidity (NTU) | Low (< 4 NTU) | Low (< 4 NTU) | High (> 9 NTU) | High (> 9 NTU) |
| Origin/formation | Local formation | Disko and Baffin Bay | Local formation | Local formation |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Table 3. Sarqardliup sermia subcatchments and runoff estimates.

| Subcatchment | C1 | C2 | C3 | SS ($\Sigma C1-3$) |
|--|--------------------|------------------|-----------------|----------------------|
| Discharge location | D1 | D2 | D3 | – |
| Bathymetry along catchment terminus | | | | |
| Average depth (m) | 116.4 | 101.5 | 39.9 | – |
| Maximum depth (m) | 150.4 | 131.8 | 49.9 | – |
| Morlighem et al. (2014) (MBM2014) | | | | |
| Catchment area (km ²) | 268.74 | 47.97 | 23.31 | 340.02 |
| Catchment area compared to SS (%) | 79 % | 14 % | 7 % | – |
| Catchment average daily runoff July 2012 $\pm \sigma_{\text{JULY}} (Q_{\text{sg}})$ (m ³ s ⁻¹) | 115.78 \pm 42.59 | 20.62 \pm 7.33 | 9.97 \pm 3.47 | 146.37 \pm 53.26 |
| Average daily July runoff compared to SS (%) | 79 % | 14 % | 7 % | – |
| Catchment average daily runoff during the field expedition (DOY 200, 203–206) $\pm \sigma_{\text{JULY}} (Q_{\text{sg}})$ (m ³ s ⁻¹) | 88.70 \pm 42.59 | 16.10 \pm 7.33 | 7.89 \pm 3.47 | 112.69 \pm 53.26 |
| Bamber et al. (2013) (BBM2013) | | | | |
| Catchment area (km ²) | 402 | 42 | 9 | 453 |
| Catchment area compared to SS (%) | 89 % | 9 % | 2 % | – |
| Catchment average daily runoff July 2012 $\pm \sigma_{\text{JULY}} (Q_{\text{sg}})$ (m ³ s ⁻¹) | 171.01 \pm 64.27 | 17.47 \pm 6.40 | 3.72 \pm 1.36 | 192.20 \pm 71.75 |
| Average daily July runoff compared to SS (%) | 89 % | 9 % | 2 % | – |
| Catchment average daily runoff during the field expedition (DOY 200, 203–206) $\pm \sigma_{\text{JULY}} (Q_{\text{sg}})$ (m ³ s ⁻¹) | 122.83 \pm 64.27 | 14.08 \pm 6.40 | 3.05 \pm 1.36 | 139.96 \pm 71.75 |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.




Figure 1. The Sarqardleq Fjord/Sarqardliup sermia outlet glacier system in West Greenland. Modified from NunaGIS 1 : 100 000 map (Asiaq, Greenland Survey). Sill locations shown in red. Figure 3 location shown in red box.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 3. July 2012 Survey of Sarqardleq Fjord. Sarqardleq Fjord bathymetry (10m colored contours below sea level within fjord) and Morlighem et al. (2014) bedmap (10m colored contours above and below sea level outside of fjord) are shown. The Sarqardliup sermia front position and coastline from a 19 June 2012 Landsat image are mapped in red and black line, respectively. Depth measurements collected during July 2012 field operations used to create the Sarqardleq Fjord bathymetry are plotted as grey dots over the contoured bathymetry. RE-MUS transects R1–R5 are shown in black, with LBL transponders mapped with red triangles. Subglacial subcatchments C1, C2, and C3 dividing lines from MBM2014 analysis are mapped in dashed blue line, with the location of D1, D2, and D3 subglacial discharge channels along the submerged terminus shown with thin black arrows. CTD casts are shown with diamonds: white diamonds are CTD casts along R1 used in REMUS cross-calibration, and the blue, gold, and grey diamonds are CTD casts 1, 2, and 3 that were taken along R5 within GMW1, GMW2, and the region between GMW1 and GMW2 (outlined in blue, gold, and white, respectively). Three proglacial stream entries to Sarqardleq Fjord are shown along the northeast and southwest fjord coastlines with thick black arrows.

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

| | |
|--------------------------|--------------|
| Title Page | |
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |
| ◀ | ▶ |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |
| Printer-friendly Version | |
| Interactive Discussion | |



Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.




Figure 4. Select REMUS Across-Fjord Sections. θ ($^{\circ}\text{C}$), S (PSU), and turbidity (NTU) sections along REMUS lines R5 (**a–c**), R3 (**d–f**), and R1 (**g–i**) from 0 to 100 m depth. Sections are oriented looking away from the terminus, with the southwestern end of the section on the left. Across-fjord transect distance is plotted as horizontal distance along section, with 0 km located at the intersection of the REMUS section with an along-fjord line running from D1 to the southwestern LBL transponder along R1 (Fig. 3). GMW1 and GMW2 regions identified by black ellipses, and labeled in blue and gold, respectively in (**a**)–(**c**). Isotherms plotted in grey, REMUS mission tracks shown in white (Table 1), and bathymetry shown in black (Fig. 3).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




Figure 5. Glacially Modified Water in Sarqardleq Fjord. θ ($^{\circ}\text{C}$) **(a)** and S **(b)** profiles for R4 and R5 measurements over the full water-column depth (grey), with the average of R4 and R5 measurements and the ambient fjord waters in black and red, respectively. **(a)** and **(b)** insets show same data from 20–95 m depth over a finer θ or S range, with measurements taken within the GMW1 and GMW2 regions along R4 and R5 (Fig. 3) shown in blue and gold, respectively. θ/S plots of R4 and R5 measurements **(c)** (colors same as in **a** and **b**), with melt and runoff mixing lines. Intersection for melt and runoff mixing lines set to CTD2 properties at grounding line depth (Fig. 6b). Black square along ambient fjord water profile shows θ/S properties at sill depth (70 m). θ/S results for the Jenkins (2011) plume modeling (Table 4) of D1 (blue triangles) and D2 (gold triangles) shown. **(d)** Same data as in **(c)** over finer θ/S range indicated by thin black box in **(c)**.

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.




Figure 6. Turbidity of Glacially Modified Waters. θ ($^{\circ}\text{C}$) and S (PSU) profiles from the regions along R4 and R5 outlined in blue (GMW1 region) (a), gold (GMW2 region) (b), and white (the region between GMW1 and GMW2) (c) in Fig. 3, with turbidity plotted as the color of the point. CTD1 (a), CTD2 (b), and CTD3 (c) are plotted in grey. The GMW region in θ/S space is outlined in purple. The average of all R4 and R5 measurements and the ambient fjord waters are plotted in black and red, respectively. Black square along ambient fjord water profile shows θ/S properties at sill depth (70 m).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Linking catchment-scale subglacial discharge to glacially modified waters

L. A. Stevens et al.




Figure 7. Sarqardliup sermia catchments and discharge. **(a)** Estimated Sarqardliup sermia catchment (thick black line) and sub-catchments C1, C2, and C3 (dashed black line) from the MBM2014 analysis over Morlighem et al. (2014) bedmap (filled contours) and ice sheet surface (magenta contours). BBM2013 catchment and subcatchment outlines in thick solid and dashed grey lines, respectively. Ice sheet margin and coastlines shown in red and blue, respectively. RACMO2.3 11 km resolution grid points shown with white diamonds. Sarqardleq fjord bathymetry and outer Sarqardleq fjord CTD positions (green triangles) and depth measurements also shown. **(b)** Daily C1, C2, and C3 subcatchment MBM2014 RACMO2.3 discharge estimates (red, blue, and black lines, respectively) and daily average RACMO2.3 temperature (green line) across the Sarqardliup sermia subcatchment C1 for DOY 150–250, 2012. Daily C1, C2, and C3 subcatchment BBM2013 RACMO2.3 discharge estimates in pink, cyan, and grey lines, respectively. Dates of REMUS and CTD sampling from DOY 200–207 marked by grey bar.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion