

Peer-Review interrupted (04 Mar 2016) Author did not upload Revised Manuscript

Editor Decision: Publish subject to minor revisions (Editor review) (22 Feb 2016) by Marco Tedesco

Comments to the Author:

I would like to thank the authors for submitting the revised version of the manuscript and for addressing the concerns raised by the reviewers.

Before accepting the paper for publication, however, I would like to ask the authors to address the points below.

See answers in blue text below.

1) There are still typos and errors that need to be addressed throughout the manuscript. I am reporting below a list of what I was able to find but I strongly encourage the authors to check again the manuscript before the final submission.

We have checked again, removed remaining errors, and edited some sentences in order to make it more readable.

- Line 25. Add a comma after Here

Done

- Line 41 an should be 'a'

Done

- the last sentence of the abstract mentions runoff that 'contributed directly to global sea-level rise'. How are the authors supporting this statement ?

The gauging station is located by a Fjord and the meltwater is therefore running directly out into the Ocean and contributing to global sea level rise. In line 151, under section 2.2 "and with a direct outlet into the Kangerlussuaq Fjord" has been added.

- Line 51 'It is those that determine' . Please, rephrase

Done.

- Line 67 add a reference at the end of the sentence:

Now line 69-71. Reference added and the 3% specified to refer to the western part of the ice sheet.

- Line 68 'significant'. I would rather the authors to use this concerning statistical significance.

Now line 72. Changed to: large

- Line 111. Sentence ending with 'melt rates' would benefit of a reference.

Now line 109. *Ngheim et al, 2012* added.

- Line 117. I find the expression 'implying ...' speculative

Now line 115-116. Implying changed to *Indicating*. The fact that the bridge has not been washed away before now is strongly suggesting that the proglacial discharge has not been as high before. Given the lack of historical river stage data form before 2007, we think it is fair to use this as a proxy for highest water stage and thus proglacial discharge rate.

- Line 143. the references should be moved at the end of the sentence

Done.

- Line 189. It is not clear how the 5 km MODIS data were averaged at 100 m.

Now line 198: See the revised manuscript for clarification.

- Line 215 107 --> 7 should be superscript

Done.

- Line 288. an albedo of 0.05 seems unrealistically low and of the same order of magnitude of the MODIS error. Can the authors please add some comments on this?

Now line 303 – 305. We write that the albedo is 0.05 lower than in 2010 (...). Not that the albedo is 0.05.

- Line 293. I don't understand how this sentence fits into the context of the paper.

The sentence is deleted.

- Line 297. energyinput --> energy input

Corrected.

- Line 361. 'a couple of' is too colloquial. Please rephrase

Minor corrections applied to what is now line 372 to 377.

Moreover, the reply to reviewers is full of errors. The authors use the plural for verbs when the subject is singular. There are also several typos which highlight the poor care of the authors in checking their response before submission. I cordially invite the authors to correct those errors and pay more attention in the future about this detail. In this regard, if possible, the authors should submit a point-by-point reply to reviewers comments. This would help to understand how they addressed their comments.

As the lead author, I take full responsibility for this. I was simply not thorough enough with my replies, which have now been revised and improved.

I would encourage the authors to add the figure showing the components of the energy balance for KAN_L and KAN_U, as originally suggested by Reviewer # 1.

Three figures showing the yearly energy balance for the three weather station sites, AWS_L, AWS_M, AWS_U is shown in the supplementary material. There is referred to these figures in line 297 and 298 in the manuscript.

The plot in Fig. 3 stretches below the x-axis. Please correct the plot.

Temperature below -10 °C has been cut out of the plot and is not shown. The plot is very compact and it takes up too much space to include the lowest temperatures without stretching below the x-axis of Fig. 3A. The temperature figure would take up too much space if the entire temperature span in spring and autumn is included.

Lastly, I encourage the authors to, please, show how they addressed the points above in their reply to this comment.

See answers above in blue.

Answer to reviewers

Thanks to the two anonymous reviewers and comments from C. Charalampidis who provided constructive feedback and suggestions for improvement of the manuscript.

We have updated all the calculations, figures and numbers in the text so that they are based on a catchment delineation derived from the subglacial topography as presented in Lindbäck et al. (2015). Given the difference on catchment shape and size, the numbers change. However, the general pattern with the remarkable difference in the discharge response to the energy input between the two years 2010 and 2012 remain the same. In addition to this, the following changes have been made to the manuscript:

- The abstract have been shortened.
- The former supplementary figure is now included in the manuscript as Figure 2.
- The method section 2.5, about the SEB model have been expanded.
- A method description about the firn saturation model have been added as section 2.6.
- The method description of the catchment delineation (section 2.8) have been rewritten.
- Minor changes have been made to the result section.
- The discussion have been edited in order to make a better flow in the manuscript.

Specific answers to Referee #1

Page C1804, line 10, "I assume that the values in TW are not based solely on PDDs but on the full energy budget"?

All data and graphs regarding the Σ PDD has been replaced with calculations from the energy balance model, with the updated catchment. In the previous version, the text was not always clear enough about whether we used the Σ PDD's or the Energy balance model. The data presented in Table 1 was based on the SEB-model and not a PDD calculation.

Page C1804, line 11: "I recommend that the authors provide additional detail about the van As (2011) model ..."

As explained above, we have expanded our method explanation about the energy balance model. This have now been corrected and it should now be clear what we have done. The SEB model description have been expanded, although still kept relatively short due to the overall length of the manuscript. The model is well tested and used in numerous studies and described in detail in the papers cited in section 2.5.

Page C1804, line 15: "It might be useful to show even show the components of the energy budget for AWS_L and AWS_U":

We have now added three figures in the supplementary material showing the energy balance for the three weather stations AWS_L, AWS_M and AWS_U. The figures is accompanied be a description.

Specific answers to Referee #2

Page C2004, line 18: "Mikkelsen et al 2015 acknowledge that ice catchment delineation is a potential source of uncertainty...."

The concerns raised about the catchment by Referee #2 is fair and we believe to have addressed that in the best possible way. We now use a catchment derived from basal topography in Lindbäck et al. (2015) and therefore address the specific comment (3).

In the revised method section, we address the problem with potential for subglacial water piracy (Specific comment (1)) and refer to Lindbäck et al. (2015) that find that Watson River catchment to shift during the season due to differences in subglacial water-pressure. However, the subglacial catchment, along with its hypsometry, remains effectively constant over the melting season. Therefore the calculated melt rates based on results from the energy balance model is considered robust.

Page C2005, line 2: "(2) Multi watershed approach...."

We do not present a multi watershed approach in the manuscript as such (specific comment (2)). However, the results presented in the previous version and the current revised version is based on two different catchment areas and arrive on the same conclusions. This support that the results presented here are robust. In Lindbäck et al. (2015), previous catchment delineations for the Watson River Catchment is compared and at lower elevations (where the melt rates are highest), they differ only little.

Page C2005, line 9: "(3): "(3) Basal topography resolution...."

Lindbäck et al. (2015) represent all the current available basal topography datasets together with a large amount of additional data presented in that study. Therefore, the best available basal topography dataset has been used for the revisions of the calculations.

Page C2005, line 11: (3) "... Furthermore, MODIS data is being used to map lakes and lake drainages and is also being resampled to a 5km grid ..."

See line 227 to 238 for a clarified description:

"...Fifty-two cloud-free MODIS images with an initial resolution of 500 m were sharpened to 250 m and processed to derive the surface area and volume of supraglacial lakes...."

Specific answers to C. Charalampidis

We have no reason to doubt our calculations and the results presented in Table 2 and we are convinced that the differences lie in the point vs. catchment wide study. It is also worth noting that the absolute differences in melt at elevations above the mean ELA is small and the relative difference therefore is very sensitive to these differences.

Answer to the second and third point by C. Charalampidis: We do not include melt from the elevation interval between 1850 and 2050 m a.s.l. in this revised Figure 3F (previously Figure 2F). We do show the calculated melt from that interval on Figure 3D&E and include it as a part of Table 2. The reason for not

including it in the calculated totals we use in the conclusion and Figure 3F is that satellite imagery indicates that free surface water occurred below 2050 m elevation. However, we don't know exactly at what elevation the melt potentially runoff rather than percolating and refreezing. Therefore we still show the calculated melt from the 1850 to 2050 m elevation interval in Table 2 and Figure 3D&E. Again we don't see any reason to doubt the calculations. Even very small melt rates at this elevation sum up to relatively large amounts due to the hypsometric effect on the catchment that result in a large area for this elevation interval.

As explained above we have rewritten parts of the manuscript and believe it is sufficient to meet the comments on this point. We see no problem in having a youtube link in the text; it's an easy and illustrative way to provide valuable information about the river for the readers not familiar with the location. We are after all living in the 21'st century.

1 Extraordinary runoff from the Greenland Ice Sheet in 2012 amplified by
2 hypsometry and depleted firn-retention

3

4 A.B. Mikkelsen^{1,2}, Hubbard, A.L.^{3,4}, MacFerrin, M.⁵, Box, J.E.⁶, Doyle, S.H.⁴, Fitzpatrick, A.⁴,
5 Hasholt, B.¹, ~~_-~~Bailey, H.⁷, Lindbäck, K.⁸ and Pettersson, R.⁸

6

7 [1]{Department of Geosciences and Natural Resource Management, University of Copenhagen,
8 Denmark}

9 [2]{Centre for Permafrost (CENPERM), University of Copenhagen, Øster Voldgade 10,
10 Copenhagen, DK-1350, Denmark-}

11 [3]{Centre for Arctic Gas Hydrate, Environment and Climate, Department of Geology,
12 University of Tromsø, Dramsveien 201, N-9037 Norway-}

13 [4]{~~Centre for Glaciology, Centre for Glaciology,~~ Department of Geography and Earth
14 Sciences, Aberystwyth University, Aberystwyth, SY23 3DB, UK}

15 [5]{Cooperative Institute for Research in Environmental Sciences (CIRES), University of
16 Colorado, Boulder, CO, USA}

17 [6]{Department of Glaciology and Climate, Geological Survey of Denmark and Greenland,
18 Copenhagen, Denmark}

19 [7] {Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research, Periglacial
20 Research Section, 14473 Potsdam, Germany}

21 [8] Department of Earth Sciences, Uppsala Universitet, Villav. 16, 752 36 Uppsala, Sweden:-
22 Department of Earth Sciences, Villav. 16 752 36 Uppsala Universitet, Sweden.

23 Correspondence to: A.B. Mikkelsen (bechmikkelsen@gmail.com)

24 Abstract

25 It has been argued that the infiltration and retention of meltwater within firn across the
26 percolation zone of the Greenland ice sheet has the potential to buffer up to ~3.6 mm of global sea
27 level rise (Harper et al., 2012). Despite evidence confirming active refreezing processes above the
28 equilibrium line, their impact on runoff and proglacial discharge has yet to be assessed. Here, we
29 compare meteorological, melt, firn-stratigraphy and discharge data from the extreme 2010 and 2012
30 summers to determine the relationship between atmospheric forcing and melt runoff at the land-
31 terminating, Kangerlussuaq sector of the Greenland ice sheet which drains into Watson River. The
32 6.8 km³ bulk discharge in 2012 exceeded that in 2010 by 28%, despite only a 3% difference in net
33 incoming melt energy between the two years. This large disparity can be explained by a 10%
34 contribution of runoff originating from above the long-term equilibrium line (up to 1850 m a.s.l.) in
35 2012 caused by diminished firn retention. The 2012 amplified discharge 2012 response was
36 compounded by catchment hypsometry: — the disproportionate increase in the area contributing to
37 runoff as the melt-level rose high into the accumulation area.

38 Satellite imagery and oblique-aerial photographs of-reveal an active extensive network of
39 supraglacial network rivers extending 140 km from the ice margin that confirms active meltwater
40 runoff originating originating well above the equilibrium line. This runoff culminated in three days
41 of-with record discharge of 3,100 m³ s⁻¹ (0.27 Gt d⁻¹) that peaked on 11 July and washed-out the
42 Watson River bridgeBridge. Our findings corroborate meltwater infiltration processes across in the
43 percolation zone though the resulting patterns of refreezing are complex and can lead to spatially
44 extensive, perched superimposed ice layers within the firn. In 2012, such layers extended to an
45 elevation of at least 1840 m and provideding a semi-impermeable barrier to further meltwater
46 storage, and thereby promoting enhanced widespread runoff from the accumulation area of the
47 Greenland ice sheet that contributed directly to proglacial discharge and global sea-level rise.

48

50 The Greenland ice sheet is losing mass at 0.7 mm yr⁻¹ equivalent of global sea-level rise, the
 51 majority of which is attributed to surface ablation that is set to increase under atmospheric warming
 52 (Enderlin et al., 2014; Hanna et al., 2013). Although surface melt water production can be readily
 53 calculated by regional climate models (e.g. Fettweis et al., 2011), such estimates do not equate
 54 directly to sea-level rise, due to the hydrological processes that buffer and store melt on, within and
 55 beneath the ice sheet. ~~It has been argued that retention near-at~~ the ice sheet surface ~~appear-has to~~
 56 ~~have~~ the greatest capacity to offset future sea-level rise, ~~and~~ particularly refreezing across the wet-
 57 snow/percolation zone above the equilibrium line ~~altitude (ELA) is important~~ (Pfeffer et al., 1991).
 58 Within the percolation zone, melt generated at the surface infiltrates and refreezes within the snow-
 59 pack, increasing its density, forming firm and thereby retaining potential runoff (Pfeffer et al., 1991;
 60 Braithwaite et al., 1994). Harper et al. (2012) analysed a series of cores and ground penetrating
 61 radar profiles collected across an 85 km transect above the ~~ELA-equilibrium line~~ at ~69.5°N to
 62 quantify the water storage capacity of the percolation zone. Their analysis revealed repeated
 63 infiltration events in which surface melt penetrated to more than 10 m depth and refroze as
 64 superimposed ice layers. Although the resulting patterns of vertical densification were complex,
 65 they ~~argue-argued~~ ~~proposed~~ that over a ~~period-number~~ of decades such infiltration ~~will fill all of the~~
 66 available pore space ~~and thereby providing~~ a storage sink of between 322 to 1,289 Gt of melt –
 67 equivalent to buffering ~0.9 to ~3.6 mm of global sea level rise.

Commented [AM1]: Rephrased.

68 Below the ~~ELA-equilibrium line~~ in spring, melt water is initially stored within the snow-pack,
 69 but once the pore-space is saturated, it runs off the previous summer's ice surface (Irvine-Fynn et
 70 al., 2011). This runoff either flows directly into the subglacial environment via supraglacial river
 71 networks and moulins, or is temporarily stored in supraglacial lakes. Such lakes can individually
 72 capture up to 10⁷ m³ (≈0.01 Gt) of water and are estimated to cover up to 3% of the ~~western sector~~
 73 ~~of the ice sheet ablation area on the western part of the ice sheet~~ (Box and Ski, 2007; Fitzpatrick et
 74 al., 2014). ~~They~~ ~~hence,~~ ~~these lakes~~ have the capacity to buffer large volumes of water on
 75 timescales from weeks to months, ~~or and if they do not drain, then~~ potentially ~~over-years~~ ~~if they do~~
 76 ~~not drain~~ (e.g. Fitzpatrick et al., 2014; Selmes et al., 2011). Once filled, the lakes contribute directly
 77 to proglacial discharge either by over-flowing into ~~downstream~~ moulins ~~that have advected~~
 78 ~~downstream~~ or by rapid in situ drainage into the subglacial environment (e.g. Das et al., 2008;
 79 Doyle et al., 2013; Tedesco et al., 2013). It ~~has-been-is noted-observed~~ that supraglacial lakes often

Commented [HLB2]: Will, or could?

80 drain in clusters that ~~may could then~~ cause major peaks in proglacial discharge (Doyle et al., 2013;
81 Fitzpatrick et al., 2014). Ice-dammed proglacial lakes also provide a temporary buffer to proglacial
82 discharge ~~and that are known to drain can flood suddenly rapidly~~ (Carrivick and Quincey, 2014;
83 Mikkelsen et al., 2013).

84 Quantifying these water storage mechanisms across the ice ~~surface-sheet~~ is important since the
85 consequence of enhanced melt on mass balance and sea level contribution depends on the fraction
86 of melt that escapes to the ocean. The ~~elevation-area~~ of the ice sheet undergoing melt will ~~rise~~
87 ~~expand to higher elevations~~ under predicted atmospheric warming, and this could force runoff from
88 well within the ice sheet interior and contribute to enhanced sea level rise (Hanna et al., 2008;
89 Huybrechts et al., 2011; Smith et al., 2015). Expansion of the melt area with warming is further
90 amplified by the ice sheet hypsometry. As the ice surface flattens toward higher elevations, a linear
91 increase in the melt level results in a disproportionate gain in the net surface area exposed to melt
92 conditions. If, however, a significant fraction of that melt is subsequently intercepted and stored by
93 local percolation and refreezing within the snow-pack above the ~~ELA-equilibrium line~~, or otherwise
94 at lower elevations in supra- and pro-glacial lakes, then ~~proglacial~~ discharge and sea-level rise ~~will~~
95 ~~be-is~~ buffered on a time-scale of weeks to decades. Although these storage terms have been
96 estimated for the ice sheet (Box and Ski, 2007; Carrivick and Quincey, 2014; Fitzpatrick et al.,
97 2014; Harper et al., 2012; Humphrey et al., 2012), their combined impact on runoff and proglacial
98 discharge in an integrated study has yet to be quantitatively assessed.

99 1.1 The exceptional 2010 and 2012 melt-seasons

100 The record warm Greenland summers of 2010 and 2012 have been ~~studied-documented~~ using
101 regional atmospheric modelling (Tedesco et al., 2013), microclimatological observations (Bennartz
102 et al., 2013; van As et al., 2012), microwave and optical remote sensing (Nghiem et al., 2012; Smith
103 et al., 2015; Tedesco et al., 2011), and in situ data ~~informing an hypsometric analysis~~ (McGrath et
104 al., 2013). In both years, a blocking high pressure ~~system~~, associated with a strongly negative
105 summer North Atlantic Oscillation (NAO) anomaly, was present in the mid-troposphere over
106 Greenland (Hanna et al., 2014). The resulting circulation pattern advected warm southerly winds
107 over the western flank of the ice sheet, forming an insulating heat-bubble over Greenland (Neff et
108 al., 2014) that promoted enhanced surface heating.

Formatted: Indent: First line: 0.5 cm

109 During summer 2010, higher than average near-surface air temperatures in western and south-
110 western regions of the ice sheet led to early and prolonged summer melting and metamorphism of
111 surface snow, significantly reducing surface albedo and thereby enhancing sunlight absorption (van
112 As, 2012; Box et al., 2012; Tedesco et al., 2013). Similarly, ~~in 2012~~ high-near-surface air
113 temperatures and a low surface albedo enabled high melt rates (Ngheim et al., 2012). During 2012,
114 exceptional melt events were concentrated in two periods in mid and late July. On 12 July, a ridge
115 of warm air stagnated over Greenland and melt occurred over 98.6% of the ~~entire~~ surface of the ice
116 sheet – even extending to the perennially-frozen, high-elevation interior ~~ice~~ at the ice divide
117 (McGrath et al., 2013; Nghiem et al., 2012). In the Kangerlussuaq sector, the focus of this study, the
118 11 July 2012 melt-event had a severe and direct hazardous impact with the wash-out and partial
119 destruction of the Watson River ~~bridge~~ Bridge on the 11 July 2012 (<https://youtu.be/RauzduvIYog>),
120 ~~indicating-revealing~~ that proglacial discharge was at its highest stage since the early 1950's when
121 the bridge was constructed. A second phase of exceptional conditions returned in late July 2012
122 when ~~over~~ 79.2% of the ice sheet surface was again exposed to exceptional melt (Nghiem et al.,
123 2012). Bennartz et al. (2013) found that low-level clouds played an important role by increasing
124 near-surface air temperatures via their effect on radiative absorption: ~~Such clouds were sufficiently~~
125 low ~~enough~~ to enhance the downward infrared irradiance whilst ~~being~~ optically thin enough to
126 allow solar radiation to penetrate.

127 These conditions had the capacity to force rapid and extreme ice sheet melt and runoff that was
128 visible from space and in time-lapse camera sequences of, for example, proglacial flooding (Smith
129 et al., 2015) and turbulent plumes active at the fronts of tidewater glaciers (Chauché et al., 2014;
130 Nick et al., 2012). Nevertheless, the challenge of ~~estimating-measuring~~ discharge at marine-
131 terminating glaciers, and the lack of proglacial gauging ~~measurements-stations~~ in Greenland, means
132 that this inference can only be assessed ~~on-at a~~ broad, regional scales using satellite-derived
133 estimates of mass ~~change-balance~~ (e.g. GRACE; Ewert et al., 2012). Hence, ~~these-the years of two~~
134 ~~years-of~~ exceptionally warm atmospheric forcing ~~in 2010 and 2012 provide-present~~ an ideal natural
135 experiment and ~~ease-study-opportunity~~ to assess and quantify the catchment-wide efficacy and
136 spatio-temporal footprint of melt, storage, and runoff processes across the ice sheet.

137 Here, by reference to the two successive extreme melt seasons of 2010 and 2012, we quantify
138 the efficacy of surface melt storage processes across the Greenland ice sheet using a hydrological-
139 budget approach. We compare surface melt with proglacial discharge across a well-defined, land-

Commented [HLB3]:

summer 2012 ?

terminating catchment that drains the Kangerlussuaq (K-transect) sector of the ice sheet. By drawing on satellite imagery, photographs and a series of snow-pits and firm-cores above the equilibrium line, we relate the calculated residual difference in the hydrological budget to the spatial extent and effectiveness of potential storage terms within the percolation zone.

2 Study area & methods

2.1 Study area

We focus on the ~ 12,500 km² hydrological catchment that drains into Watson River from the land-terminating Kangerlussuaq sector on the western margin of the ice sheet. The catchment is 95% glaciated and comprises four main outlet glaciers centred on Russell Glacier (Figure 1). Within this catchment, the ice surface rises ~90 km from the ice margin at 550 m a.s.l. to the mean (1990–to 2010) ELA-equilibrium line altitude (ELA) of 1,553 m a.s.l. (van de Wal et al., 2012; 2015), and extends a further ~150 km across the accumulation area to the ice divide at ~2,550 m a.s.l.

2.2 Proglacial discharge measurements

Proglacial river discharge was gauged near the Watson River bridge in Kangerlussuaq, located 22 km from the ice sheet margin. Due to orographic shielding by Sukkertoppen Ice Cap the Kangerlussuaq region is exceptionally dry, with a mean annual precipitation of 149 mm (Box et al., 2004; van den Broeke et al., 2008). Land surface water losses from evaporation and sublimation further minimise the land area contribution to runoff compared to the ice sheet component (Hasholt et al., 2013). Watson River discharge was determined using the stage/discharge relationship presented in Hasholt et al. (2013). Water stage was recorded by pressure transducers on a stable cross section ~100 m upstream from the bridge. The discharge Q is given by

$$Q = V \times A, \quad \text{Eq. 1}$$

where V represents the mean velocity in the river cross-section and A is the cross-sectional area. The surface velocity (V) was measured by means of a float and converted into mean cross-sectional velocity by applying a reduction factor of 0.95 (Hasholt et al., 2013). The cross-sectional area (A) used for discharge calculations is based on the deepest sounding of the channel bottom after the winter ice melts in spring. The combined uncertainty in the cross-sectional area and velocity

167 measurements is estimated to be 15% (Hasholt et al., 2013). However, here we also conservatively
168 include the possibility of a systematically deeper cross section due to bed erosion within the deepest
169 of the two channels during the runoff season. Therefore we estimate the upper limit in the ~~yearly~~
170 ~~annual~~ cumulative discharge for 2010 and 2012 at +44% and +32% respectively. The instantaneous
171 ~~possible-potential~~ error varies with the discharge rate, and is plotted together with the measured
172 discharge (Figure 2D and E).

173 During the flood event on 11 July 2012 the water level exceeded the previously observed
174 maximum water stage by 1.65 m (15%) and the stage-discharge relationship was extrapolated
175 accordingly. Our stage-discharge relationship was also altered by the partial removal of a road dam
176 (part of the bridge construction), which opened up two new, shallow channels between and south of
177 the two original channels (Figure 3). We measured the cross-sectional area of the two new channels
178 after the flood had subsided, and by combining these ~~with~~ measurements ~~of with estimates of the~~
179 stage from time-stamped time-lapse photographs, we estimate that these new channels were 1.5 and
180 2.5 m deep at peak flow.

181 The surface velocity in these new channels was calculated assuming the conservation of energy
182 in fluids:

$$v = \sqrt{2gh} \quad \text{Eq. 2}$$

183 where v is the surface velocity of the water, g is the gravitational acceleration (9.82 m s^{-2}) and h is
184 the water level. ~~The u~~ncertainty in v for the two new channels is ~~predominantly-mainly~~ attributed
185 to the determination of stage from time-lapse photos, which we conservatively estimate at ~30%.
186 The two original bedrock channels remained intact and we assume that the hydraulic conditions in
187 these channels did not change substantially during the flood event. For the period after the bridge
188 foundation was partially washed out, the ~~amount of~~ discharge in the new channels is added to that
189 calculated based on the stage/discharge relationship for the original channels. We estimate that the
190 formation of the two new channels during the flood event resulted in a ~~low-small~~ relative (i.e. < 3%)
191 contribution to the total discharge ~~and they therefore do not substantially alter our results.~~

192 2.3 Meteorological measurements

193 Automatic weather stations (AWS) are located at three elevations: 732 ~~m a.s.l.~~(AWS_L), 1280
194 ~~m a.s.l.~~(AWS_M) and 1840 m a.s.l. (AWS_U) (see van As et al., 2012). Each AWS, recorded near-

Formatted: Superscript

195 surface (2-3_m) air temperature, humidity, wind speed, upward and downward shortwave and long
196 wave irradiance, and air pressure.

197 2.4 Snow and ice albedo

198 Surface albedo was determined from NASA's Terra Satellite's Moderate Resolution Imaging
199 Spectroradiometer (MODIS) interpolated onto a 5 km grid from 1 May 2010 to 31 September,
200 2012. An 11-day running median was taken to reject noise caused by contrails and cloud shadows
201 (Box et al., 2012). From these data, an albedo time series was formed for the glaciated part of the
202 Watson River catchment area defined as 67 ± 0.2 °N, and west of 44 °W. The data were averaged in
203 100 m elevation intervals ~~on basis of Scambos & Haran (2002).~~ The resulting albedo product was
204 divided into three approximately equal-area bands corresponding to the physiographic regions
205 dominated by surface impurity darkness (1000 to 1450 m a.s.l.), lakes (1500 ~~to~~ 1650 m a.s.l.) and
206 wet-snow (1700 ~~to~~ 1850 m a.s.l.) (Figure 1) (see also Wientjes et al., 2012; Wientjes and
207 Oerlemans, 2010).

Commented [AM4]: Added

208 2.5 Surface energy budget model

209 The surface energy budget (SEB) was calculated daily across the glacierized catchment
210 following van As et al. (2012). The model calculates radiative, turbulent, rain and subsurface
211 (conductive) energy fluxes using data from the three AWS measurements as input, interpolated into
212 the same 100 m elevation bins as the albedo data. The MODIS albedo data were used in the
213 calculation of net shortwave radiation. The sensible and latent energy fluxes were calculated from
214 near surface gradients of wind speed, temperature and humidity using a stability correction. The
215 surface mass balance (SMB) ~~is~~ was calculated as the sum of solid precipitation, surface melt and
216 sublimation. The model was validated against independent K-transect measurements (e.g. van de
217 Wal et al., 2012) and its performance was found to be within 4% of the observed values. The net
218 energy available for melt across the entire glacierized-catchment was determined by integrating the
219 calculated energy flux ($W\ m^{-2}$) for each elevation interval by area. For the purpose of quantifying
220 the potential net melt available for runoff, refreezing and retention parameterisations were disabled.

221 2.6 Firm ~~Saturation-saturation m~~Model

222 Based upon firm core stratigraphy and density measurements at AWS_U, ~~a mass conservation a~~
223 ~~simple~~ model was ~~produced-applied to illustrate-investigate~~ when ~~saturation and~~ horizontal water
224 flow ~~might may-would~~ occur if melt water were not permitted to percolate beneath ~~the massive~~
225 ~~2010-semi-impermeable, superimposed~~ ice layers. Water generated by melt at the surface, minus
226 evaporation/sublimation, fills the available pore-space of the firm beneath and raises the saturated
227 water table level. In situ measurements and/or reasonable ranges were assigned for model input
228 values, including the density of fresh snow, the average depth and density of the packed snow layer
229 above the firm, the density of refrozen ice and the amount of water attributed to sublimation and
230 evaporation. Ten million (10^7) Monte Carlo model iterations were run over the range of input
231 variables to produce 95% confidence intervals of the daily water levels and potential firm saturation
232 dates at AWS_U.

233 2.7 Supraglacial lake drainage

234 To determine the extent and timing of supraglacial lake drainage events within the Watson River
235 catchment, an automatic lake classification was applied to daily MODIS MOD09 imagery
236 following Fitzpatrick et al. (2014). Fifty-two cloud-free MODIS images with an initial resolution of
237 500 m were ~~processed-sharpened to 250 m and processed~~ to derive the surface area and volume of
238 supraglacial lakes. ~~The smallest lake classified was 0.0625 km² which equates to a single 250 x 250~~
239 ~~m pixel.~~ Lake areas ~~are-were~~ classified using an empirically-determined threshold of the
240 Normalised Difference Water Index (NDWI; Huggell et al., 2002). Lake volume ~~is-was~~ derived
241 using a reflective index approach after Box and Ski (2007) calibrated against lake bathymetry data
242 ~~that was-were~~ acquired in 2010 (Doyle et al., 2013), and subsequently validated against in-situ
243 depths from an independent supraglacial lake at 67° N, 48°W, at ~1420 m.a.s.l. (Fitzpatrick et al.,
244 2014). The error in our lake area and depth ~~are-were~~ is an estimated ~~at~~ ±0.2 km² per lake and 1.5 m
245 per pixel respectively. Change in stored volume in each lake was converted to mean discharge rates
246 between cloud-free observations (Figure 2 D ~~and &~~E).

Formatted: Superscript

248 2.8 Catchment delineation

249 A well documented source of uncertainty in calculating runoff stems from the delineation of
250 hydrologically complex watersheds ~~characterised by with~~ rapidly evolving supraglacial stream,
251 river and lake networks (e.g. van As et al., 2012; Fitzpatrick et al., 2014; Smith et al., 2015).
252 Furthermore, ~~the supraglacial drainage system is plays only but a relatively minor part (be it albeit a~~
253 ~~readily observable one) part~~ of the ~~entire water transport water transport story story~~ and ~~the~~
254 subsequent ~~routing of routing of~~ meltwater ~~on injection injected~~ into the ~~basal environment~~
255 ~~subglacial hydrological system~~ via moulins and fractures remains ~~poorly accounted woefully~~
256 ~~unconstrained~~. Here we adopted a novel watershed delineation approach based on ~~the subglacial~~
257 catchment and ~~drained drainage~~ routing determined from ~~subglacial hydropotential hydraulic~~
258 ~~potential~~ analysis ~~(see presented by Lindbäck et al. (, 2015). In their analysis, Lindbäck et al.~~
259 (2015) demonstrate that the ~~subglacial spatial extent footprint and footprint~~ of the Watson River
260 catchment ~~can evolved may shift migrate northwards by and capture up to up to~~ ~30% of the area of
261 ~~the adjacent Isunnguata Sermia catchment, according owing under to variable varying~~ subglacial
262 ~~water hydrological and water~~ pressure conditions ~~over the course of during~~ the melt-season.
263 However, ~~they also the study also note reveals~~ that despite significant hydrological piracy between
264 adjacent catchments, the actual contributing area of the Watson River subglacial catchment, along
265 with its ~~surface~~ hypsometry, remains effectively constant. ~~Moreover, Lindbäck Lindbäck~~ et al.
266 (2015) also demonstrate that across the lower ablation area (500 to 1250 m a.s.l.) where meltwater
267 ~~production~~ rates are highest, ~~this assumption holds the subglacial footprint is fixed~~ even under
268 transient water-pressure conditions. ~~We are thus Hence, we are~~ confident that ~~our the~~ catchment
269 delineation ~~adopted in this study~~, based on ~~a steady state~~ subglacial hydropotential analysis ~~and s,~~
270 ~~and along with~~ the ~~resulting associated melt and runoff estimates calculations,~~ are robust ~~and~~
271 ~~defensible within the errors of given available~~ data-sets ~~used utilised~~.

Commented [AM5]: Two words?

272 2.9 Measurements of firn and snow pack density

273 To assess firn and snow-pack densification, 15 snow-pits and three 7.6-cm-diameter ice cores
274 were obtained from eight sites between 1280 and 1840 m a.s.l. in April 2012. Two cores (#1 and
275 #2) were drilled 10 m apart ~~in the direct vicinity of near~~ AWS_U while core #3 was drilled at a site
276 located 400 m to the south of AWS_U. ~~The e~~Core stratigraphy was analysed at ~1 cm vertical
277 resolution before cores were cut into 10 cm sections and weighed to determine the density profile of

278 the snowpack and firn. A transect of 0.5 to 1 m-deep snow-pits between AWS_M and AWS_U
279 were examined to investigate spatial variations in firn and snowpack density (Figure 1).

280

281 3 Results

282 Near-surface air temperatures from three AWS reveal insightful differences in the temporal and
283 altitudinal distribution of ~~melt~~ energy available for melt between 2010 and 2012. Melt commenced
284 earlier in 2010 with the lowest AWS_L reaching 6°C daily average air temperature by mid-May
285 (Figure 2A). At AWS_L, melt with air temperature 5°C above the seasonal average persisted until
286 15 September. The duration of the 2010 melt-season (119 days) was without precedent for the
287 Kangerlussuaq sector of the ice sheet since 1973 (van As et al., 2012). At the uppermost AWS_U,
288 located ~300 m above the 1991-2009 baseline ELA of 1524 m (van de Wal et al., 2012), above-
289 freezing temperatures did not prevail until 8 July, 2010. Thereafter daily temperatures remained
290 above melting-freezing until September making 2010 exceptional for melt compared to the long-
291 term average.

292 During the 2012 melt season, air temperatures ~~at elevations~~ above the mean ELA-equilibrium
293 line indicated ~~a widespread~~ surface melting from mid-June onwards ~~and including~~ two week-long
294 periods with extreme daily air temperatures at AWS_U of ~~of~~ well above 3 ~~to~~ 4°C (Figure 2A)
295 ~~during, coinciding with~~ high barometric-pressure and ~~associated~~ clear sky conditions. ~~During~~ In
296 the 5-days leading up to the extreme mid-July 2012 melt event, air temperatures at AWS_M and
297 AWS_U were within 1°C despite ~~being separated by~~ 70 km horizontal and ~~having~~ 500 m elevation
298 vertical separation. Hence, from mid June ~~and~~ throughout July 2012, the environmental lapse rate
299 was exceptionally low— indicating that melting conditions likely prevailed across an extensive, the
300 relatively flat ~~lower~~-accumulation area. By ~~the~~ 12 July, surface melting extended across the entire
301 accumulation area up to the topographic peak of the ice sheet ice sheet divide, and, indeed, the
302 entire ice sheet including Summit Camp and the NEEM drill site where wet snow conditions halted
303 airborne ski-equipped CH130 operations (McGrath et al., 2013; Nghiem et al., 2012). Below 1000
304 m a.s.l., the mean 2012 summer air temperatures were in contrast 0.75°C lower than in 2010,
305 though still higher than the long-term mean. This, in part is ~~partly~~-explained by the delayed 2012
306 melt onset that; commenced in late May 2012 (Figure 2A).

Commented [KL6]: ice divide?

307 ~~Somewhat surprisingly, (The net cumulative energy available for surface melt across the~~
308 ~~glacierized portion of the~~ catchment are virtually equivalent by the end of ~~the~~ 2010 and 2012 ~~melt-~~
309 ~~seasons--summers~~ despite ~~contrasting--quite different prevailing melt season development weather~~
310 ~~conditions~~ (Figure 2B). The total energy available for melt ~~across the catchment~~ in 2010 and 2012
311 calculated ~~using--from~~ the SEB model ~~for the catchment up--up~~ to an elevation of 1840 m a.s.l. was
312 ~~just--only~~ 3% less in 2010~~2~~, compared to 2012~~0~~ (Table 1; ~~.)--See also the supplementary material for~~
313 ~~yearly energy balances for the three weather station sites for 2010 and 2012 respectively).~~

314 MODIS albedo time-series (Figure 2C) binned into three elevation bands equating to the extent
315 of the dark-, lake- and wet-snow zones, respectively (Figure 1) exhibit complex patterns of change
316 through space and time. ~~In~~ 2012, the albedo decline lags behind 2010 (Figure 2C) due to the early-
317 ~~May melt--season~~ onset in ~~May,~~ 2010 promoted by low 2009/2010 winter snow accumulation (van
318 As et al., 2012). By mid-June, albedo across the dark zone for both years declined to 0.4. For the
319 remainder of the melt season, the 2010 dark zone albedo was ~~~0.05~~ lower than in 2012 (Figure 2C),
320 consistent with warmer temperatures and enhanced melt at low elevations ~~during the--in summer~~
321 2010 ~~melt--season~~. Across the lake and wet-snow zones, a similar pattern of albedo decline is
322 observed up until mid-June. From this time onward, in contrast to the dark zone, it is ~~the~~ 2012
323 albedo that is consistently ~~and~~ as much as 0.2 lower than 2010, with the exception of a week--long
324 ~~period, --snow fall albedo reset on 5 August 2012~~ when albedo was reset due to snow-fall on ~~5~~
325 ~~August 2012.~~ Enhanced black carbon deposition from North American wildfire may have played a
326 ~~key role in driving the exceptionally low albedo at high elevations in 2012 (Keegan et al., 2014).~~

327 The seasonal evolution of daily Watson River discharge and catchment-integrated melt varies
328 considerably between 2010 and 2012 (Figure 2D to F). In 2010 the integrated melt and ~~proglacial~~
329 discharge increased at a ~~slower--lower~~ rate than in 2012, despite higher cumulative energy input
330 aided by ~~high--elevated~~ temperatures ~~and--and--combiend with~~ lower albedo. ~~Mean daily 2012~~
331 integrated discharge ~~in 2012~~ peaked at $3100 \text{ m}^3 \text{ s}^{-1}$ (equivalent to $\sim 0.27 \text{ km}^3 \text{ d}^{-1}$; Figure 4E) in mid-
332 July, ~~and which--that~~ washed-out Watson River bridge. With lower temperatures during the week
333 commencing ~~the~~ 15 July, melt and discharge dropped to below 2010 levels but returned to high
334 values ~~(--of at least 1500 m³ s⁻¹--)~~ for 11 days ~~starting on--from~~ 26 July, 2012, coincident with the
335 second phase of exceptionally warm conditions. ~~By the end of the melt-season, (The annual--final~~
336 total ~~annaul~~ discharge in 2012 of 6.8 km^3 ~~--15/+32%--~~ exceeded ~~the--that 2010 total~~ of 5.3 km^3 ~~in~~
337 ~~2010~~ by $\sim 28\%$.

Commented [KL7]: Error bars? Maybe would make the editor more happy.

338 Throughout the 2010 melt-season ~~it becomes apparent that there is~~, a steady increase in the
339 ~~residual difference~~ between calculated ~~integrated~~ melt across the catchment and cumulative
340 ~~proglacial discharge~~, ~~is apparent which which~~ by the end of the season equates to 33% (~1.8 km³)
341 of residual melt retained (R') within the catchment (Figure 2F) ~~compared to the measured discharge~~.
342 In the period leading up to 11 July, 2012, a similar ~~increase in residual R'~~ as ~~compared to 2010~~
343 ~~suggests indicates substantial~~ meltwater storage ~~within the catchment~~. However, ~~After~~ 11 July
344 2012 ~~however~~, the residual R' drops by 40% ~~from more than equating to~~ 1 km³ ~~of bulk discharge~~
345 ~~released~~ within 5 days. ~~Throughout the remainder of the summer, R' and reduces further~~ ~~diminishes~~
346 ~~by the end of the season so~~ ~~indicating that only~~ ~ 0.2 km³ of meltwater is ~~retained retained in the~~
347 ~~catchment, and that meltwater retention after 11 July 2012 was limited by the end of the melt-~~
348 ~~season. This contrasting catchment response to forcing between the two years is nicely~~
349 ~~demonstrated by plotting The plot of cumulative energy input versus cumulative discharge in for~~
350 2010 and 2012 (Figure 4) ~~demonstrates a contrasting catchment response to varying surface energy~~
351 ~~budget between the two years. The resulting slope of the cumulative measured discharge versus~~
352 ~~cumulative calculated energy input energy forcing against discharge response is considerably~~
353 steeper in 2012 than ~~in~~ 2010. Hence, for a given energy input, there ~~was is~~ a ~~disproportionately~~
354 ~~higher larger catchment runoff and Watson River~~ discharge response in 2012 compared to 2010,
355 particularly ~~so~~ during the 11 to 14 July 2012 ~~melt event flooding when the discharge response to the~~
356 ~~energy input is even stronger~~.

357 ~~Table 2 lists calculated The melt totals from for different each~~ elevation bands ~~along~~ with bulk
358 Watson River discharge and their difference ~~are listed in Table 2 s~~. Below the ~~long-term mean~~
359 ~~ELAELA of 1550 m~~, 2010 and 2012 ~~calculated melt total have roughly equivalent are within~~ (7%
360 ~~difference) calculated melt of each other~~. By contrast, ~~above the ELA at in the two~~ elevations ~~bands~~
361 ~~up to 1550 -- 1850 m and 1850 -- 2050 m a.s.l., the~~ calculated melt was 97 and 232% ~~respectively~~
362 larger in 2012 compared to 2010 ~~for the elevations 1550 to 1850 m a.s.l. and 1850 to 2050 m a.s.l.~~
363 ~~respectively (only only melt up to 1850 m a.s.l. is included in Figure 2F and 3)~~. Despite this, the
364 ~~absolute~~ difference in total calculated melt between the two years, ~~was is~~ still ~~within only~~ 2%,
365 ~~depending on the elevation band to which melt is included. Y yet~~, the difference in measured
366 proglacial discharge between the two years ~~peaks at is~~ 28%. Thus, the runoff response to ~~surface~~
367 ~~energy input atmospheric forcing is again demonstrated to be was significantly more pronounced~~
368 ~~higher~~ in 2012, reflected in the larger residual between calculated melt and measured proglacial

369 discharge (Figure 2F), ~~and further illustrated by the contrasting discharge response to energy flux~~
370 ~~compared to 2010 (Figure 4).~~

371 Examination of the ~~The timing-timing between~~ of catchment-integrated melt and ~~Watson River~~
372 proglacial discharge (Figure 2D and E) ~~demonstrates-reveals~~ that meltwater routing through the
373 glacial and proglacial system has a lag of between 1 to 5 days ~~during over~~ each melt-season. In June
374 2012, the proglacial discharge response to melt was dampened and delayed. Prior to the 11 July
375 2012 extreme melt and discharge, the integrated modelled melt closely resembles the proglacial
376 discharge hydrograph but with a ~3 day lag. Henceforth, ~~through-during the remainder of~~ July and
377 the beginning of August 2012, there is a significantly shorter lag between discharge responses ~~eds~~ to
378 melt production ~~with a shorter lag~~. The implication here is that once local meltwater production had
379 been mobilised, even at high elevations above the ~~ELA-equilibrium line~~, the resulting runoff transits
380 ~~within 3 days~~ through a drainage network up to 160 km ~~distant long from the gauging station within~~
381 3 days, eventually thereby contributing to the proglacial discharge peak. Such rapid transit times a
382 drainage system imply with supra- and sub-glacial ~~mean-flow transit velocities-velocities >in excess~~
383 of 2 km h⁻¹ (~0.6 m s⁻¹) through what has to be may be considered an efficient - linked - ~~drainage~~
384 system. These results are on-comparison-comparable to similar transit velocities derived from
385 tracer-experiments conducted up to 57 km from the ice margin in 2011 (Chandler et al., 2013). The
386 second phase of intense melt, commencing on 26 July 2012 ~~resulted-was followed by in~~ a rapid rise
387 in proglacial discharge with a lag of just 2 days. Peak melt during this period occurred on 3 August
388 2012 with the associated peak in proglacial discharge occurring ~~two days later on the 5 August 2012~~.
389 The onset of discharge abatement ~~is-was~~ concurrent with declining air temperatures from 6 August
390 2012 onwards.

391 The release of water stored in supraglacial lakes accounts for a ~~very~~-minor component of
392 proglacial discharge. In 2012 the majority of lake drainages occurred well before any peaks in
393 proglacial discharge (Figure 2E and F). The calculated mean drainage rate of <100 m³ s⁻¹ for 2012
394 ~~clearly~~ indicates that the volume of lake drainage water contributed less than 2% of the total bulk
395 discharge (Figure 2 D&E). The maximum short-term contribution from lake drainage (0.10 km³)
396 occurred on 23 June 2012 with the ~~concurrent-synchronous~~ drainage of a local cluster of five lakes
397 (Figure 2E). Over the following week, approximately 70% of all ~~the~~-water stored in supraglacial
398 lakes across the entire catchment was released (Figure 2E), which ~~could have-could have potentially~~
399 accounted for ~~as-much-as~~ half of the Watson River discharge. However, this ~~synchronous~~/multiple

400 lake drainage event occurred ~12 days before the proglacial discharge peak of 11 July 2012.
401 Supraglacial lakes ~~can empty drain~~ in as little as 2 hours (Das et al., 2008; Doyle et al., 2013) and it
402 is ~~very~~-likely that this ~~stored~~ water ~~stored in supraglacial lakes exited discharged out of~~ the
403 catchment well before 11 July. One ~~small~~ ~0.02 km³ lake drainage event between 5 and 8 July
404 ~~would would have likely to have~~ contributed ~~some~~ ~2% to the extraordinary discharge measured
405 between July 10 and 14 (0.9 km³).

406 Analysis of MODIS and Landsat imagery ~~reveal indicate~~ that no proglacial ice-dammed lakes
407 within the catchment drained prior to the ~~mid-July flooding~~ event, including one that appears to
408 drain regularly in August/September each year (~~Mikkelsen et al., 2013~~). On 11 September 2010 and
409 12 August 2012, ~~the a~~ partially filled proglacial lake ~~described in Mikkelsen et al., (2013)~~ did drain
410 (~~described in Mikkelsen et al., 2013~~), and even though ~~it is evident it is recorded~~ in the Watson
411 River hydrography, the net contribution to ~~the~~ proglacial discharge is minor (Figure 2D and E).

413 4 Discussion

414 Our ~~observations analysis reveals show~~ that ~~even though~~ the net atmospheric forcing
415 represented by the ~~total~~ incoming energy flux for 2010 and 2012 was ~~equivalent similar~~(Figure 4),
416 ~~yet~~ the ensuing runoff response was markedly different (Figure 4). Widespread melt in 2010 has
417 been ascribed to atmospherically-sourced heating coupled with ~~the a strong~~ albedo feedback
418 promoted by low winter snowfall and early melt onset (Tedesco et al., 2011; Box et al., 2012; van
419 As et al., 2012). Yet low albedo and high air temperatures alone ~~cannot do not~~ explain the 28%
420 increase in discharge in 2012 compared to 2010. ~~MODIS-Our~~ analysis ~~also~~ confirms that the release
421 of stored water from supraglacial lakes played a relatively minor role in ~~the~~ peak and total
422 proglacial ~~hydrograph discharge~~ in 2012 (Fig 2D and E). At most, the supraglacial lake contribution
423 to the 11 July 2012 peak discharge of 3,100 m s⁻¹ was ~2%. Our results indicate that only a
424 relatively small proportion of the total melt generated at the surface ~~is was~~ stored in supra- and pro-
425 glacial lakes and that the buffering effect of lakes on runoff and discharge is ~~thus therefore~~ limited
426 (Figure 2D and E). That is not to dismiss the key role of supraglacial lakes in ice sheet hydrology,
427 since it is the ~~ephemeral critical~~ storage of ~~large volumes of surface~~ meltwater in them that ~~initiate~~
428 ~~enable the critical volume required to initiate and propagate~~ new hydrofractures ~~and moulins and~~
429 ~~allow them to propagate~~ to the bed - ~~which eventually develop into moulins~~ (Krawczynski et al.,

Commented [AM8]: No hyphen between words?

430 2009; Doyle et al., 2013; Tedesco et al., 2013). ~~In this manner,~~ ^sSupraglacial lakes are hence key-a
431 prerequisite to ~~creat~~establishing ~~ng~~ efficient englacial pathways ~~for discharging~~ ~~for injecting~~
432 surface water into the subglacial environment ~~over the melt season~~ (Das et al., 2008; Doyle et al.,
433 2013).

434 We ~~infer~~ invoke three mutually compatible explanations for the exceptional discharge response
435 observed in 2012: 1) ~~that~~ significant melt occurred above the equilibrium line in addition to as well
436 ~~as~~ below ~~the ELA~~it, 2) ~~that~~ ice surface hypsometry amplified the total melt originating from the
437 accumulation zone by disproportionately increasing the contributing area ~~when as~~ melt-levels rose
438 ~~above the ELA~~, and; 3) ~~there was reduced~~ firm-retention and storage capacity was reduced within
439 the accumulation zone, ~~which that~~ thereby promoting widespread ~~large-scale~~ runoff. It is
440 significant that such a large runoff contribution from the percolation zone could only have been
441 attained if firm-retention capacity was either filled or otherwise severely reduced in 2012 and it is
442 this hypothesis that herein forms the central tenet of our discussion. In support of this we present
443 three additional lines of evidence: A) snow pit observations and ice-firm core stratigraphy acquired
444 in April 2012 from the percolation zone, B) observations of surface water networks obtained from
445 satellite imagery and oblique photographs in the vicinity of AWS_U ~~from the vicinity of AWS_U~~
446 (Figure 6), and; C) results of our SEB-modelling experiments where total integrated melt is
447 assumed to runoff without any retention or refreezing.

448 ~~Our~~ The core stratigraphic analysis (Figure 5A to C) reveals significant-a number of perched
449 superimposed ice layers that would be impermeable and hence potentially could be capable of
450 blocking surface meltwater infiltration into deeper unsaturated ~~firm-pore-space layers across the~~
451 percolation zone. In addition to the shallow ice-firm cores presented (Figure 5), a persistent and
452 continuous decimetre-thick layer of refrozen, superimposed ice was also observed in 15 snow pits
453 dug along a transect ~~from~~ extending from ~~below the ELA equilibrium line (1500 m a.s.l.)~~ to
454 AWS_U (Figure 1). Severely reduced firm-retention due to such a superimposed, perched ice lens is
455 further supported by energy-balance-mass conservation modelling of the near-surface water table at
456 AWS_U (Figure 5D). Here, two potential sets of blocking layers at different levels within the
457 snow-pack equate to the thick superimposed ice lenses observed in the firm cores acquired at
458 AWS_U (Figure 5A to C). For the shallowest of these scenarios, melt and retention calculations
459 predict complete saturation and free surface water available for active runoff by 11 July 2012. ~~Our~~
460 These results are consistent with a recent study by Machguth et al. (2015) who also demonstrate

Commented [KL9]: Rephrase that is, there is, it is...
(considered bad English)

461 reduced meltwater retention across the percolation zone of western sector of the Greenland ice
462 sheet.

463 Evidence for firm saturation and active surface runoff are furnished independently by the
464 identification of an active supraglacial channel network in Landsat satellite imagery and from
465 oblique photographs taken 13 August 2012 in the vicinity of AWS_U (Figure 6). ~~In the~~ Landsat
466 imagery ~~indicates that~~ ~~ry~~, wet snow, meltwater channels and lakes can be identified up to at least
467 1750 m a.s.l. on 23 June 2012, and an active stream network ~~up to~~ at least 1840-1800 m a.s.l. from 5
468 July 2012 onwards. In early August, 2012 an active channel network was confirmed first-hand
469 during a scheduled maintenance visit to AWS_U (Figure 6B and C). ~~The~~ That a well developed
470 supraglacial hydrological network ~~that is~~ clearly evident-observed well above the long-term ELA
471 equilibrium line in the period leading up to the 2012 peak discharge event confirms ~~the~~ the
472 assessment of firm retention conditions and the snow-pack modelling presented here. Moreover, The
473 oblique-aerial photos of stream networks to 1840 m a.s.l. provide unequivocal-clear evidence ~~for~~ of
474 widespread surface runoff from the percolation zone across this the region western sector of the
475 Greenland ice sheet.

476 ~~MODIS analysis confirms that the release of stored water from supraglacial lakes played a~~
477 ~~relatively minor role in the peak and total proglacial hydrograph in 2012 (Fig 2D and E). At most,~~
478 ~~change in supraglacial lake storage contributed just only 2 % to the 11 July 2012 peak discharge of~~
479 ~~3,100 m³ s⁻¹ was ~2%. Our results indicate that only a relatively small proportion of the total melt~~
480 ~~generated at the surface is stored in supra- and pro-glacial lakes and that the buffering effect of~~
481 ~~lakes on runoff and discharge is therefore limited (Figure 2D and E). That is not to dismiss the~~
482 ~~importance of supraglacial lakes in ice sheet hydrology, since it is the ephemeral storage of surface~~
483 ~~meltwater in them that enable the critical volume required to initiate and propagate new~~
484 ~~hydrofractures and moulins to the bed (Krawczynski et al., 2009). In this manner, supraglacial~~
485 ~~lakes are key to creating efficient englacial pathways for discharging surface water into the~~
486 ~~subglacial environment over the melt season (Das et al., 2008; Doyle et al., 2013).~~

487 If ~~forecasted-predicted~~ future atmospheric warming is realised, then the combined impact of
488 reduced firm retention capacity and ice sheet hypsometry will become increasingly apparent through
489 amplification of runoff and discharge response with interior melting. If, as we hypothesise, the
490 extraordinary 2012 discharge was substantially in-partly derived from runoff originating above the
491 ELA-equilibrium line due to an impermeable, superimposed ice lens that formed during previous

492 warm summers, then the 2012 record-warm event itself will ~~drive-lead to~~ the formation of even
493 thicker, superimposed ice layers extending yet further into the interior (~~i.e.g. McGrath et al.,~~
494 ~~2013~~). Hence, we infer a strong positive feedback where a disproportionate and amplified runoff
495 response to future melt events leads to yet more abrupt and severe proglacial discharge, as the 11
496 July 2012 flood~~ding~~ documented here.

497 In light of these findings, the firn-buffering mechanism proposed for the EGIG line some 120
498 km north of our study area and extrapolated across the entire ice sheet by Harper et al. (2012) would
499 appear to be somewhat diminished, at least in the Kangerlussuaq sector. Based on their data and
500 analysis (Figure 2B and 3C in Harper et al., 2012) and assuming an equivalent location, our
501 AWS_U site, located 50 km beyond, and 300 m above the ELA, should have had a buffering
502 capacity equating to a fill-depth of between 2 m and 10 m of ~~melt-water equivalent-of-melt~~. In July
503 2012, ~~at up to and including~~ AWS_U at 1840 m a.s.l. this was not the case and ~~under~~-saturated
504 ~~snow-pack~~ conditions, ~~melt was~~ forced ~~melt~~ to runoff from the percolation zone into an ~~active~~
505 ~~well-developed supraglacial hydrological-river~~ network ~~thereby-that~~ directly contribut~~ed~~
506 proglacial discharge and sea-level rise. ~~The next decade will reveal if 2010 and 2012 were~~
507 ~~exceptions or are part of an emerging new trend. The three years subsequent to the 2012 melt and~~
508 ~~runoff extreme, i.e., 2013-2015, have been marked by low temperatures, reduced melting and~~
509 ~~anomalously high accumulation which will have, to some extent, recharged the buffering capacity~~
510 ~~of the lower accumulation area. Either way, it will be critical to understand the future runoff~~
511 ~~response to variable atmospheric forcing and to determine what portion of the melt generated is~~
512 ~~intercepted and stored and what fraction contributes directly to proglacial discharge and global sea-~~
513 ~~level rise.~~

Commented [KL10]: i.e. = that is or e.g. for example?

~~A key implication of our study is that expected climate warming will change the limit of upper elevation ice sheet runoff to a higher level sooner. The hypsometric effect that amplifies runoff by the contributing area increasing exponentially with elevation (Figure 4) combined with efficient drainage (2-3 day transit times for water; (results here and in Smith et al. 2015), we may thus expect the ice sheet sea level contribution to be faster than inferred by Harper et al. (2012).~~

Formatted: Heading 11, Left, Tab stops: Not at 1.25 cm + 2.5 cm + 3.75 cm + 4.99 cm + 6.24 cm + 7.49 cm + 8.74 cm + 9.99 cm + 11.24 cm + 12.49 cm + 13.74 cm + 14.98 cm

Commented [KL11]: I don't understand this sentence. What do you mean? Will the limit go to a higher elevation? Refrase.

Commented [KL12]: Rephrase. Messy sentence.

5 Conclusions

Comparison of melt and discharge across the Kangerlussuaq sector in 2010 and 2012 has enabled us to assess, ~~resolve~~, and attribute the contrasting runoff response of the Greenland ice sheet to extreme atmospheric forcing. The bulk discharge of 6.8 km³ ~~measured and flooding of in~~ Watson River in 2012 was unprecedented since the Kangerlussuaq ~~B~~bridge was constructed in the early 1950's and exceeded the previous record set in 2010 by ~28%. Throughout the 2010 melt-season, there was a steady increase in the residual difference between calculated melt across the catchment and cumulative proglacial discharge, which by the end of the season equated to 33% (~1.8 km³) melt retained within the catchment up to an elevation of 1850 m a.s.l. In the period ~~leading~~ up to 11 July 2012 a similar pattern of storage indicates significant ~~meltwater-catchment~~ retention. However, after ~~the~~ 11 July ~~flooding~~ the residual fell by 40% and ~~reduced~~ ~~decreased~~ ~~diminished~~ further by the end of September with only 3% (~0.2 km³) of ~~generated~~ melt ~~generated within the catchment~~ retained. ~~The abrupt change signifies a sudden decrease in retention associated with essentially complete surface snow ablation below areas with snow that became water saturated.~~ Surface melt energy versus proglacial discharge demonstrates an amplified response to melt energy forcing in 2012 as compared to 2010, particularly ~~during after the 11 -- 14~~ July ~~flood~~ing. In 2010 local melting from above the ELA-equilibrium line infiltrated, and was stored within the firn as superimposed ice layers and hence did not contribute to ~~river-proglacial~~ discharge. ~~In~~ ~~By contrast, in 2012 though,~~ our observations-analysis and modelling ~~indicate-reveals~~ severely reduced firn-layer infiltration and retention capacity due an extensive perched, thick and low semi-impermeable ice lens, ~~which that most likely formed in in previous, anomalously warm melt-seasons, including 2010-years (e.g. 2007 and including 2010).~~

Commented [AM13]: Comma after lens?

~~The next decade will reveal if 2010 and 2012 were exceptional melt seasons or are part of an emerging new trend. The three years subsequent to the 2010 and 2012 melt and runoff extremes, i.e., 2013-2015 were marked by low total melt and in some cases anomalously high accumulation.~~

545 ~~The effect will have been to recharge the buffering capacity of the lower accumulation area to some~~
546 ~~degree. Either way, it is critical to understand the ice sheet runoff response to such events to~~
547 ~~determine what portion of the melt generated is intercepted and stored and what fraction directly~~
548 ~~contributes to proglacial discharge and sea level rise. Our results reveal that the firm retention and~~
549 ~~buffering effects that are argued to dominate the percolation zone were much reduced across the~~
550 ~~Kangerlussuaq sector in 2012. This resulted, and that there was in a near-instantaneous runoff and~~
551 ~~proglacial discharge response with a disproportionately greater area of the ice sheet above the ELA~~
552 ~~from above the accumulation area contributing to runoff and thereby contributing directly to global~~
553 ~~sea level rise.~~

Commented [KL14]: Rewrite. Too long sentence, hard to follow.

554 Acknowledgements

555 We thank Dirk van As ~~and Horst Machguth~~ for assistance in the field, ~~advice and during~~
556 ~~preparations on~~ the manuscript. ~~We thank Dirk van As for weather station data, runoff~~
557 ~~calculations, initiating firm investigations, fieldwork logistics, writing and and~~ supervision of
558 Andreas Bech Mikkelsen's PhD project. ~~Likewise Horst Machguth for field assistance.~~ Gratitude
559 also ~~goes~~ to Paul Smeets, Institute for Marine and Atmospheric Research, Utrecht University for
560 providing oblique areal ~~pictures-photographs~~ taken at AWS_U on 13 August 2011. ~~We~~
561 ~~acknowledge~~ ~~We acknowledge financial support from the the~~ Greenland Analogue Project (GAP) –
562 Sub Project A ~~that for funding AWS_U the weather stations~~ and field logistics, the commission
563 on scientific investigations in Greenland, grant no. 07-015998, 09-064628 and 2138-08-0003 and
564 the Danish National Research Foundation founding Centre for Permafrost (CENPERM), funded by
565 the Danish National Research Foundation, DNRF number 100, Department of Geosciences and
566 Nature Resource Management, University of Copenhagen, Denmark for financial support of the
567 discharge measurements. ~~The on-ice weather station in this study was maintained by GEUS and~~
568 ~~Aberystwyth University.~~ ~~J-BBox~~ is ~~here~~ supported by Denmark's "Det Frie Forskningsråd", Nature
569 and Universe grant DFF – 4002-00234. The National Aeronautics and Space Administration
570 (NASA) award NNX10AR76G provided funding for firm table modelling work through the
571 Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder,
572 USA. AF and SD were supported by NERC and Aberystwyth University doctoral scholarships
573 respectively and fieldwork ~~infrastructure~~ was ~~further supported-funded~~ by NERC Projects
574 NE/G005796/1, NE/G010595/1, NE/H024204/1 and a Royal Geographical Society Gilchrist

575 Fieldwork Award. A-H. ~~kindly~~ acknowledges ~~support-salary~~ from the Centre for Arctic Gas
576 Hydrate, Environment and Climate ~~by funding from through~~ the Research Council of Norway
577 (Grant No. 223259).

578 **References**

- 579 Bennartz, R., Shupe, M. D., Turner, D. D., Walden, V. P., Steffen, K., Cox, C. J., Kulie, M. S.,
580 Miller, N. B. & Pettersen, C. (2013). July 2012 Greenland melt extent enhanced by low-
581 level liquid clouds. *Nature*, **496**, 83-86.
- 582 Box, J. E., Bromwich, D. H. & Bai, L.-S. (2004). Greenland ice sheet surface mass balance 1991–
583 2000: Application of Polar MM5 mesoscale model and in situ data. *Journal of Geophysical*
584 *Research*, **109**.
- 585 Box, J. E., Fettweis, X., Stroeve, J. C., Tedesco, M., Hall, D. K. & Steffen, K. (2012). Greenland ice
586 sheet albedo feedback: thermodynamics and atmospheric drivers. *The Cryosphere*, **6**, 821-
587 839.
- 588 Box, J. E. & Ski, K. (2007). Remote sounding of Greenland supraglacial melt lakes: implications
589 for subglacial hydraulics. *Journal of Glaciology*, **53**, 257-265.
- 590 Braithwaite, R. J., Laternser, M. & Pfeffer, T. (1994). Variations of near-surface firn density in the
591 lower accumulation area of the Greenland ice sheet, Pakitsoq, West Greenland. *Journal of*
592 *Glaciology*, **40**, 477-485.
- 593 Carrivick, J. L. & Quincey, D. J. (2014). Progressive increase in number and volume of ice-
594 marginal lakes on the western margin of the Greenland Ice Sheet. *Global and Planetary*
595 *Change*, **116**, 156-163.
- 596 Chandler, D. M., Wadham, J. L., Lis, G. P., Cowton, T., Sole, A., Bartholomew, I., Telling, J.,
597 Nienow, P., Bagshaw, E. B., Mair, D., Vinen, S. & Hubbard, A. (2013). Evolution of the
598 subglacial drainage system beneath the Greenland Ice Sheet revealed by tracers. *Nature*
599 *Geoscience*, **6**, 195-198.
- 600 Chauché, N., Hubbard, A., Gascard, J. C., Box, J. E., Bates, R., Koppes, M., Sole, A.,
601 Christoffersen, P. & Patton, H. (2014). Ice–ocean interaction and calving front morphology
602 at two west Greenland tidewater outlet glaciers. *The Cryosphere*, **8**, 1457-1468. Das, S. B.,
603 Joughin, I., Behn, M. D., Howat, I. M., King, M. A., Lizarralde, D. & Bhatia, M. P. (2008).
604 Fracture propagation to the base of the Greenland Ice Sheet during supraglacial lake
605 drainage. *Science*, **320**, 778-781.
- 606 Das, S., Joughin, I., Behn, M., Howat, I., King, M., Lizarralde, D., & Bhatia, M. (2008). Fracture
607 propagation to the base of the Greenland ice sheet during supraglacial lake drainage.
608 *Science*. 320, 778–781. doi:10.1126/science.1153360, 2008.
- 609 Doyle, S. H., Hubbard, A. L., Dow, C. F., Jones, G. A., Fitzpatrick, A., Gusmeroli, A., Kulesa, B.,
610 Lindback, K., Pettersson, R. & Box, J. E. (2013). Ice tectonic deformation during the rapid
611 in situ drainage of a supraglacial lake on the Greenland Ice Sheet. *The Cryosphere*, **7**, 129-
612 140.
- 613 Doyle, S. H., Hubbard, A. et al. (2015), Amplified melt and flow of the Greenland ice sheet driven
614 by late-summer cyclonic rainfall, *Nat. Geosci.*, doi:10.1038/ngeo2482.
- 615 Enderlin, E. M., Howat, I. M., Jeong, S., Noh, M. J., van Angelen, J. H. & van den Broeke, M. R.
616 (2014). An improved mass budget for the Greenland ice sheet. *Geophysical Research*
617 *Letters*, **41**, 866-872.

Commented [KL15]: Please check all references in text. References are missing!

Formatted: Font: Italic

- 618 Ewert, H., Groh, A. & Dietrich, R. (2012). Volume and mass changes of the Greenland ice sheet
619 inferred from ICESat and GRACE. *Journal of Geodynamics*, **59–60**, 111-123.
- 620 Fettweis, X., Belleflamme, A., Erpicum, M., Franco, B. & Nicolay, S. (2011). Estimation of the sea
621 level rise by 2100 resulting from changes in the surface mass balance of the Greenland Ice
622 Sheet. In: Blanco, J. & Kheradmand, H. (eds.) *Climate Change: Geophysical Foundations
623 and Ecological Effects*. Croatia: InTech.
- 624 Fitzpatrick, A. A. W., Hubbard, A. L., Box, J. E., Quincey, D. J., van As, D., Mikkelsen, A. P. B.,
625 Doyle, S. H., Dow, C. F., Hasholt, B. & Jones, G. A. (2014). A decade (2002–2012) of
626 supraglacial lake volume estimates across Russell Glacier, West Greenland. *The
627 Cryosphere*, **8**, 107-121.
- 628 Hanna, E., Fettweis, X., Mernild, S. H., Cappelen, J., Ribergaard, M. H., Shuman, C. A., Steffen,
629 K., Wood, L. & Mote, T. L. (2014). Atmospheric and oceanic climate forcing of the
630 exceptional Greenland ice sheet surface melt in summer 2012. *International Journal of
631 Climatology*, **34**, 1022-1037.
- 632 Hanna, E., Huybrechts, P., Steffen, K., Cappelen, J., Huff, R., Shuman, C., Irvine-Fynn, T., Wise, S.
633 & Griffiths, M. (2008). Increased Runoff from Melt from the Greenland Ice Sheet: A
634 Response to Global Warming. *Journal of Climate*, **21**, 331-341.
- 635 Hanna, E., Navarro, F. J., Pattyn, F., Domingues, C. M., Fettweis, X., Ivins, E. R., Nicholls, R. J.,
636 Ritz, C., Smith, B. & Tulaczyk, S. (2013). Ice-sheet mass balance and climate change.
637 *Nature*, **498**, 51-59.
- 638 Harper, J., Humphrey, N., Pfeffer, W. T., Brown, J. & Fettweis, X. (2012). Greenland ice-sheet
639 contribution to sea-level rise buffered by meltwater storage in firn. *Nature*, **491**, 240-3.
- 640 Hasholt, B., Mikkelsen, A. B., Nielsen, M. H. & Larsen, M. A. D. (2013). Observations of Runoff
641 and Sediment and Dissolved Loads from the Greenland Ice Sheet at Kangerlussuaq, West
642 Greenland, 2007 to 2010. *Zeitschrift für Geomorphologie*, **57** (Suppl. 2), 3-27.
- 643 Huggell, C., Kaab, A., Haeblerli, W., Teyssie, P. & Paul, F. (2002). Remote sensing based
644 assessment of hazards from glacier lake outbursts: a case study in the Swiss Alps. *Canadian
645 Geotechnical Journal*, **39**, 316-330.
- 646 Humphrey, N. F., Harper, J. T. & Pfeffer, W. T. (2012). Thermal tracking of meltwater retention in
647 Greenland's accumulation area. *Journal of Geophysical Research-Earth Surface*, **117**.
- 648 Huybrechts, P., Goelzer, H., Janssens, I., Driesschaert, E., Fichet, T., Goosse, H. & Loutre, M. F.
649 (2011). Response of the Greenland and Antarctic Ice Sheets to Multi-Millennial Greenhouse
650 Warming in the Earth System Model of Intermediate Complexity LOVECLIM. *Surveys in
651 Geophysics*, **32**, 397-416.
- 652 Irvine-Fynn, T. D. L., Hodson, A. J., Moorman, B. J., Vatne, G. & Hubbard, A. (2011).
653 Polythermal glacier hydrology: a review. *Reviews of Geophysics*, **49**, RG4002.
- 654 Krawczynski, M., Behn, M., Das, S., and Joughin, I. (2009). Constraints on the lake volume
655 required for hydro-fracture through ice sheets, *Geophys. Res. Lett.*, **36**, 1–5,
656 doi:10.1029/2008GL036765.
- 657 Keegan, K. M., Albert, M. R., McConnell, J. R. & Baker, I. (2014). Climate change and forest fires
658 synergistically drive widespread melt events of the Greenland Ice Sheet. *Proceedings of the
659 National Academy of Sciences of the United States of America*, **111**, 7964-7967.

- 660 [Lindbäck, K., Pettersson, R., Hubbard, A., Doyle, S., van As, D., Mikkelsen, A., Fitzpatrick, A.](#)
661 [\(2015\). et al. \(2015\)](#)
- 662
- 663 [Subglacial water drainage, storage, and piracy beneath the Greenland ice sheet. *Geophysical*](#)
664 [Research Letters](#), **42** (18), 7606-7614.
- 665 Machguth, H., MacFerrin, M., van As, D., Box, J., E., Charalampidis, C., Colgan, W., Fausto, R.,
666 S., Meijer, H., A., J., Mosley-Thompson, E. and van de Wal, R., S., W. Greenland meltwater
667 storage in firn limited by near-surface ice formation (2016). *Nature Climate Change*, 1-6.
- 668 McGrath, D., Colgan, W., Bayou, N., Atsuhiko, M. & Konrad, S. (2013). Recent warming at
669 Summit, Greenland: Global context and implications. *Geophysical Research Letters*, **40**,
670 2091-2096.
- 671 Mikkelsen, A. B., Hasholt, B., Knudsen, N. T. & Nielsen, K. (2013). Jökulhlaups and sediment
672 transport in Watson River, Kangerlussuaq, West Greenland. *Hydrology Research*, **44**, 58-67.
- 673 Neff, W., Compo, G. P., Ralph, F. M. & Shupe, M. D. (2014). Continental heat anomalies and the
674 extreme melting of the Greenland ice surface in 2012 and 1889. *Journal of Geophysical*
675 *Research-Atmospheres*, **119**, 6520-6536.
- 676 Nghiem, S. V., Hall, D. K., Mote, T. L., Tedesco, M., Albert, M. R., Keegan, K., Shuman, C. A.,
677 DiGirolamo, N. E. & Neumann, G. (2012). The extreme melt across the Greenland ice sheet
678 in 2012. *Geophysical Research Letters*, **39**, L20502.
- 679 Nick, F. M., Luckman, A., Vieli, A., van der Veen, C. J., van As, D., van de Wal, R. S. W., Pattyn,
680 F., Hubbard, A. L. & Floricioiu, D. (2012). The response of Petermann Glacier, Greenland,
681 to large calving events, and its future stability in the context of atmospheric and oceanic
682 warming. *Journal of Glaciology*, **58** (208), 229-239.
- 683 Pfeffer, W. T., Meier, M. F. & Illangasekare, T. H. (1991). Retention of Greenland Runoff by
684 Refreezing— Implications for Projected Future Sea-Level Change. *Journal of Geophysical*
685 *Research-Oceans*, **96**, 22117-22124.
- 686 Rennermalm, A., Moustafa, S., Mioduszewski, J., Chu, V., Forster, R., Hagedorn, B., Harper, J.,
687 Mote, T., Robinson, D., Shuman, C. & others (2013). Understanding Greenland ice sheet
688 hydrology using an integrated multi-scale approach. *Environmental Research Letters*, **8**,
689 015017.
- 690 Russell, A. J., Carrivick, J. L., Ingeman-Nielsen, T., Yde, J. C. & Williams, M. (2011). A new cycle
691 of jokulhlaups at Russell Glacier, Kangerlussuaq, West Greenland. *Journal of Glaciology*,
692 **57**, 238-246.
- 693 Selmes, N., Murray, T. & James, T. D. (2011). Fast draining lakes on the Greenland Ice Sheet.
694 *Geophysical Research Letters*, **38**, L15501.
- 695 Scambos, T. A., & Haran, T. (2002): An image-enhanced DEM of the Greenland ice sheet. *Ann*,
696 *Glaciol.*, **34**, 291-298.
- 697 Smith, L. C., Chu, V. W., Yang, K., Gleason, C. J., Pitcher, L. H., Rennermalm, A. K., Legleiter, C.
698 J., Behar, A. E., Overstreet, B. T., Moustafa, S. E., Tedesco, M., Forster, R. R., LeWinter,
699 A. L., Finnegan, D. C., Sheng, Y. & Balog, J. (2015). Efficient meltwater drainage through

Formatted: Font: Italic

Formatted: Font: Bold

700 supraglacial streams and rivers on the southwest Greenland ice sheet. *Proceedings of the*
701 *National Academy of Sciences*, **112**, 1001-1006.

702 Tedesco, M., Willis, I., Hoffman, M., Banwell, A., Alexander, P. & Arnold, N. (2013). Ice dynamic
703 response to two modes of surface lake drainage on the Greenland ice sheet. *Environmental*
704 *Research Letters* **8**, 034007.

Formatted: Font: Italic

Formatted: Font: Bold

705 Tedesco, M., Fettweis, X., Mote, T., Wahr, J., Alexander, P., Box, J. E. & Wouters, B. (2013).
706 Evidence and analysis of 2012 Greenland records from spaceborne observations, a regional
707 climate model and reanalysis data. *The Cryosphere*, **7**, 615-630.

708 Tedesco, M., Fettweis, X., van den Broeke, M. R., van de Wal, R. S. W., Smeets, C. J. P. P., van de
709 Berg, W. J., Serreze, M. C. & Box, J. E. (2011). The role of albedo and accumulation in the
710 2010 melting record in Greenland. *Environmental Research Letters*, **6**, 1-6.

711 van As, D., Hubbard, A. L., Hasholt, B., Mikkelsen, A. B., van den Broeke, M. R. & Fausto, R. S.
712 (2012). Large surface meltwater discharge from the Kangerlussuaq sector of the Greenland
713 ice sheet during the record-warm year 2010 explained by detailed energy balance
714 observations. *The Cryosphere*, **6**, 199-209.

715 van de Wal, R. S. W., Boot, W., Smeets, C. J. P. P., Snellen, H., van den Broeke, M. R. &
716 Oerlemans, J. (2012). Twenty-one years of mass balance observations along the K-transect,
717 West Greenland. *Earth System Science Data*, **4**, 31-35.

718 van de Wal, R. S. W., Greuell, W., van den Broeke, M. R., Reijmer, C. H. & Oerlemans, J. (2005).
719 Surface mass-balance observations and automatic weather station data along a transect near
720 Kangerlussuaq, West Greenland. *Annals of Glaciology*, **42**, 311-316.

721 van de Wal, R. S. W., Smeets, C. J. P. P., Boot, W., Stoffelen, M., van Kampen, R., Doyle, S. H.,
722 Wilhelms, F., van den Broeke, M. R., Reijmer, C. H., Oerlemans, J. & Hubbard, A. (2015).
723 Self-regulation of ice flow varies across the ablation area in south-west Greenland. *The*
724 *Cryosphere*, **9**, 603-611.

725 van den Broeke, M., Smeets, P., Ettema, J. & Munneke, P. K. (2008). Surface radiation balance in
726 the ablation zone of the west Greenland ice sheet. *Journal of Geophysical Research*, **113**,
727 D13105.

728 Wientjes, I. G. M., De Van Wal, R. S. W., Schwikowski, M., Zapf, A., Fahrmi, S. & Wacker, L.
729 (2012). Carbonaceous particles reveal that Late Holocene dust causes the dark region in the
730 western ablation zone of the Greenland ice sheet. *Journal of Glaciology*, **58**, 787-794.

731 Wientjes, I. G. M. & Oerlemans, J. (2010). An explanation for the dark region in the western melt
732 zone of the Greenland ice sheet. *The Cryosphere*, **4**, 261-268.

733

734

735 **Table 1:** Energy inputs in 2010 and 2012 (TW).

Energy inputs— 0 to 1850 m.a.sl.	2010	2012	2012 _{to} -2010
Energy available for melt	2.43 x 10 ⁶	2.37 x 10 ⁶	—3-%

736

737


738

739 **Table 2:** Melt contributions (km³) from different elevation intervals integrated through end of melt
740 season, 1 Oct each year

	2010	2012	Difference
	km ³	km ³	%
Below mean ELA	6.8	6.3	—7
1550 to 1850 m	0.4	0.7	97
1850 to 2050 m	0.1	0.3	232
Total — up to 1850 m	7.1	7.0	—2
Total — up to 2050 m	7.2	7.3	1
% melt above mean ELA (1550 to 1850 m)	5.6 (%)	10.0 (%)	78
Measured proglacial discharge at Oct. 1	5.3	6.8	28
Integrated melt up 1850 m – measured discharge	1.8	0.2	—90
Integrated melt up 2050 m – measured discharge	1.9	0.5	—74

741

742 **Figures**



743


744 **Figure 1:** (A) The location of the study area (cyan) and catchment (red) in Greenland is shown on
 745 the inset map. (B) Map of the study area overlain with the location of the AWS, gauging station,
 746 catchment area, and snow pit sites. The background Landsat 7 image, which was acquired on 16
 747 July 2012, reveals ~~surface water in that~~ supraperiglacial lakes and streams ~~occurred-formed~~
 748 at an exceptional and unprecedented elevation of ~~~180~~ ~1800 m asl. The non-linear increase in the size of the
 749 catchment with increasing elevation is shown in (C) and (D) shows an example of the impact on
 750 melt area with a rise in the snow line of 250 m with a 500 m displacement in different start
 751 elevations (hypsometric effect).

752



753


754 **Figure 2:** Photograph taken at 18:00 West Greenland Summer Time on 11 July 2012 during the
755 flood with the Watson River [B](#)ridge being washed-out. Courtesy of Jens Christiansson.



757 **Figure 3:** Meteorological records, discharge measurements and modelled melt runoff for the study
 758 area during 2010 and 2012, including (A) daily average air temperature at AWS_L and AWS_U.
 759 (To avoid cluttering temperatures below $-10\text{ }^{\circ}\text{C}$ is not shown. Likewise the plot the air
 760 temperatures at AWS_U is only plotted during summer and the air temperature at AWS_M, which
 761 usually lies between that of AWS_L and AWS_U is not plotted at all). (B) the calculated cumulative
 762 energy input, (C) the albedo at three different elevation bands, (D, E) the proglacial discharge,
 763 supraglacial lake drainage volume, and modelled melt runoff, and (F) the cumulative proglacial
 764 discharge, modelled melt runoff, and residual between the two. The dashed vertical purple line
 765 demarks the bridge wash out on 11 July 2012. The uncertainty in discharge estimates is shown
 766 using grey lines on (d) and (e) and by grey shading on (f). Where the uncertainty estimates for 2010
 767 and 2012 overlap on (f) a darker shade of grey is used.


Commented [KL16]: ?

768
769
770



771
772 **Figure 4:** The cumulative measured discharge as a function of the calculated energy input for the
 773 catchment up to 1850 m a.s.l. The flooding period of 11 to 14 July is marked with a bold red line.


774



775
776 **Figure 5:** (A-C) Density profiles of three shallow firn cores (A-C respectively) drilled at AWS_U
 777 in May-April 2012. The water table is indicated in light blue and ice lenses observed in the core

778 straigraphy are indicated in cyan. Magenta and red lines indicate two potential sets of "blocking" ice
779 lenses observed in the firn. (D) A model simulation of the near-surface water table at AWS_U for
780 each of the two blocking lens assumptions in A-C, with 95% confidence intervals in grey. Red ticks
781 on the horizontal axes indicate days above freezing when surface melt would occur. As snow melts
782 above the blocking lenses the water table rises simultaneously until it meets the lowering snow
783 surface. Light blue is free air. The daily snow surface is observed by the adjacent AWS_U AWS.
784 The two dashed orange vertical lines indicate 11 July, the date of the Watson River bridge
785 destruction, and 16 July, when the Landsat image from Figure 1 shows horizontal water transport in
786 the vicinity of AWS_U.

787



788

789 **Figure 6:** (A) Zoom in on Landsat 7 image from 16 July 2012 showing free surface water in the
790 area around AWS_U. The extent is marked on Figure 1. The scan line correction failure was
791 interpolated using the ENVI 'replace bad data' routine based on Band 8 and visible surface water
792 was enhanced using a modified normalized difference water index (Fitzpatrick et al., 2014). (B and
793 C) Oblique aerial photographs of the active supraglacial channel network emerging from AWS_U


794 well within the accumulation zone at 1840 m a.s.l. and 140 km from the ice sheet margin on 13
795 August 2012. Courtesy of Paul Smeets.

796

797

798

799



800 **Supplementary Figures**

801 Energy balance for the three weather stations AWS_L, AWS_M and AWS_U (located at 680, 1270
802 and 1840 m a.s.l. respectively), as based on the surface energy balance model explained in section
803 2.5. The energy balance is shown as the yearly averaged energy fluxes for the respective 100 m
804 elevation interval corresponding to the weather stations at each elevation interval. The componets
805 shown are SSH = sub-surface heat flow, LHF = latent heat flow, LRnet = net long wave radiation,
806 SHF = sensible heat flow, SRnet = net short-wave radiation. M = energy available for melt. Energy
807 input from rain is omitted on the figure given it is contributing with a maximum of 0.1 W/m^2 when
808 averaged over a year. When the number is positive, the energy flux is directed towards the surface
809 and vice versa when it is negative.

810

811 For AWS_L, the main difference between year 2010 and 2012 is a 10.2 W/m^2 smaller SRnet
812 influx of energy over the year. The energy input from SHF is 2.6 W/m^2 smaller for the averaged
813 year and the loss of energy through LRnet is 3.9 W/m^2 samller in 2012 compared to 2010. The
814 removal of energy through LHF is 1.4 W/m^2 smaller in 2012 compared to 2010, where SSH is 1.24
815 W/m^2 larger in 2012. Overall the resulting energy available for melt is 6.5 W/m^2 smaller for the
816 KAN_L elevation in 2012, as compared to 2010.

817

818 For AWS_M, the energy input for SRnet and SHF was 4.7 and 2.4 W/m^2 smaller respectively in
819 2012 compared to 2010. The removal of energy via LRnet and LHF was respectively 5 and 1.4 W
820 m^2 samller in 2012 compared to 2010. SSH represented a positive flux towards the surface, that
821 was 0.4 W/m^2 larger in 2012. The resulting energy available for melt is almost equal between the
822 two years with a 0.6 W/m^2 larger energy input in 2010.


823

824 For AWS_U, the energy input for SRnet was 0.6 larger in 2012 relative to 2010, where SHF was
825 3.7 W/m^2 smaller in 2012 compared to 2010. The removal of energy via LRnet and LHF was
826 respectively 5.1 and 0.8 W/m^2 samller in 2012 compared to 2010. SSH represented a positive flux
827 towards the surface, that was 1.5 W/m^2 larger in 2012. The resulting averaged energy available for
828 melt in 2012 was 4.3 W/m^2 larger than in 2010.


Formatted: Superscript

Formatted: Highlight


829



830



831



832

833

834