

The Cryosphere Discuss., 2, 843–872, 2008
www.the-cryosphere-discuss.net/2/843/2008/
© Author(s) 2008. This work is distributed under
the Creative Commons Attribution 3.0 License.



The Cryosphere Discussions is the access reviewed discussion forum of *The Cryosphere*

A new 1 km digital elevation model of Antarctica derived from combined radar and laser data – Part 2: Validation and error estimates

J. A. Griggs and J. L. Bamber

Bristol Glaciology Centre, School of Geographical Sciences, University of Bristol, Bristol, UK

Received: 18 September 2008 – Accepted: 26 September 2008
– Published: 25 November 2008

Correspondence to: J. A. Griggs (j.griggs@bristol.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

TCD

2, 843–872, 2008

**A new Antarctic
digital elevation
model: validation and
errors**

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We have developed a new digital elevation model (DEM) of Antarctica from a combination of satellite radar and laser altimeter data. Here, we assess the accuracy of the DEM by comparison with airborne altimeter data from four campaigns covering a wide range of surface slopes and ice sheet regions. RMS differences varied from 4.84 m, when compared to a densely gridded airborne dataset over the Siple Coast region of West Antarctica to 29.28 m when compared to a more limited dataset over the Antarctic Peninsula where surface slopes are high and the across track spacing of the satellite data is relatively large. The airborne data sets were employed to produce an error map for the DEM by developing a multiple linear regression model based on the variables known to influence errors in the DEM. Errors were found to correlate highly with surface slope, roughness and density of satellite data points. Errors ranged from typically ~ 1 m over the ice shelves to between about 4 and 10 m for the majority of the grounded ice sheet. In the steeply sloping margins, along the Peninsula and mountain ranges the estimated error is several tens of metres. Slightly less than 7% of the area covered by the satellite data had an estimated random error greater than 20 m.

1 Introduction

Digital elevation models (DEMs) of the Antarctic are important datasets for a wide range of applications from fieldwork planning, calculating drainage basins, determining balance velocities to dynamical modelling of the ice sheet. The limitations of currently available DEMs (Bamber and Gomez-Dans, 2005) and the availability of new, higher quality altimeter data for the Antarctic continent has lead us to produce a new DEM. The methodology and data used is described in detail in a companion paper (Bamber et al., 2008) and so we repeat only the most salient points here. The new DEM was created by combining high-accuracy, but relatively low spatial resolution, laser altimeter data from the GLAS instrument onboard IceSat (Zwally et al., 2007) with radar altimeter data

TCD

2, 843–872, 2008

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

from the geodetic phase of the ERS-1 satellite, which provided high spatial resolution but lower vertical accuracy particularly in highly sloping areas (Brenner et al., 2007). The ERS-1 data are the same as those used to create an earlier DEM derived solely from radar altimetry (Bamber and Bindschadler, 1997). The resulting DEM was gridded with 1 km postings.

For most applications of an ice sheet DEM, it is important to i) minimise the errors and ii) to determine what the errors are, how they vary spatially and as a function of variables such as surface slope and roughness. Previously published DEMs included some limited validation and error estimation but this was, in general, confined to small areas such as the two traverses used to validate the RAMP DEM (Liu et al., 1999) or to comparison with visible imagery (Bamber and Bindschadler, 1997). A recent unpublished IceSat DEM of Greenland (DiMarzio et al., 2007) was validated using airborne data but a similar analysis was not performed on the equivalent Antarctic DEM. Thus, no error estimates are available for this product. Here, we assess the accuracy of our new DEM using a range of extensive airborne altimeter data sets covering both East and West Antarctica, interior plateau regions, marginal areas and ice shelves. The results of this comparison were used to calculate an error map for the entire continent.

2 Validation

As mentioned, the accuracy of the new DEM was estimated by comparison with airborne altimeter data. The airborne data had a variable accuracy in the range 0.08–1.91 m depending on the campaign. It should be noted that the airborne data measured the surface elevation at a higher spatial resolution than the satellite sensors and so some differences will be due to the airborne data ranging into crevasses and rifts below the surface which were not resolved by the spaceborne instruments. In addition, temporal differences between the time stamp of the DEM and the airborne data must also be considered particularly in areas of fast-flow and known surface elevation change such as Pine Island and Thwaites glaciers (Shepherd et al., 2002) and Kamb

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice Stream.

Four airborne datasets were compared to the DEM to assess its accuracy. These datasets were

- CECS/NASA over the Antarctic Peninsula and Pine Island, Thwaites, Pope, Smith and Kohler glaciers
- AGASEA over the Amundsen Sea sector
- SOAR CASERTZ over the south-eastern Ross Embayment and
- ISODYN/WISE in northern Victoria Land.

The locations of these data are shown in Fig. 1 and the acronyms explained below. The comparison with the DEM is discussed for each data set, next. In all cases, bilinear interpolation was used to calculate the DEM elevation at the exact location of the airborne measurement.

2.1 CECS/NASA

The CECS/NASA data were collected on a joint Centro de Estudios Científicos (CECS), Chile and NASA organised survey (Rignot et al., 2004). The areas covered were Pine Island, Thwaites, Pope, Smith and Kohler glaciers in the Amundsen Sea sector and the Antarctic Peninsula. The survey operated from a Chilean P-3 aircraft carrying the NASA/Wallops Airborne Topographic Mapper (ATM) (Krabill et al., 2000). Flights of 750–1000 km length were collected in November and December 2002. The survey operated with a long baseline GPS (~1400 km) which gave an accuracy of around 20–30 cm, a factor two lower than achieved in Greenland where shorter baselines were used.

Figure 2 shows the difference between the CECS/NASA data for the Antarctic Peninsula. The differences are overlaid on a shaded relief plot of the DEM. Over the Larsen Ice Shelf, where both datasets would be expected to be most accurate, the agreement

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is good with a mean difference of 46 cm and a standard deviation of those differences of 3.22 m. On the highly sloping areas of the Peninsula, the agreement is poorer with mean differences of $26.20\text{ m} \pm 40.63\text{ m}$ (one standard deviation). This is an area of the DEM with little satellite data and interpolation lengths are long. Over the entire Peninsula region, the difference between the airborne data and the DEM has a mean bias of -1.51 m , modal bias of 16.64 m and RMS difference of 29.28 m.

Figure 3 shows the difference between the ICE/CECS data and the DEM for the Amundsen Sea sector. Here, better agreement is seen than over the Peninsula due to the greater amount of satellite data present in the DEM and the more gentle relief. Over Pine Island glacier and ice shelf, the agreement is better than over Thwaites glacier with a mean difference of $-2.28\text{ m} \pm 11.80\text{ m}$ compared to a mean difference of $-7.37\text{ m} \pm 21.15\text{ m}$ for Thwaites. A small section over the centre of Thwaites glacier (indicated by the red box on Fig. 3) shows differences of consistently around 70 m with the DEM being higher than the airborne data. This is a region, which is believed to be close to, or oceanward of, the 1996 limit of tidal flexure and so is likely on the floating ice tongue of Thwaites glacier (Rignot et al., 2004). The ice tongue surface does not resemble a simple ice shelf but is a collection of broken up icebergs attached together by an ice melange of sea ice, ice sheet debris and windblown snow (Rignot and MacAyeal, 1998). The ice tongue rifts and calves into large tabular icebergs along a significant proportion of the grounding line (Rignot, 2001). It is therefore likely the airborne data is ranging, in places, to the low elevation melange and the satellite data to the top of the icebergs. 70 m is around the likely freeboard of icebergs calved directly from Thwaites glacier. As the ice tongue is a rapidly changing ice feature, it would not be expected that the DEM would capture the surface well as the DEM is created from data recorded over a period of several years. Overall, the agreement between the DEM and CECS/NASA data in the Amundsen Sea sector is good with the histogram of differences showing a mean bias of -3.58 m and a modal bias of 0.55 m, skewed towards negative differences and a RMS difference of 15.94 m.

A new Antarctic digital elevation model: validation and errorsJ. A. Griggs and J. L. Bamber

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2.2 AGASEA

The AGASEA (Airborne Geophysical Survey of the Amundsen Sea Embayment, Antarctica) dataset (Young et al., 2008) was collected by the University of Texas Institute for Geophysics between 12 December 2004 and 30 January 2005. The data were collected with a 15 km grid spacing over the Thwaites and Smith glacier catchments using a Twin Otter fitted with an ice penetrating radar, gravimeter and magnetometer in addition to a nadir pointing laser distance meter for measuring surface elevation. In addition to flying the grid, targeted profiles were flown re-flying three IceSat ground tracks. The laser altimeter configuration results in a 1 m wide ground spot size with an along track resolution of approximately 20 m. Cross-over analysis of the data showed an RMS difference for the entire dataset of ± 20 cm and ± 8 cm when a reference dataset consisting of datapoints closest to the base reference GPS station was considered. The data exhibited an approximately 20 cm bias from the IceSat dataset in flat areas for which the source is unknown (Young et al., 2008). This bias increased to 27 cm when all areas were considered.

Figure 4 shows the difference between the AGASEA airborne data and DEM along with a histogram of the differences. The agreement between the airborne data and the DEM is good in flat and low slope areas such as the area marked by the red box where the mean difference is $11 \text{ cm} \pm 3.70 \text{ m}$. The agreement is poorer in fast flow areas with the coastal region of the dataset having a mean difference of $-9.07 \text{ m} \pm 21.25 \text{ m}$. This area is where crevassing is more prevalent and so agreement is expected to be poorer due to the difference in spatial resolution of the two types of measurement. This is manifested in differences which are highly variable exhibiting both positive and negative differences close to each other. The DEM can be seen to not fully resolve nunataks due to the small spatial scale compared to the interpolation length. The mean bias observed across the whole dataset is -2.9 m with a modal bias of 0.7 m . The AGASEA dataset shows good agreement with the DEM with an RMS difference of 11.62 m .

TCD

2, 843–872, 2008

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.3 SOAR CASERTZ

The SOAR CASERTZ data (Support Office for Aerogeophysical Research – Corridor Aerogeophysics of the South East Ross Transect Zone) data were recorded by the University of Texas Institute for Geophysics and the US Geological Survey (Blankenship et al., 2001). Flights were conducted over an area encompassing the onset area and catchment region of Whillans and Kamb Ice Streams (formerly, Ice Streams B and C), all of Bindschadler Ice Stream (Ice Stream D) and the boundary between the West Antarctic rift system and the crustal provinces dominated by Byrd Subglacial Trench and the Whitmore Mountains. Data were recorded in Antarctic summer seasons between 1991 and 1997 using a laser altimeter on a Twin Otter aircraft providing approximately 8 m along track spacing. The instrument was capable of a precision of better than 10 cm in the absence of cloud cover and surface elevation profiles were repeatable to within 25 cm across track. RMS deviations were calculated at cross-over points and values varying between ± 37 cm for the earliest seasons, before the full GPS constellation was deployed, to ± 9 cm for the most recent survey. The data were provided on a 425 m resolution grid.

The agreement between the airborne data and the DEM (Fig. 5) is slightly better in the region north of 81.5° S where data from both the ERS and IceSat instruments were included in the DEM. As in the comparison with the AGASEA data, the underlying surface properties can explain the differences observed. In the region of Raymond Ice Rise where flow is slow and the surface is uncrevassed, the agreement is very good with a mean differences between the airborne data and the DEM of $28.7 \text{ cm} \pm 2.43 \text{ m}$ whereas in Bindschadler Ice Stream (D) where flow is on the order of 100's m/yr and surface crevassing is common, agreement is worse with a mean difference of $-0.47 \text{ m} \pm 6.07 \text{ m}$. Positive and negative differences indicate that the airborne data were penetrating into crevasses whereas satellite data used to produce the DEM are unlikely to adequately resolve such small-scale. Further south in the region of the, now stagnant, Kamb Ice Stream (C) a bias of -4.80 m is seen. Kamb Ice Stream

TCD

2, 843–872, 2008

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

stopped flowing about 150 years ago and has been shown to be thickening significantly (Csatho et al., 2005). The size of negative differences between the SOAR CASERTZ data and the DEM are similar to that expected from thickening of up to 0.8 m/yr over the years between the 1991–1993 time stamp of the SOAR CASERTZ data in this region and the 2004 time stamp of the DEM. The mean bias between the airborne data and the DEM is 0.2 m with a modal difference of 1.94 m and an RMS difference of 4.84 m.

The agreement between the DEM and airborne data on Siple Dome supports the use of the new surface roughness based bias correction applied to the ERS data (Bamber et al., 2008). This was one of the regions where applying a bias correction calculated from slope as a proxy for surface roughness was ineffective, as the surface of Siple Dome is highly sloping yet surface roughness is low (i.e. the second derivative of height is small). A surface slope bias correction was applied in earlier versions of the new DEM (Bamber and Gomez-Dans, 2005). Figure 6 shows the difference between the SOAR CASERTZ data and both the DEM and an earlier version which uses a surface slope based bias correction. The improved agreement in this region justifies the more physically realistic bias correction to the ERS data.

2.4 ISODYN/WISE

The ISODYN/WISE data (Ferraccioli et al., 2007) were collected as part of a joint Italian – British Antarctic Survey campaign in the northern Victoria Land. Data were collected using a radar altimeter (Vaughan et al., 2006) on 68 flights between November 2005 and January 2006. The spatial sampling of the data was approximately 22 m. A cross-over analysis showed a standard deviation of 1.91 m for all cross-overs and 1.29 m when considering only those cross-overs on the flatter plateau area of the campaign region. Figure 7 shows the difference between the ISODYN/WISE data and the DEM. Agreement is better on the inland flatter areas as would be expected. A mean bias of 1.05 m, a modal bias of 2.61 m and RMS difference of 11.54 m was obtained for the ISODYN/WISE data and the DEM.

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3 Discussion

The results of the comparison of the DEM with the 5 independent airborne altimetry datasets is shown in Table 1. The accuracy and random error in the new DEM is good in most areas, worsening in areas of steeper relief and poorer satellite coverage. To put these results in context, we have compared them with other DEMs of Antarctica. The AGASEA dataset have been compared with the RAMP (Liu et al., 1999), ERS-1 (Bamber and Bindschadler, 1997) and IceSat (DiMarzio et al., 2007) DEMs in a previous study (Young et al., 2008). A bicubic interpolation scheme was used to interpolate the DEMs to the location of the airborne data. This, however, makes a negligible difference to the statistics of the differences when compared with the bilinear interpolation used here. The results from the earlier study along with those from the comparison of our new DEM are given in Table 2, for the AGASEA data set. Values are shown for both the entire area as well as for a central, plateau area, highlighted by the red box on Fig. 4. The RMS difference for the whole area is lowest for the new DEM and around 10% higher than the ICESat DEM for the plateau region.

As the AGASEA comparison is limited in spatial extent and sampling density, we have also compared the older DEMs to the SOAR CASERTZ dataset which covers an extensive area of grounded and floating ice with dense sampling and decimetre level accuracy (Blankenship et al., 2001). In addition it covers regions of the new DEM which include just GLAS data as well as areas containing both GLAS and ERS-1 data. Table 3 shows the results of this comparison. In the case of the ERS-1 DEM, we only make a comparison in area south of 81.5° as, south of this latitude, the DEM was created using cartographic data with accuracy considerably poorer than elsewhere (Bamber and Gomez-Dans, 2005). Not surprisingly, the new DEM has a lower RMS and standard deviation compared with the ERS-1 DEM but also compared to the ICESat DEM, although the differences here are on the order of 20 cm, equivalent to $\sim 4\%$. However, some small scale features on the surface are still not fully captured as seen by the pattern of alternating positive and negative differences in Fig. 5. A profile of the dif-

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ferences between the SOAR CASERTZ data and the DEMs, along the track shown in blue on Fig. 5, is plotted in Fig. 8. All the DEMs are missing small scale features at the scale of ~ 5 km. The reason for this in the new DEM may be due to the interpolation methodology used or possibly the movement of some small scale features in the interval between the satellite and terrestrial observations..

4 Error map

The validation shows the quality of the DEM in the selected areas where airborne data exists. We use the results of these comparisons to derive an error map for the new DEM. Our implicit assumption is that differences are predominantly due to errors in the DEM. In other words, i) we neglect errors in the airborne data, ii) we ignore the effect of differences in spatial resolution and iii) assume that there are sufficient airborne validation data to fully characterise the errors in the DEM. The errors in the airborne data have been found from crossover analysis to be of the decimetre scale which justifies our first assumption. In the case of the second assumption, we consider a single profile of airborne data and compare that to the DEM. We find no differences at wavelengths less than 2 km, above the noise level of the airborne instrument, which would create a different result had the airborne data had a 1 km footprint. Our third assumption is justified later. The consequence of these assumptions is that our map is likely to slightly over estimate the random errors in the DEM by assigning error from all sources to the DEM.

Multiple regression can be used to predict a spatially complete error estimate by defining the relationship between a set of spatially complete parameters and the spatially incomplete measured errors between the DEM and airborne data. A multiple regression model states that the mean of the required variable, Y , in this case the RMS error in the DEM, can be expressed as a linear combination of the k dependent variables, $X_l (l=1, 2, \dots, k)$ for all n points in our airborne study regions. The multiple

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

regression model is expressed as

$$Y_i = a_0 + \sum_{l=1}^k a_l X_{li} + E_i \quad (1)$$

where E_i for $i=1, \dots, n$ are zero mean, independent identically distributed, random errors in the multiple regression and a_0, a_1, \dots, a_k are unknown model parameters. The validation data could be used to ascertain the values of a_0, a_1, \dots, a_k as long as the validation data covered the full range of values of the dependent variables. The model could then be applied to the whole of Antarctica to create a map of RMS error. The airborne datasets used in the model were the full AGASEA and SOAR CASERTZ datasets and the ISODYN/WISE data from elevations over 2200 m. The data were chosen to include the best quality airborne data over the widest range of surface slopes and type. The modelled parameter was RMS error. The RMS error was calculated from all the differences between the airborne data and the DEM in each DEM gridbox containing airborne data. 98.5% of RMS errors were calculated from 4 or more differences. To determine the range of dependent variables, X_l used in the model, a backwards elimination technique was used. In this method, a multiple regression model was created using all possible dependent variables. The dependent variable which reduces the regression sum of squares by the smallest amount was then determined and a second model created excluding this variable. An F-test was conducted to determine whether the model with all variables was significantly different from the reduced model. If the excluded variable did not contribute significant variance to the model, as determined by the F-test, it was removed from the full model. The process was repeated until only dependent parameters which contribute significant variance were included in the full model. The full set of dependent variables were

1. Surface slope (X_1). ERS data contained a surface slope bias which was removed but random error in the ERS data has been shown to increase as a function of increasing surface slope. The pointing accuracy of IceSat is 1 arcsec (Schutz

TCD

2, 843–872, 2008

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A new Antarctic
digital elevation
model: validation and
errors**

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2005) which will give an increasing large error in elevation for increasing surface slope with a 2° slope having an error of 10 cm.

2. Surface roughness (X_2). ERS data contains a surface roughness bias as they range to the tops of undulations in the surface. The mean bias was removed but a random component may remain.
3. Number of datapoints (X_3). Increasing numbers of satellite datapoints within each DEM gridbox decreases the random error on the measurement of the elevation in that gridbox.
4. Standard deviation (X_4). The standard deviation of all satellite datapoints within each DEM gridbox gives an approximate measure of the error and variance in elevation within a grid cell.
5. Deviation from the quasi-regular grid (X_5). All satellite datapoints were averaged to a quasi-regular grid before being interpolated to the regular 1 km grid of the final DEM. The difference between the quasi-regular grid and the final DEM gives a measure of the error introduced by the use of an interpolation method which produces a smooth surface by allowing the interpolation surface to deviate from the input data.
6. Distance to real datapoint (X_6). The further any DEM gridbox was from one which contains satellite data then the larger the effect of the interpolation of the satellite data will be and the greater the chance of missing true changes in the surface elevation.

Figure 9 shows plots of the dependent variables against the RMS error calculated from the difference between the airborne data and the DEM. All dependent parameters were binned into 200 equally sized bins over the full range of values shown. The RMS error is correlated with all dependent variables. At high values of all dependent variables, scatter increases, however, few datapoints exist and this contributes

to uncertainty in the regression model as expressed by E_j . The full set of dependent parameters produce a regression model of the form

$$Y = 1.023 + 14.286X_1 + 10.887X_2 - 0.025X_3 + 0.206X_4 + 0.121X_5 + 0.861X_6 \quad (2)$$

The model was tested to ensure that a regression relation existed by testing the null hypothesis that all regression coefficients were equal to zero. The relation was significant at less than the 0.1% level. The dependent parameter which contributed the least variance to the model was X_3 , the number of satellite datapoints within each DEM gridbox. This was also the parameter which had the smallest error associated with it. However, the parameter contributed significant variance to the model at the 99% confidence level and so no dependent variable could be eliminated from the model and Eq. (2) is the full regression model for the RMS error in the DEM.

The ability of the regression model to fully characterise the error in the DEM can be determined by the use of a bootstrap technique. We create 10 000 regression models which each contain a random 95% of the available airborne data. These models can then be used to calculate the 95% confidence levels on the regression model. We find that the 95% confidence levels on our regression model are in general less than 0.5 m, increasing over mountainous regions, but remaining much less than 25% of the size of the error in all areas apart from those with the very lowest value of RMS error. This justifies our third assumption in using this method to create an error estimate for our DEM.

The regression model was applied to all DEM gridboxes north of 86° S. South of this limit, there were no satellite data and the DEM was filled with cartographic data as described in Bamber et al. (2008). These data do not have the same properties as the satellite derived areas. In the areas of cartographic data, a value of the RMS error was calculated as the RMS difference between the DEM and cartographic data from the same source in a latitude band between 81.5° S and 86° S. The error in the area south of 86° was found to be 77.09 m. We also limit the maximum size of the error from the regression model to this value. This threshold is applied in some mountain-

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ous regions, coastal areas with large interpolation lengths and in some parts of the Peninsula, accounting for less than 1% of all gridboxes.

Figure 10 shows the calculated value of RMS error across the whole Antarctic continent. The figure has a range of 0 to 20 m but larger values of exist in coastal areas, particularly in the Peninsula, where little satellite data exist, and the amount of interpolation present in the DEM was large. The pattern of RMS error calculated from the regression model is as would be expected with errors of around 1–2 m on the ice shelves, where the instruments are known to perform well and spatial sampling is almost complete. The error increases slightly on the Ross and Filchner-Ronne ice shelves south of 81.5° S, moving from an area close to the orbital limit of ERS with very dense satellite coverage to an area with only GLAS data and poorer coverage.

5 Conclusions

In Part 1, a new DEM of Antarctica was presented. Airborne validation of this new DEM showed low biases and random errors, roughly halving the errors compared to the DEM without the benefit of the GLAS data (Bamber and Gomez-Dans, 2005). We have shown how the errors are a function of the underlying properties of the surface and satellite datasets. The highest errors were obtained in the Peninsula, where the terrain is rugged and satellite data limited. The lowest errors, unsurprisingly, were on the floating ice shelves.

Using a multiple regression model, we have created, what we believe to be, the first rigorous error map to accompany a large scale DEM of Antarctica. Multiple regression modelling showed that all six parameters considered have a significant effect on the RMS error of the DEM and all were, therefore, included in the model. These parameters were surface slope, surface roughness, number of satellite datapoint in gridbox, standard deviation of those datapoints, deviation of the interpolated surface from the mean of the satellite datapoints and the distance of an interpolated gridbox to the closest satellite data. The error map shows the expected pattern of low error on smooth,

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

flat surfaces such as the ice shelves and in areas over sub-glacial lakes, with increasing error with increasing slope and surface roughness and in regions with less satellite data. The error map will allow users of the DEM to assess its accuracy and suitability for their given application.

5 *Acknowledgements.* The authors would like to thanks the following data contributors: W. Krabill from NASA Goddard Wallops Flight Facility for the CECS/NASA data, Duncan Young from the University of Texas for the AGASEA data and Hugh Corr from the British Antarctic Survey for the ISODYN/WISE data. The SOAR data are based on work supported by the National Science Foundation under Grants: OPP-9120464, 9319369 and 9319379 and we'd like to thank SOAR
10 and the University of Texas for allowing us access. This work was funded by NERC grant NE/E004032/1.

References

- Bamber, J. and Gomez-Dans, J. L.: The accuracy of digital elevation models of the Antarctic continent, *Earth Planet. Sc. Lett.*, 237, 516–523, 2005.
- 15 Bamber, J. L. and Bindschadler, R. A.: An improved elevation dataset for climate and ice-sheet modelling: validation with satellite imagery, *Ann. Glaciol.*, 25, 439–444, 1997.
- Bamber, J. L., Gomez-Dans, J. L., and Griggs, J. A.: A new 1 km digital elevation model of Antarctica derived from combined radar and laser data – Part 1: Data and methods, *The Cryosphere*, 2, 811–841, 2008.
- 20 Blankenship, D. D., Morse, D., Finn, C. A., Bell, R. A., Peters, M. E., Kempf, S. D., Hodge, S. M., Studinger, M., Behrendt, J. C., and Brozena, J. M.: Geologic controls on the initiation of rapid basal motion for West Antarctic ice streams: A geophysical perspective including new airborne radar sounding and laser altimetry results, in: *The West Antarctic Ice Sheet, Behavior and Environment*, edited by: Alley, R. B. and Bindschadler, R. A., American Geophysical Union, 105–121, 2001.
- 25 Brenner, A. C., DiMarzio, J. R., and Zwally, H. J.: Precision and accuracy of satellite radar and laser altimeter data over the continental ice sheets, *IEEE T. Geosci. Remote*, 45, 321–331, doi:310.1109/TGRS.2006.887172, 2007.
- Csatho, B., Ahn, Y., Yoon, T., van der Veen, C. J., Vogel, S., Hamilton, G., Morse, D.,

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A new Antarctic digital elevation model: validation and errorsJ. A. Griggs and J. L. Bamber

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Smith, B., and Spikes, V. B.: ICESat measurements reveal complex pattern of elevation changes on Siple Coast ice streams, Antarctica, *Geophys. Res. Lett.*, 32, L23S04, doi:10.1029/2005GL024289, 2005.

DiMarzio, J. P., Brenner, A. C., Schutz, B. E., Shuman, C. A., and Zwally, H. J.: GLAS/ICESat 500 m laser altimetry digital elevation model of Antarctica. Boulder, Colorado, USA, National Snow and Ice Data Center, Digital Media, 2007.

Ferraccioli, F., Jordan T., Armadillo E., Bozzo E, Corr H., Caneva G., Robinson C., Frearson N., and Tabacco I.: Collaborative Aerogeophysical Campaign targets the Wilkes Subglacial Basin, the Transantarctic Mountains and the Dome C region, in: The Italian-British Antarctic Geophysical and Geological Survey in Northern Victoria Land 2005–2006 – towards the International Polar Year 2007–2008, edited by: Bozzo, E. and Ferraccioli, F., *Terra Antarctica Reports*, 13, 1–36, 2007

Krabill, W., Abdalati, W., Frederick, E., Manizade, S., Martin, C., Sonntag, J., Swift, R., Thomas, R., Wright, W., and Yungel, J.: Greenland ice sheet: High-elevation balance and peripheral thinning, *Science*, 289, 428–430, 2000.

Liu, H. X., Jezek, K. C., and Li, B. Y.: Development of an Antarctic digital elevation model by integrating cartographic and remotely sensed data: A geographic information system based approach, *J. Geophys. Res.-Sol. Ea.*, 104, 23 199–23 213, 1999.

Rignot, E. and MacAyeal, D. R.: Ice-shelf dynamics near the front of the Filchner-Ronne Ice Shelf, Antarctica, revealed by SAR interferometry, *J. Glaciol.*, 44, 405–418, 1998.

Rignot, E.: Evidence for rapid retreat and mass loss of Thwaites Glacier, West Antarctica, *J. Glaciol.*, 47, 213–222, 2001.

Rignot, E., Thomas, R. H., Kanagaratnam, P., Casassa, G., Frederick, E., Gogineni, S., Krabill, W., Rivera, A., Russell, R., Sonntag, J., Swift, R., and Yungel, J.: Improved estimation of the mass balance of glaciers draining into the Amundsen Sea sector of West Antarctica from the CECS/NASA 2002 campaign, *Ann. Glaciol.*, 39, 231–237, 2004.

Schutz, B. E., Zwally, H. J., Shuman, C. A., Hancock, D., and DiMarzio, J. P.: Overview of the ICESat Mission, *Geophys. Res. Lett.*, 32, L21S01, doi:10.1029/2005GL024009, 2005.

Shepherd, A., Wingham, D. J., and Mansley, J. A. D.: Inland thinning of the Amundsen Sea sector, West Antarctica, *Geophys. Res. Lett.*, 29, 1364, doi:10.1029/2001GL014183, 2002.

Vaughan, D. G., Corr, H. F. J., Ferraccioli, F., Frearson, N., O'Hare, A., Mach, D., Holt, J. W., Blankenship, D. D., Morse, D. L., and Young, D. A.: New boundary conditions for the West Antarctic ice sheet: Subglacial topography beneath Pine Island Glacier, *Geophys. Res. Lett.*,

33, L09501, doi:09510.01029/02005GL025588, 2006.

Young, D. A., Kempf, S. D., Blankenship, D. D., Holt, J. W., and Morse, D.: New airborne laser altimetry over the Thwaites Glacier catchment, West Antarctica, *Geochem. Geophys. Geos.*, 9, Q06006, doi:06010.01029/02007GC001935, 2008.

- 5 Zwally, H. J., Schutz, B. E., Bentley, C. R., Bufton, J., Herring, T., Minster, J. F., Spinhirne, J., and Thomas, R.: GLAS/ICESat L2 Antarctic and Greenland Ice Sheet Altimetry Data V428, 25th September 2003 to 27th November 2006, Boulder, CO: National Snow and Ice Data Center, Digital Media, 2007.

TCD

2, 843–872, 2008

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Table 1. Statistics of the comparisons between the DEM and each airborne dataset.

	Number of points	Mean bias (m)	Modal bias (m)	Standard deviation(m)	RMS difference(m)	FWHM (m)
CECS/NASA peninsula	107561	-1.51	16.64	29.24	29.28	4.0
CECS/NASA Amundsen	203123	-3.58	0.55	15.54	15.94	9.0
AGASEA	1681920	-2.9	0.7	11.24	11.62	6.4
AGASEA (central area)	138077	0.11	0.94	3.70	3.71	2.6
SOAR CASERTZ	1615531	0.2	1.94	4.84	4.84	4.5
ISODYN/WISE	2182625	1.05	2.61	11.49	11.54	3.8
ISODYN/WISE (elevations over 2200 m)	910854	3.03	2.85	6.18	6.89	3.8

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Table 2. Comparison of the new DEM with the results from older DEMs when compared to the AGASEA airborne datasets. Results from the older DEMs are all taken from Young et al. (2008).

	Bias whole area (m)	RMS whole area (m)	Bias central area (m)	RMS central area (m)
New DEM	0.72	13.95	1.26	4.89
ERS-1 DEM	-1.4	27.4	-0.46	5.2
RAMP DEM	0.4	55	1.07	5.7
IceSat DEM	-0.2	18.4	0.4	4.4

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Table 3. Comparison of the new DEM with the results from older DEMs when compared to the SOAR CASERTZ airborne datasets.

	Number of points	Mean bias (m)	Modal bias (m)	Standard deviation (m)	RMS difference (m)	FWHM (m)
New DEM	1615531	0.2	1.94	4.84	4.84	4.5
ERS-1 DEM	711784	−2.48	1.29	5.84	6.35	5.0
Icesat DEM	1615531	0.64	1.61	5.08	5.12	5.0

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. Validation datasets. Red dots indicate the position of the CECS/NASA data, blue dots indicate AGASEA data, green dots indicate SOAR CASERTZ data and yellow dots indicate the position of ISODYN/WISE data.

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**A new Antarctic
digital elevation
model: validation and
errors**

J. A. Griggs and J. L. Bamber




Fig. 2. Difference between the CECS/NASA data and the DEM in the peninsula region overlaid on a shaded relief of the DEM. Inset shows a histogram of the differences.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**A new Antarctic
digital elevation
model: validation and
errors**

J. A. Griggs and J. L. Bamber




Fig. 3. Difference between the CECS/NASA data and the DEM in the Amundsen Sea sector overlaid on a shaded relief of the DEM. Inset shows a histogram of the differences.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A new Antarctic digital elevation model: validation and errorsJ. A. Griggs and J. L. Bamber




Fig. 4. Difference between the AGASEA data and the DEM in the Amundsen Sea sector overlaid on a shaded relief of the DEM. Inset shows a histogram of the differences.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A new Antarctic digital elevation model: validation and errorsJ. A. Griggs and J. L. Bamber




Fig. 5. Difference between the SOAR CASERTZ data and the DEM in the Siple coast region overlaid on a shaded relief of the DEM. Inset shows a histogram of the differences.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber




Fig. 6. (a) Difference between the SOAR CASERTZ data and the DEM over Siple Dome. (b) Difference between the SOAR CASERTZ data and an earlier version of the DEM created using a slope based bias correction on the ERS data.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber




Fig. 7. Difference between the ISODYN/WISE data and the DEM in northern Victoria Land overlaid on a shaded relief of the DEM. Inset shows a histogram of the differences.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**A new Antarctic
digital elevation
model: validation and
errors**

J. A. Griggs and J. L. Bamber




Fig. 8. Difference between the SOAR CASERTZ dataset and the three DEM along a profile across the Siple Coast. Black shows the difference to the new DEM, red shows the difference to the IceSat DEM and blue shows the difference to the ERS-1 DEM.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A new Antarctic digital elevation model: validation and errorsJ. A. Griggs and J. L. Bamber




Fig. 9. Plots of the dependent variables in the multiple regression model against the RMS error in the DEM. **(a)** shows surface slope, **(b)** shows surface roughness, **(c)** shows the number of satellite datapoints, **(d)** shows the standard deviation of the satellite datapoints within each DEM gridbox, **(e)** shows the deviation of the interpolated surface from the mean satellite datapoints in each gridbox and **(f)** shows the distance from each grid to the nearest one containing satellite data.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 10. Map of the distribution of possible RMS error in the DEM calculated using a multiple regression based on airborne validation data.

TCD

2, 843–872, 2008

A new Antarctic digital elevation model: validation and errors

J. A. Griggs and J. L. Bamber

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion