

Fluctuations of  
a Greenlandic  
tidewater glacier

J. M. Lea et al.

# Fluctuations of a Greenlandic tidewater glacier driven by changes in atmospheric forcing: observations and modelling of Kangiata Nunaata Sermia, 1859–present

J. M. Lea<sup>1</sup>, D. W. F. Mair<sup>1</sup>, F. M. Nick<sup>2,3</sup>, B. R. Rea<sup>1</sup>, D. van As<sup>4</sup>, M. Morlighem<sup>5</sup>,  
P. W. Nienow<sup>6</sup>, and A. Weidick<sup>4</sup>

<sup>1</sup>Department of Geography and the Environment, University of Aberdeen, Elphinstone Road, AB24 3UF, UK

<sup>2</sup>The University Centre in Svalbard (UNIS), P.O. Box 156, 9171 Longyearbyen, Norway

<sup>3</sup>Centre for Ice and Climate, Niels Bohr Institute, University of Copenhagen, Copenhagen, 2100, Denmark

<sup>4</sup>Geological Survey of Denmark and Greenland (GEUS), Copenhagen, Denmark

<sup>5</sup>University of California, Irvine, Department of Earth System Science, Croul Hall, Irvine, CA 92697-3100, USA

<sup>6</sup>School of GeoSciences, University of Edinburgh, Drummond Street, Edinburgh, EH8 9XP, UK

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 4 April 2014 – Accepted: 14 April 2014 – Published: 24 April 2014

Correspondence to: J. M. Lea (j.lea@abdn.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

TCD

8, 2005–2041, 2014

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Many tidewater glaciers in Greenland are known to have undergone significant retreat during the last century following their Little Ice Age maxima. Where it is possible to reconstruct glacier change over this period, they provide excellent records for comparison to climate records, and calibration/validation for numerical models. These records therefore allow tests of numerical models that seek to simulate tidewater glacier behaviour over multi-decadal to centennial timescales. Here we present a detailed record of behaviour from Kangiata Nunaata Sermia (KNS), SW Greenland, between 1859–2012 and compare it against available oceanographic and atmospheric temperature variability between 1871–2012. We also use these records to evaluate the ability of a well-established one-dimensional flow-band model to replicate behaviour for the observation period. The record of terminus change demonstrates that KNS has advanced/retreated in phase with atmosphere and ocean climate anomalies averaged over multi-annual to decadal timescales. Results from an ensemble of model runs demonstrate that observed dynamics can be replicated, with changes in atmospheric forcing not needing to be offset by changes in oceanic forcing sensitivity. Furthermore, successful runs always require a significant atmospheric forcing component, while an oceanic forcing component is not always needed. Although the importance of oceanic forcing cannot be discounted, these results demonstrate that changes in atmospheric forcing are likely to be a primary driver of the terminus fluctuations of KNS from 1859–2012.

## 1 Introduction

Calving from tidewater glaciers (TWGs) accounts for up to 50 % of the mass loss from the Greenland Ice Sheet (Van den Broeke et al., 2009). Thus determining controls on tidewater glacier dynamics over decadal to centennial timescales is crucial to understanding their contribution to sea level in a warming climate (Alley et al., 2010; Vieli

TCD

8, 2005–2041, 2014

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Nick, 2011). The ability to achieve this in Greenland has been restricted in part by the relative lack of TWG terminus observations prior to the satellite age, and evidence of terminus locations being spread across a disparate array of sources. However, the synthesis of these sources has previously allowed multi-decadal to centennial records of TWG glacier behaviour to be reconstructed (e.g. Csatho et al., 2008; Bjørk et al., 2012; Weidick et al., 2012).

The generation of such records provide potentially excellent calibration and validation records for numerical modelling efforts (Vieli and Nick, 2011). That is to say that numerical models that are capable of replicating observed terminus behaviour over decadal to centennial timescales will be better placed to predict the future behaviour of a TWG over similar timescales. Despite this, there remain few examples of modelling efforts that have attempted to calibrate their results against multi-decadal observational records (e.g. Colgan et al., 2012). The ability of most numerical models to replicate dynamics over such timescales using realistic inputs therefore remains largely untested.

By undertaking calibration/validation exercises, the sensitivity of terminus position to different climatically forced processes can also be evaluated (e.g. Nick et al., 2013; Cook et al., 2013; Lea et al., 2014a). This is achieved by comparing the sensitivity of a modelled glacier to climate forcing against observed behaviour (Nick et al., 2013). With a knowledge of realistic ranges of forcing, this allows evaluation of the relative importance of each in contributing to the observed TWG behaviour.

Changes in oceanic forcing are significant drivers of TWG retreat in Greenland (Murray et al., 2010; Straneo et al., 2010; Rignot et al., 2012), but their relative importance between glaciers appears to be dependent on geographical location, glacier geometry (Nick et al., 2013), and potentially fjord connectivity with the open ocean (Straneo et al., 2012). Model based studies have also helped to demonstrate the sensitivity of some major outlet glaciers to air temperature changes (via enhanced runoff increasing crevasse water depth; Nick et al., 2013; Cook et al., 2013).

Where multi-decadal to centennial timescale climate data exist alongside records of terminus position, these provide the potential for robust evaluation of both numer-

ical models and the importance of different drivers of TWG terminus change. In this study we present a detailed record of terminus fluctuations at Kangiata Nunaata Sermia (KNS), SW Greenland from 1859–present. We then use this to evaluate the ability of a well-established numerical ice-flow model, driven by climate data, to replicate the pattern and timing of change at KNS during this period. Results of this are used to evaluate the sensitivity of KNS to climate forcing data over centennial timescales.

## 2 Field site and climate data

KNS is the largest TWG on the west coast of Greenland, south of Jakobshavn Isbræ (Fig. 1; Van As et al., 2014). It is known to have undergone significant retreat since its Little Ice Age maximum (Weidick et al., 2012), retreating a total of 22.6 km, with at least 12 km of this retreat occurring prior to 1859 when climate forcing data are unavailable (Lea et al., 2014a). It is situated  $\sim 100$  km inland from Nuuk at the head of Godthåbsfjord, and currently has a calving flux of  $\sim 6 \text{ km}^3 \text{ a}^{-1}$  (Van As et al., 2014).

A continuous record of mean monthly air temperature is available at Nuuk from 1866–present (Vinther et al., 2006; Cappelen et al., 2012). Temperatures at Nuuk are known to be strongly correlated to those near to KNS throughout the year (Taurisano et al., 2004), meaning that the Nuuk record can be used as an indicator of the atmospheric forcing at KNS.

As with all TWGs around Greenland, there are no long observational records of fjord water temperatures adjacent to KNS, though detailed hydrographic studies of the fjord have been undertaken recently (Mortensen et al., 2011, 2013). A shallow  $\sim 80$  m sill at the entrance to the fjord at Nuuk has been suggested to limit the connectivity of the fjord to warm ocean waters at depth. In fjords where shallow sills do not exist, the incursion of these warm ocean waters are thought to have significantly affected the stability of TWGs (Rignot et al., 2012; Straneo et al., 2012). The presence of the shallow sill in Godthåbsfjord also results in significant tidal mixing at the fjord entrance, allowing sea surface waters to be incorporated at depth which are then advected into

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the fjord (Mortensen et al., 2011). These intermediate level mixed waters have been proposed to significantly influence the energy available for submarine melting at the termini of the TWGs in Godthåbsfjord (Mortensen et al., 2013).

Due to the impact of surface waters near the fjord entrance on the energy balance of the fjord (Mortensen et al., 2011, 2013), and the potentially restricted influence of warm coastal currents at depth (Straneo et al., 2012), we suggest that sea surface temperatures (SSTs) provide a good indicator of the relative oceanographic forcing affecting KNS. Such data have also been used to good effect elsewhere as an indicator of oceanographic forcing where observations at depth are unavailable (e.g. McFadden et al., 2011; Bevan et al., 2012). The HadISST1  $1^\circ \times 1^\circ$  dataset provides SST estimates for the period 1871–present (Rayner et al., 2003), with annual averages for the area immediately offshore from Nuuk ( $62^\circ$  to  $64^\circ$  N  $51^\circ$  to  $53^\circ$  W) used as an indicator of oceanographic conditions affecting Godthåbsfjord. Although the data used will in part be based on interpolation (especially in the earlier part of the record), the data have been validated for west Greenland against independent records back to 1875 (Hanna et al., 2009). This therefore provides confidence in the results obtained from the HadISST1 dataset.

### 3 Glacier reconstruction data

By 1859 KNS is known to have retreated between 12–15 km from its Little Ice Age (LIA) maximum extent (Lea et al., 2014a). The post-LIAmax glacial geomorphology of KNS has been mapped, while previous analysis of a photograph taken in the 1850s, and a map published in 1859 places the terminus position somewhere inside the limit of a significant glacier readvance/stillstand (Lea et al., 2014a). We refer to this as the Akullersuaq Stade (after the headland that its maximum extent adjoins), previously named as the “1920 Stade” (Weidick et al., 2012). This is done due to the uncertainty of whether the glacier was at its maximum in 1920.

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Where the full terminus cannot be observed in photographs, terminus position is determined indirectly using GIS based analyses described below, in conjunction with evidence from maps (e.g. Lea et al., 2014a). Subsequent to 1921, intermittent direct observations of the terminus are available enabling mapping of terminus positions from a combination of ground-based, oblique-aerial, vertical-aerial, and satellite imagery (list of sources in Table 1).

Landsat panchromatic band imagery was used to map terminus position for 1987–2012. Cloud-free Landsat scenes were selected for analysis, acquired as late in the melt season as possible/just after its end. The start of November was used as the latest date from which images could be selected, since beyond this, mélange in the fjord has been observed to freeze, causing the terminus to advance (Mortensen et al., 2011; Sole et al., 2011). The majority of images were acquired during September or October, though for 1993 and 2003 cloud-free images were only available for dates in August (30 August 1993 and 8 September 2003 respectively). No suitable images were available for the years 1988–1991 and 1998, meaning that annual resolution rates of terminus change were acquired for 1992–1997 and 1999–2012 (Table 1).

Where more than 1 year separated terminus observations, annually averaged rates of change were calculated. This provides a continuous record of the trends in behaviour, and inter-annual variability of KNS for the period spanning 1859–2012. This behaviour could then be directly compared to atmospheric and oceanic climate data.

Each terminus position was determined using an adaptation of the box method (Moon and Joughin, 2008; Howat and Eddy, 2011), called the Curvilinear Box Method (CBM; see Lea et al., 2014b, for details). This has a marked advantage over the centreline tracking or standard box methods as it is capable of accounting for changes in terminus geometry, while also accurately tracking changes in fjord orientation. Furthermore, the box used to calculate terminus change is always centred on the glacier/fjord centreline, which is also the flowline used for the numerical model. Consequently, terminus positions and observed distances of change derived using the CBM can be compared directly to model output.

## 4 The TWG Model

The numerical model used is specifically designed to simulate the dynamics of TWGs along a flowband (Nick et al., 2010). It has been successful in replicating the dynamics of marine terminating outlets in both Greenland (e.g. Vieli and Nick, 2011; Nick et al., 2012; Lea et al., 2014a) and Antarctica (Jamieson et al., 2012, 2014), and also been used to make centennial timescale projections of the future contribution of Greenland's major TWG outlets to global sea level (Nick et al., 2013). The model accounts for basal, lateral and longitudinal shear stresses, and includes a robust treatment of grounding line dynamics (Pattyn et al., 2012). Bed topography data for the majority of the catchment are provided by Bamber et al. (2001), though the lower 40 km is generated using a mass continuity based bed reconstruction (Morlighem et al., 2011), validated by OIB/CReSIS flightlines (Gogineni et al., 2001). Where available, fjord bathymetry data are also used where KNS has retreated following its LIAMax (Weidick et al., 2012). Sensitivity analyses conducted by Lea et al. (2014a; their Fig. 10) for this bed configuration demonstrated that the model exhibits broadly comparable patterns of retreat behaviour where bed topography is varied within an uncertainty of  $\pm 50$  m.

A constant height vs. SMB relation is used to calculate SMB for the ablation zone of KNS (Eq. 1). This is derived from the average RACMO SMB model output for 1958–2007 (Van Angelen et al., 2013).

$$b(x) = 0.0018 \times h(x) - 2.693 \quad (1)$$

Where  $b(x)$  = SMB for position  $x$  on the model flowline, and  $h(x)$  = glacier elevation for position  $x$  on the flowline. Due to the tendency for over-estimation of accumulation in RACMO in this region (Van As et al., 2014), positive SMB values in the upstream section of the modelled glacier are prescribed to allow the glacier to maintain its contemporary elevation profile. Irrespective of this, SMB variability has previously been demonstrated to be of minimal importance to results of modelled TWG dynamics over the timescales that are being investigated (Lea et al., 2014a). The model is initialised

TCD

8, 2005–2041, 2014

### Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



using a glacier geometry approximating that of the Akullersuaq Stade maximum (ASM), derived from geomorphological mapping of associated trimlines (Fig. 1). Constants and parameter values used are summarised in Table 2, while the initial tuning procedure followed for this configuration is the same as that used by Lea et al. (2014a). Surface runoff (Van As et al., 2014)/air temperature (JJA average) and SST (annual average) data are used to drive changes in crevasse water depth ( $d_w$ ) and submarine melting ( $M$ ) respectively. Seasonal variability in basal and lateral sliding is not included due to its negligible importance over multidecadal timescales (Nick et al., 2013). Parameters which control the model sensitivity to climate forcing are derived using Monte Carlo methods described below.

### 4.1 Relating $d_w$ to air temperature

Changes in the value of  $d_w$  have previously been related to runoff variability (e.g. Nick et al., 2010; Cook et al., 2012, 2013), and have been successfully used as a climate linked forcing directly affecting terminus change (Nick et al., 2013). However, the only previously used scaling of surface runoff to  $d_w$  requires a baseline  $d_w$  value to be prescribed, which it cannot fall below (Nick et al., 2013, their Eq. S3). We present a new, unrestricted parameterisation that relates seasonal changes in monthly surface runoff to  $d_w$ , and allows  $d_w$  to freely evolve due to changes in annual runoff (Eq. 2).

$$d_{wNew} = d_{wPrev} + \alpha_1 \left( R_{year} \beta_{month} - \frac{R_{base}}{12} \right) \quad (2)$$

Where  $d_{wNew}$  = new crevasse water depth for a particular month,  $d_{wPrev}$  = crevasse water depth from the previous month,  $\alpha_1$  = coefficient relating crevasse water depth sensitivity to changes in runoff,  $R_{year}$  = total runoff for a given year ( $\text{Gtyr}^{-1}$ ),  $\beta_{month}$  = fraction of annual runoff occurring in a particular month,  $R_{base}$  = a baseline annual runoff total ( $\text{Gtyr}^{-1}$ ), equivalent to the annual volume of water that is either refrozen within the glacier or drains from the crevasse to the bed. This assumes that the rate of refreezing/drainage of water from crevasses is constant from year to year. Where annual

runoff exceeds  $R_{\text{base}}$ , the average annual  $d_w$  will therefore increase, and where runoff falls below  $R_{\text{base}}$ , the average annual  $d_w$  will decrease. Dividing annual runoff into each month's contribution also allows the direct incorporation of  $d_w$ 's seasonal variability. The value of  $d_w$  will therefore reach its annual minimum prior to the onset of the melt season, and peak in August. The coefficient  $\alpha_1$  allows the sensitivity of  $d_w$  to changes in runoff to be adjusted, and is used as a tuning parameter.

## 4.2 Definition of $\beta_{\text{month}}$

The fraction of annual runoff occurring in each month,  $\beta_{\text{month}}$ , is derived from analysis of each month's average runoff from the catchments of both KNS and Akullersuaq Sermia (AS) over the period 1960–2012, as given by high resolution surface mass balance (SMB) modelling of the region (Van As et al., 2014). The runoff values for KNS and AS are summed since the glaciers were confluent for much of the time since their LIAMax, including a significant portion of the period of interest of this study (see below; Wedick et al., 2012). Monthly runoff estimates were generated using both Modèle Atmosphérique Régional (MARv3.2; Fettweis et al., 2011), and the Regional Atmospheric Climate Model (RACMO2; Van Angelen et al., 2013). The variability in the monthly fraction of annual runoff is shown for both models in Fig. 2, with each producing similar patterns and magnitudes of monthly variability. We took the median result from the monthly averages of the two models. This pattern of monthly variability was kept constant from year to year for each model run.

While the model can be forced directly with annual modelled runoff values for the period 1960–2012 (Van As et al., 2014), no such values are available for the century before. Runoff values prior to 1960 are therefore estimated using the relation that exists between average June, July, August (JJA) air temperatures ( $A_{\text{JJA}}$ ) from Nuuk for 1960–2012 (Cappelen et al., 2012) and the modelled runoff values ( $r = 0.75$ ). A regression equation is generated from this (Eq. 3), allowing runoff estimates ( $\text{Gt a}^{-1}$ ) for the period 1866–1959 to be made from the Nuuk air temperature ( $^{\circ}\text{C}$ ) record (Vinther et al., 2006;

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Where  $x_{\text{conf}}$  = distance of the terminus through the confluence, and  $X_{\text{conf}}$  = the total flowline distance over which the confluence occurs. Due to uncertainty regarding the precise scaling of runoff to  $d_w$  as KNS retreats through its confluence with AS, and other confluence effects,  $\alpha_2$  is used as a tuning parameter within the model.

The extra ice flux contribution from AS when confluent with KNS is estimated to be approximately one sixth of that of KNS, based on the contemporary across glacier velocity profiles (Joughin et al., 2010), and terminus widths of AS and KNS. This extra flux is added to the modelled glacier as positive SMB at the confluence of KNS and AS, distributed along the flowline proportionate to the contemporary AS across glacier velocity profile (Lea et al., 2014a).

#### 4.4 Relating submarine melt rate to sea surface temperature

Submarine melt rate ( $M$ ) has previously been linearly related to deep ocean temperature (DOT) variability using a scaling coefficient (Nick et al., 2013; their Eq. S2). Using this parameterisation, the highest rates of  $M$  (expressed in this study in  $\text{km}^3 \text{a}^{-1}$ ) would also be associated with the highest inter-annual variability of  $M$ . This study therefore takes a slightly different approach in that (1)  $M$  is scaled to sea surface temperature (SST) rather than DOT, for reasons relating to fjord circulation explained above, and (2) we introduce a constant (minimum) baseline  $M$  rate,  $M_{\text{base}}$ , which is added to the linear relation to SST. We therefore calculate  $M$  ( $\text{km}^3 \text{a}^{-1}$ ) according to

$$M = M_{\text{base}} + \alpha_3 T_{\text{year}} \quad (6)$$

Where  $\alpha_3$  = submarine melt rate scaling coefficient, and  $T_{\text{year}}$  = annual average SST. This allows different minimum background  $M$  rates to be tested for different model runs, with different sensitivities of  $M$  to changes in SST superimposed upon this.

## 4.5 Model experiments and evaluation

Tuning parameters  $\alpha_1$ ,  $\alpha_2$ ,  $\alpha_3$  and  $M_{\text{base}}$  were varied randomly within prescribed limits for a total of 1500 Monte Carlo style model runs. The limits for each of the tuning parameters were: (1)  $\alpha_1$ , between 0 and 1.5, (2)  $\alpha_2$ , between 0.3 to 0.8, (3)  $M_{\text{base}}$ , between 0 to  $0.7 \text{ km}^3 \text{ a}^{-1}$ , and (4)  $\alpha_3$ , between 0 to 0.3. These ranges of  $\alpha_1$  and  $\alpha_2$  were chosen to reflect a wide range of potential forcing scenarios, while the values of  $M_{\text{base}}$  and  $\alpha_3$  were chosen so total  $M$  could potentially range from  $0 \text{ km}^3 \text{ a}^{-1}$  to values that exceed  $M$  rates that have been observed for other TWGs in western Greenland (Rignot et al., 2010; Enderlin and Howat, 2013). This allowed the feasibility of different potential drivers of the observed terminus change to be comprehensively assessed. Runs were conducted for the period 1871–2012, given that this is the period that both atmospheric and oceanic climate records are available for. The model was initialised at approximately the ASM profile and terminus position, as defined by the geomorphology, and given the duration of the spin up period to stabilise for the given forcing scenario. During spin up,  $d_w$  was allowed to freely evolve by up to  $\pm 3 \text{ m a}^{-1}$  to allow the terminus to stabilise at the ASM, with  $R_{\text{base}}$  and  $T_{\text{year}}$  held constant. These were defined as the 1871–1920 runoff average ( $3.107 \text{ Gtyr}^{-1}$ ) and SST average ( $2.605 \text{ }^\circ\text{C}$ ) respectively. These values were used for spin up as it is known the ASM was attained at some point within this window.

Model results were evaluated against their ability to replicate observed terminus dynamics, where absolute terminus positions are known (i.e. 1921 to 2012). The period from 1871–1920 therefore effectively becomes a transient spin up period, where the model is driven using real climate data though terminus position is only known within a range. The ability of each model run to replicate observed dynamics was determined using a weighted regression ( $R^2$ ) calculation, with the weighting of each terminus observation calculated according to

$$w_n = \frac{D_{n+1} - D_{n-1}}{2(D_k - D_1)} \quad \text{for } n = 1, 2, \dots, k \quad (7)$$

Where  $w$  = weighting in regression calculation,  $n$  = terminus observation,  $k$  = total number of terminus observations, and  $D$  = date of terminus observation. Each terminus observation is therefore temporally weighted according to the median length of time elapsed between the terminus observations that occur before and after observation  $n$ . This ensures that the evaluation of model performance is not biased towards the last  $\sim 20$  years where there is a comparatively high density of observations. Model runs were counted as successful where (1) the difference between the modelled and observed 1921 position was  $< 500$  m, (2) the weighted  $R^2 > 0.85$ , and (3) the gradient of the resulting line of regression was  $> 0.85$ .

## 5 Glacier reconstruction results

The geomorphology shows distinct upper and lower sets of lateral moraines on both sides of the fjord, with fluted moraines occupying the intervening space (Fig. 1a). The upper set are associated with the LIA maximum (Lea et al., 2014a), while the lower set were formed during the Akullersuaq Stade. Fridtjof Nansen's (1890) account of the first traverse of Greenland in 1888, includes a drawing from a photograph showing AS and KNS to be confluent, though the terminus position itself is not visible. Although the original image could not be traced or an exact date of acquisition determined, it is likely to have been taken some time near to the publication date of 1890.

Maps from 1859, 1860, 1866 and 1885 all show the terminus of KNS to be adjoining Akullersuaq and fully confluent with AS (Kleinschmidt, 1859; Poulsen, 1860; Brede, 1866; Rink, 1866; Jensen, 1885). While it is possible that some details on the maps were copied following Kleischmidt (1859), the addition of detail such as lakes on plateaus near to KNS by Jensen (1885) provides confidence that this map faithfully records the contemporary terminus position. There is nothing to suggest that KNS became diffluent from AS at any time from 1859–1885. However, due to a lack of map detail and the Nansen (1890) drawing not including the terminus, these sources cannot be used to provide absolute terminus positions.

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The earliest images of KNS are from the 1850s and 1903. Both are taken from approximately the same position, with the terminus partially obscured by foreground topography (Weidick et al., 2012). The presence of medial moraines in each image demonstrates that KNS was confluent with AS. Lea et al. (2014a) quantified the terminus position uncertainty for the 1850s photograph using viewshed analysis. Similar analysis has been undertaken for the 1903 image, showing that the uncertainty in terminus position is the same as for the 1850s image (Fig. 3). The maximum terminus extents for both images are therefore located behind a headland corresponding to the ASM on the eastern side of the fjord (Figs. 1a and 3).

It is not currently possible to say when the ASM was attained from any observational evidence, only that it occurred sometime between 1859–1920. The climate anomalies for the period (compared to 1961–1990 baselines) show that air temperature (AT) and SST anomalies were, on average, antiphased for the period 1871–1903 (Fig. 4c and d), though AT and SST anomalies are in phase (negative/near-baseline) for 1903–1920. Conditions are therefore more likely to have been conducive for glacier advance during the latter period.

Terminus position was mapped directly for the remaining images, providing a record of 29 terminus positions spanning the period 1921–2012 (Figs. 1 and 4). The first direct terminus observation (1921) shows a slight retreat from the ASM. Subsequent to this, KNS retreated a total of 9.7 km at a non-uniform rate up to 2012, interrupted by short periods of readvance (Fig. 4a and b). Averaged retreat rates of  $-116 \text{ m a}^{-1}$  are observed between 1921–1946, before a rapid retreat of 3.9 km within the 2 year period from 1946–1948 (Figs. 1a and 4). Between 1948–1968 KNS retreated on average by  $-97 \text{ m a}^{-1}$ , before readvancing by  $+60 \text{ m a}^{-1}$  up to 1979 (Fig. 4b). A terrestrial photograph taken in 1965 with the majority of the terminus obscured shows the termini of KNS and AS to be fully diffluent.

The 1921–1968 period of sustained retreat was accompanied by positive average AT and SST anomalies (Fig. 4c and d). The highest AT anomalies occurred during the

period 1928–1941, though the largest retreat (between 1946–1948) occurred during a comparatively less extreme period of positive AT and SST (Fig. 4).

From 1979 to 1987 KNS retreated by  $-658$  m in total ( $-82$   $\text{m a}^{-1}$ ), before readvancing by  $+758$  m from 1987–1992 ( $+152$   $\text{m a}^{-1}$ ). Using the near complete 20 year annual record of terminus fluctuations from 1992–2012, KNS advanced for 4 out of 5 years between 1992–1997, followed by retreat in 11 out of 13 years from 1999–2012 at an average rate of  $-103$   $\text{m a}^{-1}$ . The latter included 8 annual retreats of  $> 100$  m, with the largest retreats occurring in 2004 ( $-438$  m) and 2005 ( $-316$  m). These periods of advance and retreat behaviour occurred during periods of in-phase negative and positive climate anomalies respectively.

Where temporal density of observations was high, terminus behaviour that was antiphased with the prevailing climate anomalies was also observed. Examples of this include a retreat of  $-626$  m observed in 1995 during negative climate anomalies, while two terminus advances occur in 2008 and 2009 despite markedly positive AT and SST anomalies (Fig. 4). At annual resolution, the magnitude of terminus retreat/advance was also found to be unrelated to the magnitude of either climate anomaly for each particular year.

## 6 Model results

From a total of 1500 model runs conducted, 29 runs (1.9 %) successfully replicated the observed dynamics of KNS according to the criteria outlined above (Fig. 5a). Following the initiation of climate forcing in 1871 (Fig. 5b and c), the results of each run are highly comparable up to 1884, with little modelled terminus change observed. Following this, for the period 1884 to  $\sim 1910$ , 6 of the 29 runs (21 %) show evidence of multi-annual terminus retreats and equivalent readvances of  $> 750$  m with periodicities of 2–4 years. A further 7 runs (24 %) show evidence of at least one short lived ( $< 5$  year) oscillation in terminus position of  $> 750$  m between 1884 to 1920. None of these model runs signifi-

### Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cantly exceed the ASM position, and are thus in agreement with the geomorphological evidence presented, and the position of the 1921 terminus observation.

All model runs retreat to the observed 1936 position between modelled years 1929–1936, via a single retreat event of  $\sim 1$  km. Subsequent to this, modelled retreat to the observed 1946 position is gradual, before the model successfully replicates a large topographically controlled retreat from the 1946 position. There was varying success in modelling the exact timing of this retreat (observed between 1946–1948), with the model ensemble predicting it to occur anywhere between 1943–1962. The position where the modelled terminus restabilises following the retreat through the AS confluence is generally too far advanced by  $\sim 1$  km compared to the position following the 1946–1948 retreat. All model runs then go on to over-predict terminus extent for the 1968 observation by between 0.35 to 1.59 km.

Though no model runs exactly match the precise inter-annual terminus fluctuations from 1968–2012, they do capture the general multi-annual to decadal pattern of retreat observed. This is characterised by general terminus stability within a range of  $\pm 500$  m for the period 1968 to  $\sim 1999$ , before the terminus begins to retreat  $\sim 2$  km towards the 2012 position. All of the successful model runs identified predict KNS to be in a more retreated position in 2012 than observed by a range of 0.32 to 5.04 km. Where a significant difference between observed and modelled terminus positions has occurred by the end of the model run in 2012, the divergence begins in 2010 at the earliest.

The distributions of tuning parameters for successful runs are shown in Fig. 6, with the distribution of all histograms shown to be non-normal. Submarine melting related tuning parameters,  $\alpha_3$ , and  $M_{\text{base}}$ , tended towards the mid to lower ends of the ranges tested (Fig. 6c and d). Values of  $\alpha_3$  peak between 0.075 to 0.1, though there is no clearly defined peak in the distribution of  $M_{\text{base}}$  values.

In contrast, none of the  $d_w$  related tuning parameters ( $\alpha_1$  and  $\alpha_2$ ) approach 0 (Fig. 6a and b), with the lowest values being 0.412 and 0.389 respectively. Construction of a correlation matrix comparing all tuning parameter values for all successful runs also demonstrates a significant inverse relationship between the value of  $\alpha_1$ , and the AS

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



average terminus change rates can be mitigated by averaging change over timescales up to, or greater than a decade. For example, extending the 1992–1997 average ( $51 \text{ ma}^{-1}$  retreat) to cover the period 1987–1997 ( $91 \text{ ma}^{-1}$  advance) provides a more representative impression of multi-annual terminus behaviour, since 5 out of the 6 observations available show terminus advance. Interpreting absolute terminus change rate values should therefore be done with caution, and in most cases will be more representative of the average direction of terminus change rather than the absolute magnitude of annual change.

Taking into account uncertainties due to topographic controls on terminus stability, observations of terminus change over a period of several years are more likely to allow a more accurate evaluation of a TWG's response to climate forcings. However, for this study, deconvolving the relative importance of AT vs. SST in driving terminus change is difficult using observations alone, given that both climate drivers vary in phase for 1903–present. It could potentially be argued that AT is the primary driver of change, since the 33 year period of positive anomaly SST from 1871–1903 had relatively little impact on the terminus stability of KNS. However, fjord geometry could also have been a significant factor stabilising the terminus during this time. Arguably this becomes less likely when it is considered that while SST was similar for the period 1921–1948, positive AT allowed KNS to retreat through the same section of fjord and through its confluence with AS within  $26 \pm 1$  years (Fig. 4). However, given the lack of certainty in terminus position between 1871–1920, it is not possible to robustly verify these arguments.

## 7.2 Implications of modelling

The observed terminus behaviour of KNS from 1921–2012 was successfully replicated by 29 of 1500 model runs using surface runoff and SST records as drivers of terminus change. This demonstrates that the parameterisations used to scale these climate records to  $d_w$  and  $M$  respectively can successfully be used to simulate the observed pattern of behaviour of a tidewater glacier over centennial timescales. Where the ob-

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



servational record is of sufficient detail to resolve inter-annual terminus fluctuations (1992–2012), the model does not replicate these. This is to be expected given (1) the flowband nature of the model and associated depth and width integrations over each grid cell, meaning that fluctuations of terminus configurations such as the creation of calving bays cannot be replicated (e.g. Fig. 1b), (2) the uncertainty in fjord bathymetry and geometry potentially affecting relative terminus stability, and (3) the use of single terminus observations as notionally definitive indicators of annual terminus change, where the stochastic nature of calving and associated sub-annual terminus fluctuations make any direct one-to-one comparison of modelled results to annual resolution observations inappropriate. Valid comparison of model results to observations should therefore only be attempted over multi-annual timescales where terminus dynamics within calving bays, sub-annual calving events and fine scale uncertainties in fjord topography become comparatively less significant.

For successful model runs, the interrelationships between the parameter values that determine  $d_w$  and  $M$  sensitivity to the climate records also inform the relative importance of changes in atmospheric and oceanic forcing in driving terminus change. The lack of any significant relationship between  $\alpha_1$  and  $\alpha_3$  demonstrates that a change in model sensitivity to surface runoff is not offset by any change in model sensitivity to SST. Taken alone, this evidence indicates that either atmospheric forcing (via surface runoff) dominates oceanic forcing (via SST), or vice versa. However, the occurrence of runs where  $\alpha_3$  does not significantly exceed 0 (i.e. where runs experience negligible  $M$  variability) demonstrate that the model can successfully reproduce observed behaviour with nearly no changes in oceanic forcing from year to year. Although some successful model runs did have significant inter-annual  $M$  variability (e.g. the maximum range of  $M$  values for an entire 141 year model run was  $0.76 \text{ km}^3 \text{ a}^{-1}$ ), each model run always requires significant atmospheric forcing variability to allow it to replicate observations. The importance of oceanic forcing variability can therefore not be entirely discounted.

The model demonstrates that knowledge of atmospheric forcing variability (via runoff), without needing to vary oceanic forcing, can be sufficient to reproduce real-

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



istic patterns of observed glacier behaviour at KNS over the last century. However, the precise physical mechanism by which air temperature could drive observed change requires further investigation. For example, though a combination of modelled and empirically estimated runoff values have been used to drive changes in  $d_w$  to force the model, subglacial runoff variability is also known to drive rates of submarine melting at the terminus (Jenkins, 2011; Xu et al., 2012; Sciascia et al., 2013). Therefore we do not rule out that the centennial behaviour observed could also be explained by calving driven by seasonal changes in submarine melt rates, that are in turn a function of subglacial runoff (e.g. Sciascia et al., 2013).

The relative insensitivity to changes in oceanic forcing is not necessarily surprising given the hydrographic setting of KNS – located at the end of a > 100 km long fjord system that is thought to be largely insulated from changes in ocean conditions due to the presence of a shallow sill at its entrance (Mortensen et al., 2011, 2013). This has previously been used to suggest that recent changes in ocean conditions (e.g. Straneo and Heimbach, 2013) have not affected the dynamics of KNS significantly (Straneo et al., 2012). The results presented here are therefore compatible with this argument.

The over-estimation of terminus retreat by 2012 of every successful run is thought to result from the poor knowledge of fjord width geometry beyond the contemporary glacier terminus. Upstream of the 2012 terminus, the lateral ice margins are used to define model glacier width, leading to a likely over-estimation of the prescribed fjord width. The divergence between the actual and prescribed fjord width is therefore likely to increase upglacier, increasing the likelihood of model error in this area. This explains why significant divergence from the observational record only occurs once the modelled terminus has retreated ~ 1.5 km beyond the 2012 terminus. Any attempt at modelling the future fluctuations of KNS will therefore require both improvements to subglacial topography estimates and comprehensive assessments of fjord width uncertainties as part of any predictions.

## 8 Conclusions

Utilising multiple lines of evidence, it has been possible to reconstruct terminus fluctuations of KNS from 1859–2012. This study therefore completes the record of terminus fluctuations of KNS from its LIAMax, in 1761, up to the present (Lea et al., 2014a), providing one of the longest, and most detailed records of observed TWG change in Greenland. Results from numerical modelling show that the fluctuations of KNS can be simulated through parameterisations that link surface runoff to a crevasse water depth based calving criterion. Changes in both/either crevasse water depth and/or runoff driven rates of submarine melt are therefore suggested as potential drivers of observed change. Although ocean driven changes in submarine melt rates are not always required for the model to replicate the observed length variations of KNS, results do not allow their importance to be discounted entirely.

Observations of KNS show it to respond in phase with AT and SST anomalies over multi-annual to decadal timescales from at least 1921–2012. However, where inter-annual comparisons to AT and SST are possible (1992–2012), climatically anti-phased terminus fluctuations are observed. This highlights the inherent noisiness of terminus response over short timescales, the potential importance of antecedence, and the dangers of using similarly short calibration periods for predictive modelling efforts.

Results from numerical modelling successfully capture the terminus dynamics of KNS over multi-annual to decadal timescales, though not precise inter-annual fluctuations. This is due to a combination of uncertainties in fjord topography, and the approximations inherent to the depth and width integrations associated with using a one-dimensional flow-band model.

Nevertheless, this study demonstrates that simple flow-band numerical models of tidewater glaciers can be used to capture TWG dynamics over centennial timescales. This provides validation that these models can be useful tools for both palaeo- and contemporary/prognostic modelling efforts. However, the primary challenge to their use as predictive tools remain the accurate definition of subglacial topography and fjord

### Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



width, which exert dominant controls on modelled glacier stability. Any future efforts at prognostic modelling of TWGs should therefore seek to account for these uncertainties in addition to those associated with sensitivity to climate forcing.

*Acknowledgements.* RACMO2.1 data were provided by Jan van Angelen and Michiel van den Broeke, IMAU, Utrecht University. MAR v3.2 data used for runoff calculations were provided by Xavier Fettweis, Department of Geography, University of Liège. The photogrammetric DEM used in Figs. 1 and 3 was provided by Kurt H. Kjær, Centre for GeoGenetics, University of Copenhagen. This research was financially supported by J.L.'s PhD funding, NERC grant number: NE/I528742/1. Support for F.M.N. was provided by the Conoco-Phillips/Lundin Northern Area Program CRIOS project (Calving Rates and Impact on Sea Level).

## References

- Alley, R. B., Andrews, J. T., Brigham-Grette, J., Clarke, G. K. C., Cuffey, K. M., Fitzpatrick, J. J., Funder, S., Marshall, S. J., Miller, G. H., Mitrovica, J. X., Muhs, D. R., Otto-Bliesner, B. L., Polyak, L., and White, J. W. C.: History of the Greenland Ice Sheet: paleoclimatic insights, *Quaternary Sci. Rev.*, 29, 1728–1756, doi:10.1016/j.quascirev.2010.02.007, 2010.
- Bamber, J. L., Layberry, R. L., and Gogineni, S. P.: A new ice thickness and bed data set for the Greenland ice sheet: 1. Measurement, data reduction, and errors, *J. Geophys. Res.*, 106, 33773–33780, doi:10.1029/2001JD900054, 2001.
- Bevan, S. L., Luckman, A. J., and Murray, T.: Glacier dynamics over the last quarter of a century at Helheim, Kangerdlugssuaq and 14 other major Greenland outlet glaciers, *The Cryosphere*, 6, 923–937, doi:10.5194/tc-6-923-2012, 2012.
- Brede, N.: Skizze Kaart over Vestkysten af Grønland fra Arsuk til Holsteensborg, KBK Netpublikation DK003200, Copenhagen, 1866.
- Bjørk, A. A., Kjær, K. H., Korsgaard, N. J., Khan, S. A., Kjeldsen, K. K., Andresen, C. S., Larsen, N. K., and Funder, S.: An aerial view of 80 years of climate-related glacier fluctuations in southeast Greenland, *Nat. Geosci.*, 5, 427–432, doi:10.1038/ngeo1481, 2012.
- Bruun, D.: Oversigt over Norburuiner i Godthaab og Frederikshaab Distrikter, *Meddelelser om Grønland*, 56, 55–147, 1917.
- Cappelen, J.: Greenland-DMI Historical Climate Data Collection 1873–2011, Technical Report 12-04, Danish Meteorological Institute, Copenhagen, 2012.

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Colgan, W., Pfeffer, W. T., Rajaram, H., Abdalati, W., and Balog, J.: Monte Carlo ice flow modeling projects a new stable configuration for Columbia Glacier, Alaska, c. 2020, *The Cryosphere*, 6, 1395–1409, doi:10.5194/tc-6-1395-2012, 2012.

5 Cook, S., Zwinger, T., Rutt, I. C., O’Neel, S., and Murray, T.: Testing the effect of water in crevasses on a physically based calving model, *Ann. Glaciol.*, 53, 90–96, doi:10.3189/2012AoG60A107, 2012.

Cook, S., Rutt, I. C., Murray, T., Luckman, A., Selmes, N., Goldsack, A., and Zwinger, T.: Modelling environmental influences on calving at Helheim Glacier, East Greenland, *The Cryosphere Discuss.*, 7, 4407–4442, doi:10.5194/tcd-7-4407-2013, 2013.

10 Csatho, B., Schenk, T., Van der Veen, C. J., and Krabill, W. B.: Intermittent thinning of Jakobshavn Isbræ, West Greenland, since the Little Ice Age, *J. Glaciol.*, 54, 131–144, doi:10.3189/002214308784409035, 2008.

Rink, H.: *Kyststrækning fra Frederikshåb i syd til Napasaq i nord*, KBK Netpublikation RI000029, Copenhagen, 1866.

15 Fettweis, X., Tedesco, M., van den Broeke, M., and Ettema, J.: Melting trends over the Greenland ice sheet (1958–2009) from spaceborne microwave data and regional climate models, *The Cryosphere*, 5, 359–375, doi:10.5194/tc-5-359-2011, 2011.

Gogineni, S., Tammana, D., Braaten, D., Leuschen, C., Akins, T., Legarsky, J. K., Kanagaratnam, P., Stiles, J., Allen, C., and Jezek, K.: Coherent radar ice thickness measurements over the Greenland ice sheet, *J. Geophys. Res.*, 106, 33761–33772, doi:10.1029/2001JD900183, 2001.

Hanna, E., Cappelen, J., Fettweis, X., Huybrechts, P., Luckman, A., and Ribergaard, M. H.: Hydrologic response of the Greenland ice sheet: the role of oceanographic warming, *Hydrol. Process.*, 23, 7–30, doi:10.1002/hyp.7090, 2009.

25 Howat, I. M. and Eddy, A.: Multi-decadal retreat of Greenland’s marine-terminating glaciers, *J. Glaciol.*, 57, 389–396, doi:10.3189/002214311796905631, 2011.

Hvidegaard, S. M., Sorensen, L. S., and Forsberg, R.: ASTER GDEM validation using LiDAR data over coastal regions of Greenland, *Remote Sensing Letters*, 3, 85–91, doi:10.1080/01431161.2010.527389, 2012.

30 Jamieson, S. S. R., Vieli, A., Livingstone, S. J., Cofaigh, C. Ó., Stokes, C., Hillenbrand, C. D., and Dowdeswell, J. A.: Ice-stream stability on a reverse bed slope, *Nat. Geosci.*, 5, 799–802, doi:10.1038/ngeo1600, 2012.

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Jamieson, S. S. R., Vieli, A., Cofaigh, C. Ó, Stokes, C. R., Livingstone, S. J., and Hillenbrand, C.-D.: Understanding controls on rapid ice-stream retreat during the last deglaciation of Marguerite Bay, Antarctica, using a numerical model, *J. Geophys. Res.*, 119, 247–263, doi:10.1002/2013JF002934, 2014.

Jenkins, A.: Convection-driven melting near the grounding lines of ice shelves and tidewater glaciers, *J. Phys. Oceanogr.*, 41, 2279–2294, doi:10.1175/JPO-D-11-03.1, 2011.

Jensen, J. A. D.: Vestkysten af Grønland fra Arsuk til Holstensborg 61° til 67° N, KBK Netpublikation, DK003198, Copenhagen, 1885

Joughin, I., Smith, B. E., Howat, I. M., Scambos, T., and Moon, T.: Greenland flow variability from ice-sheet-wide velocity mapping, *J. Glaciol.*, 56, 415–430, doi:10.3189/002214310792447734, 2010.

Kleinschmidt, S.: Godthåbs distrikt (hertil en Navneliste), KBK Netpublikation RI000074, Copenhagen, 1859.

Lea, J. M., Mair, D. W. F., Nick, F. M., Rea, B. R., Weidick, A., Kjær, K., Morlighem, M., van As, D., and Schofield, J. E.: Terminus-driven retreat of a major southwest Greenland tidewater glacier during the early 19th century: insights from glacier reconstructions and numerical modelling, *J. Glaciol.*, 220, 333–344, doi:10.3189/2014JoG13J163, 2014a.

Lea, J. M., Mair, D. W. F., and Rea, B. R.: Evaluation of existing and new methods of tracking glacier terminus change, *J. Glaciol.*, 220, 322–332, doi:10.3189/2014JoG13J061, 2014b.

Morlighem, M., Rignot, E., Seroussi, H., Larour, E., Ben Dhia, H., and Aubry, D.: A mass conservation approach for mapping glacier ice thickness, *Geophys. Res. Lett.*, 38, L19503, doi:10.1029/2011GL048659, 2011.

McFadden, E. M., Howat, I. M., Joughin, I., Smith, B. E., and Ahn, Y.: Changes in the dynamics of marine terminating outlet glaciers in west Greenland (2000–2009), *J. Geophys. Res.*, 116, F02022, doi:10.1029/2010JF001757, 2011.

Moon, T. and Joughin, I.: Changes in ice front position on Greenland's outlet glaciers from 1992 to 2007, *J. Geophys. Res.*, 113, F02022, doi:10.1029/2007JF000927, 2008.

Mortensen, J., Lennert, K., Bendtsen, J., and Rysgaard, S.: Heat sources for glacial melt in a sub-Arctic fjord (Godthåbsfjord) in contact with the Greenland Ice Sheet, *J. Geophys. Res.*, 116, C01013, doi:10.1029/2010JC006528, 2011.

Mortensen, J., Bendtsen, J., Motyka, R. J., Lennert, K., Truffer, M., Fahnestock, M., and Rysgaard, S.: On the seasonal freshwater stratification in the proximity of fast-flowing

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tidewater outlet glaciers in a sub-Arctic sill fjord, *J. Geophys. Res.*, 118, 1382–1395, doi:10.1002/jgrc.20134, 2013.

Murray, T., Scharrer, K., James, T. D., Dye, S. R., Hanna, E., Booth, A. D., Selmes, N., Luckman, A., Hughes, A. L. C., Cook, S., and Huybrechts, P.: Ocean regulation hypothesis for glacier dynamics in southeast Greenland and implications for ice sheet mass changes, *J. Geophys. Res.*, 115, F03026, doi:10.1029/2009JF001522, 2010.

Nansen, F.: *The First Crossing of Greenland*, Longmans, London, 1890.

Nick, F. M., Van der Veen, C. J., Vieli, A., and Benn, D. I.: A physically based calving model applied to marine outlet glaciers and implications for the glacier dynamics, *J. Glaciol.*, 56, 781–794, doi:10.3189/002214310794457344, 2010.

Nick, F. M., Luckman, A., Vieli, A., Van der Veen, C. J., Van As, D., Van de Wal, R. S. W., Pattyn, F., Hubbard, A. L., and Floricioiu, D.: The response of Petermann Glacier, Greenland, to large calving events, and its future stability in the context of atmospheric and oceanic warming, *J. Glaciol.*, 58, 229–239, doi:10.3189/2012JoG11J242, 2012.

Nick, F. M., Vieli, A., Andersen, M. L., Joughin, I., Payne, A., Edwards, T. L., Pattyn, F., van de Wal, R. S.: Future sea-level rise from Greenland's main outlet glaciers in a warming climate, *Nature*, 497, 235–238, doi:10.1038/nature12068, 2013.

Pattyn, F., Schoof, C., Perichon, L., Hindmarsh, R. C. A., Bueler, E., de Fleurian, B., Durand, G., Gagliardini, O., Gladstone, R., Goldberg, D., Gudmundsson, G. H., Huybrechts, P., Lee, V., Nick, F. M., Payne, A. J., Pollard, D., Rybak, O., Saito, F., and Vieli, A.: Results of the Marine Ice Sheet Model Intercomparison Project, MISIP, *The Cryosphere*, 6, 573–588, doi:10.5194/tc-6-573-2012, 2012.

Poulsen, J.: *Godthåb Kommune, KBK Netpublikation RI000059*, Copenhagen, 1860.

Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *J. Geophys. Res.*, 108, 4407, doi:10.1029/2002JD002670, 2003.

Rignot, E., Fenty, I., Menemenlis, D., and Xu, Y.: Spreading of warm ocean waters around Greenland as a possible cause for glacier acceleration, *Ann. Glaciol.*, 53, 257–266, doi:10.3189/2012AoG60A136, 2012.

Roussel, A.: *Farms and churches of the medieval Norse settlement in Greenland*, *Meddelelser om Grønland*, 89, 342 pp., 1941.

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sciascia, R., Straneo, F., Cenedese, C., and Heimbach, P.: Seasonal variability of submarine melt rate and circulation in an East Greenland fjord, *J. Geophys. Res.*, 118, 2492–2506, doi:10.1002/jgrc.20142, 2013.

5 Sole, A. J., Mair, D. W. F., Nienow, P. W., Bartholomew, I. D., King, M. A., Burke, M. J., and Joughin, I.: Seasonal speedup of a Greenland marine-terminating outlet glacier forced by surface melt-induced changes in subglacial hydrology, *J. Geophys. Res.*, 116, F03014, doi:10.1029/2010JF001948, 2011.

Straneo, F. and Heimbach, P.: North Atlantic warming and the retreat of Greenland's outlet glaciers, *Nature*, 504, 36–43, doi:10.1038/nature12854, 2013.

10 Straneo, F., Hamilton, G. S., Sutherland, D. A., Stearns, L. A., Davidson, F., Hammill, M. O., Stenson, G. B., and Rosing-Asvid, A.: Rapid circulation of warm subtropical waters in a major glacial fjord in East Greenland, *Nat. Geosci.*, 3, 182–186, doi:10.1038/ngeo764, 2010.

Straneo, F., Sutherland, D. A., Holland, D., Gladish, C., Hamilton, G. S., Johnson, H. L., Rignot, E., Xu, Y., and Koppes, M.: Characteristics of ocean waters reaching Greenland's glaciers, *Ann. Glaciol.*, 53, 202–210, doi:10.3189/2012AoG60A059, 2012.

15 Taurisano, A., Bøggild, C. E., and Karlsen, H. G.: A century of climate variability and climate gradients from coast to ice sheet in West Greenland, *Geogr. Ann. A*, 86, 217–224, doi:10.1111/j.0435-3676.2004.00226.x, 2004.

Van Angelen, J. H., van den Broeke, M. R., Wouters, B., and Lenaerts, J. T. M.: Contemporary (1960–2012) evolution of the climate and surface mass balance of the Greenland ice sheet, *Surv. Geophys.*, doi:10.1007/s10712-013-9261-z, 2013.

20 Van As, D., Andresen, M. L., Petersen, D., Fettweis, X., van Angelen, J. H., Lenaerts, J. T. M., van den Broeke, M. R., Lea, J. M., Bayou, N., Bøggild, C. E., Ahlstrøm, A. P., and Steffen, K.: Increasing meltwater discharge from the Nuuk region of the Greenland ice sheet and implications for mass balance (1960–2012), *J. Glaciol.*, 220, 314–322, 2014.

25 Van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E., van de Berg, W. J., van Meijgaard, E., Velicogna, I., and Wouters, B.: Partitioning recent Greenland mass loss, *Science*, 326, 984–986, doi:10.1126/science.1178176, 2009.

30 Vieli, A. and Nick, F. M.: Understanding and modelling rapid dynamic changes of tidewater outlet glaciers: issues and implications, *Surv. Geophys.*, 32, 437–458, doi:10.1007/s10712-011-9132-4, 2011.

Vinther, B. M., Andersen, K. K., Jones, P. D., Briffa, K. R., and Cappelen, J.: Extending Greenland temperature records into the late eighteenth century, *J. Geophys. Res.*, 111, D11105, doi:10.1029/2005JD006810, 2006.

5 Weidick, A. and Citterio, M.: The ice-dammed lake Isvand, West Greenland, has lost its water, *J. Glaciol.*, 57, 186–188, doi:10.3189/002214311795306600, 2011.

Weidick, A., Bennike, O., Citterio, M., and Nørgaard-Pedersen, N.: Neoglacial and Historical Glacier Changes around Kangersuneq Fjord in Southern West Greenland, *Geological Survey of Denmark and Greenland*, Copenhagen, 2012.

10 Xu, Y., Rignot, E., Menemenlis, D., and Koppes, M.: Numerical experiments on subaqueous melting of Greenland tidewater glaciers in response to ocean warming and enhanced subglacial discharge, *Ann. Glaciol.*, 53, 229–234, doi:10.3189/2012AoG60A139, 2012.

**Fluctuations of  
a Greenlandic  
tidewater glacier**

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** List of terminus observations and acquisition dates.

Acquisition date	Observation type	Source
1850s	Terrestrial photo'	H. Rink (in Weidick et al., 2012)
1859	Map	Kleinschmidt (1859)
1860	Map	Poulsen (1860)
1866	Map	Rink (1866)
1866	Map	Falbe (1866)
1885	Map	Jensen (1885)
1880s?	Sketch (after photo')	Nansen (1890)
1903	Terrestrial photo'	J. Møller in Bruun (1917)
1921	Terrestrial photo'	A. Nissen in Weidick et al. (2012)
1932	Terrestrial photo'	A. Roussell in Roussell (1941)
27 Aug 1936	Oblique photo'	Weidick et al. (2012)
10 Aug 1946	Oblique photo'	Weidick et al. (2012)
20 Aug 1948	Oblique photo'	Weidick et al. (2012)
21 Jun 1965	Terrestrial photo'	Weidick et al. (2012)
16 Aug 1968	Aerial photo'	USGS
15 Sep 1979	Terrestrial photo'	Weidick et al. (2012)
15 Sep 1987	Satellite	Landsat
19 Sep 1992	Satellite	Landsat
30 Aug 1993	Satellite	Landsat
18 Sep 1994	Satellite	Landsat
14 Oct 1995	Satellite	Landsat
14 Sep 1996	Satellite	Landsat
1 Sep 1997	Satellite	Landsat
15 Sep 1999	Satellite	Landsat
18 Sep 2000	Satellite	Landsat
22 Oct 2001	Satellite	Landsat
23 Sep 2002	Satellite	Landsat
9 Aug 2003	Satellite	Landsat
12 Sep 2004	Satellite	Landsat
24 Sep 2005	Satellite	Landsat
18 Sep 2006	Satellite	Landsat
27 Sep 2007	Satellite	Landsat
23 Sep 2008	Satellite	Landsat
19 Sep 2009	Satellite	Landsat
13 Sep 2010	Satellite	Landsat
16 Sep 2011	Satellite	Landsat
18 Sep 2012	Satellite	Landsat

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 2.** List of parameters and constants used for running the model.

Parameter/Constant	Value
Ice density – $\rho_i$	900 kg m <sup>-3</sup>
Meltwater density – $\rho_w$	1000 kg m <sup>-3</sup>
Proglacial water body density – $\rho_p$	1028 kg m <sup>-3</sup>
Gravitational acceleration – $g$	9.8 m s <sup>-2</sup>
Friction exponent – $m$	3
Friction parameters – $\mu$ and $\lambda$	1
Glen's flow law exponent – $n$	3
Glen's flow law coefficient – $A$	$4.5 \times 10^{-17}$ Pa <sup>-3</sup> a <sup>-1</sup>
Grid size	~ 250 m
Time step	0.005 a


## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

**Table 3.** Pearson correlation coefficient values for tuning parameters of successful model runs ( $n = 29$ ). Correlation coefficients with  $p$  values  $< 0.05$  are highlighted in bold.

	$\alpha_1$	$\alpha_2$	$\alpha_3$	$M_{\text{base}}$
$\alpha_1$	–	<b>–0.92</b>	0.29	<b>–0.47</b>
$\alpha_2$	<b>–0.92</b>	–	<b>–0.46</b>	0.29
$\alpha_3$	0.29	<b>–0.46</b>	–	<b>–0.43</b>
$M_{\text{base}}$	<b>–0.47</b>	0.29	<b>–0.43</b>	–

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Fig. 1.** Diagrams showing the site location (inset), terminus positions and geomorphology plotted on a hillshaded mosaic of a stereophotogrammetrically derived digital elevation model (DEM) from images acquired in 1985, and ASTER GDEM (Hvidegaard et al., 2012). **(A)** Termini and geomorphology for 1859–2012, with ASM limits delineated in yellow, and **(B)** a detailed view of termini for the period 1948–2012.

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Fig. 2.** Fraction of annual runoff occurring for each month as given by MAR and RACMO2 SMB models for KNS and AS between 1960–2012 (Van As et al., 2014). Error bars are given to 2 standard deviations.

## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.



**Fig. 3.** Reconstructed photographer position showing **(A)** the area that would be observable in the photograph shown in **(B)** that was acquired in 1903.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 4.** (A) Terminus change relative to the 2012 terminus position. Uncertainty in terminus position for 1859–1903 highlighted in grey, with a range of potential advance rates for 1903–1920 indicated. These range from a minimum of no change ( $0 \text{ m a}^{-1}$ ) to a maximum possible advance rate of  $191 \text{ m a}^{-1}$ . (B) Annually averaged rates of terminus change between observations (black dots). Includes terminus advance rates described for 1903–1921 terminus change indicated on A. (C) Summer ATA (June, July, August) at annual resolution (white bars), and red line showing the averaged ATA between terminus observations (Cappelen et al., 2012; Vinther et al., 2006). (D) Annual SSTA for the area  $61^\circ$  to  $65^\circ \text{ N}$   $51^\circ$  to  $56^\circ \text{ W}$  at annual resolution (white bars) and red line showing the averaged SSTA between terminus observations (Rayner et al., 2003).



**Fig. 5.** (A) Evolution of terminus position for model runs (coloured lines) determined to be successful according to the criteria outlined in the text, with observed terminus position also plotted (bold black line, with positions between observations linearly interpolated). (B) Combined KNS and AS runoff volume estimates for 1871–2012 that are used to drive the model (5 year moving average also plotted in red). (C) Absolute annual SST estimates used to drive the model from Rayner et al. (2003) for the area 61° to 65° N and 51° to 56° W (5 year moving average also plotted in red).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Fluctuations of a Greenlandic tidewater glacier

J. M. Lea et al.



**Fig. 6.** The distribution of the tuning parameters **(a)**  $\alpha_1$  (bin width = 0.2), **(b)**  $\alpha_2$  (bin width = 0.025), **(c)**  $\alpha_3$  (bin width = 0.025), and **(d)**  $M_{\text{base}}$  (bin width =  $0.05 \text{ km}^3 \text{ a}^{-1}$ ) for successful runs as defined by the criteria outlined in the text. Minimum and maximum x-axis values represent the full range of values tested within the 1500 model runs.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
