

**Interactive comment on “Evolution of surface velocities and ice discharge of Larsen B outlet glaciers from 1995 to 2013” by J. Wuite et al.**

Reply to T.A. Scambos (Referee)

**COMMENT:** *Review of Wuite et al, The Cryosphere Disc. The paper describes a series of ice velocity mappings of the Larsen B tributary glaciers, and flux gate estimates of their outflow for 1995 and a series of measurements since then, mainly post-2002. The authors conclude that all the glaciers are moving much faster than their 1995 rates, and that wide-embayment glaciers (e.g., Hektor-Green-Evans) have had a series of accelerations and partial decelerations.*

*This is a very good observational study – well presented, well referenced, and well written. It deserves to be published. There are really no major weaknesses here. However, interpretation of the results is somewhat cursory. I assume that with this manuscript out, future papers will be able to use the data presented here to understand the system and explain it, or model it, better.*

**REPLY:** We thank the reviewer for his comments and suggestions, below you find our response to the review. We hope this and the adjustments in the text clarify the manuscript. The velocity products generated during this study will be made available soon for the wider scientific community through our project website at: <http://glacapi.enveo.at/>

**COMMENT:** *The statement on 6278 L13-L18 is significant, but not supported, not that I can see – If there is truly evidence of summer seasonal acceleration, it should be highlighted with a clearer figure, and if a case can be made for sea ice backstress, it should be shown, or the statement should be retracted. This potential for seasonal variability has been talked about quite a bit. It would be plausible because of the similarity in climate to areas of the Greenland coast. There has been speculation about either summer melt percolation or fast ice back-stress, but no clear evidence that I’m aware of. If you have it, that would be a nice addition to the paper.*

**REPLY:** Based on the comment we checked again the velocity time series. The velocity variations for the glaciers in the study area are clearly dominated by multi-annual trends triggered by ice shelf disintegration of the northern and central sections of Larsen-B, respectively the weakening of SCAR Inlet ice shelf. Based on a time series of 22 TerraSAR-X image pairs we observe a signal of seasonal acceleration by several per cent on Crane and Jorum glaciers, but not in every year. Compared to the longer-trend this signal is rather modest. We revised the text accordingly.

**COMMENT:** *The data for Flask, Leppard, Starbuck, and the Scar Inlet shelf area is interesting, and clearly shows a system in slow-motion transition, adjusting to the loss of the main Larsen B backstress. The development of sharper shear margins, and the tension cracks on the eastern side of Scar Inlet, suggest that no further change in climate or ocean conditions may be needed for this area to rapidly calve away the current shelf and initiate the same kind of rapid acceleration and thinning seen for, e.g., Crane and Jorum post-2002. This should be stated more broadly in the conclusions (the current statement is one sentence).*

**REPLY:** Thanks for pointing out the interest in the observations of the SCAR Inlet area and the suggestions. We strengthened these issues in the discussion and conclusion sections.

**COMMENT:** *Error bars should be shown for the different missions, especially for Figures 6 and 7, where they would be more obvious (and some note on Figure 3 that they are comparable or smaller than the line thickness). Rather than clutter up these nice clear graphics, I think a set of example errors for the different velocity mappings, next to the color / mapping legends, would be adequate.*

**REPLY:** The error bars have been added to the figures.

**COMMENT:** *Similarly, Table 2 shows clearly that errors are large enough that reporting mass flux to 0.001 Gt/a is unnecessary, and in fact nearest 0.1 to 0.05 Gt/a is all that is justified.*

**REPLY:** We agree, Tables 2 & 3 are adjusted in the revised manuscript.

**COMMENT:** *I would like to see a figure similar to Fig 3 and Fig 6 showing Mapple, Pequod, and Melville Glaciers, and perhaps Punchbowl and Starbuck. I'm quite surprised at the rather large velocity increase reported for MMP. Elevation decrease was relatively minor for these glaciers in the 2000's.*

**REPLY:** The velocities and the mass turnover of these glaciers are rather small. Acceleration is confined to the lowest few kilometres of the terminus. This explains why the increase of ice export and the resulting mass deficit after ice shelf collapse have been rather small. We provided additional details on the velocities of these glaciers in the text, and updated figure 3 with additional velocity profiles for Punchbowl and Melville. The acceleration was highest at Melville Glacier whereas Starbuck Glacier did not accelerate (see velocities in Fig. 7).

**COMMENT:** *Have a look at compilations of the marine bathymetry published in Lavoie et al (The Cryosphere Discussions, discussion closed) – this may help extend the kinds of observations / speculations made regarding Crane Glacier to others in this study.*

**REPLY:** Thanks for this suggestion. The information in this publication on the location of troughs in the Larsen B embayment supports our conclusion on small mass turnover of MMP and Starbuck glaciers (no deep troughs) versus deep troughs in front for Crane, HG, and centre of Scar inlet IS (downstream of Flask and Leppard glaciers).

**COMMENT:** *6272 L17 – change to . . . their discharge was 38% and 45% respectively, higher than in 1995.*

**REPLY:** Changed

**COMMENT:** *6279 L8 – change 'since' to 'for' — for an American or British ear, at least.*

**REPLY:** Changed

**COMMENT:** *6287 L9-10 'intermitted is awkward to a U.S. ear (eye).*

**REPLY:** Changed to “alternated with”

**Interactive comment on “Evolution of surface velocities and ice discharge of Larsen B outlet glaciers from 1995 to 2013” by J. Wuite et al.**

Response to Anonymous Referee #2

**COMMENT:** This study presents a very thorough and probably the most careful and complete analysis of variations in ice discharge of outlet glaciers into the former and remnant parts of the Larsen B ice shelf so far. Based on satellite measurements of ice dynamics over various time periods and measurements or estimations of ice thickness at flux gates it is a significant and important addition to previous studies which have been primarily or solely based on change in surface elevation. Elevation change methods provide information of total ice mass change, whereas the budget (or input/output) method like the one presented allows much better insight into underlying processes. Although the information on surface mass balance in this area is very limited, I see the outcome of this observational study as an important contribution for a better understanding of changes in ice dynamics during and post ice shelf collapse. The results are clearly summarized and presented in the tables and are likely to find uptake in future studies. The paper is well structured and written. I have three major comments, and several minor comments about the analysis, description of methods, wording, and figures, but recommend full publication once this is considered.

**REPLY:** We thank the reviewer for his/her comments and suggestions, below you find our response to the review. We hope this and the adjustments in the text clarify the manuscript. The velocity products generated during this study will be made available soon for the wider scientific community through our project website at: <http://glacapi.enveo.at/>

**COMMENT:** Major comments: 1/ Baseline for the surface velocity fields are one day repeat pass interferograms in 1995 and 1999. There are meanwhile quite a few examples and theoretical studies that velocities over such short repeat pass intervals are not representative for mean velocities. E.g. Marsh et al. (2013) report that tides produce horizontal velocity variations of > 50% around the mean velocity near the grounding line at the Beardmore Glacier, which are still around 5% about 15 km upstream. However, daily fluctuations rapidly smooth out over time. It seems that the authors did either ignore or did not observe such fluctuations at the Larsen B ice shelf. In any case it needs either to be mentioned (that such fluctuations are ruled out) or at least taken into account in larger errors bars for the ice discharge in 1995 and 1999. The comment on p 6279 (line 25-27) is insufficient, as it is about the uncertainties related to vertical tidal displacements in the interferograms rather than horizontal velocity fluctuations. This could be mentioned in various places, like in section 3.1/3.2, the discussion, or in the introduction.

**REPLY:** Regarding the 1995 to 1999 velocities on the Larsen-B Ice Shelf and its tributary glaciers, comprehensive work has been performed before at Univ. of Innsbruck, reported in the Ph. D. thesis of W. Rack (Rack, 2000) and other publications (Rack et al., 2000; Rack and Rott, 2004). Therefore emphasis of this paper is on ice flow behaviour of glaciers after Larsen B collapse. The question of possible temporal variations of velocities is certainly important. Therefore we checked again the

surface motions at the flux gates, using the following ERS interferograms, acquired from the same view direction (descending orbit) with excellent signal (high coherence): 15/16 Oct. 1995, 31 Oct/1 Nov 1995, 9/10 Nov 1999. The agreement of the motion related fringes in these interferograms at the flux gates is remarkable, in spite of the fact that the tidal deformation further downstream, at the transition zone between the ice shelf and glaciers, is quite different on the three dates. The maximum difference in velocity between the 3 dates at any of the gates is  $< 5\%$ . On the other hand, the Larsen B Ice Shelf has been subject to gradual acceleration during the pre-collapse period, with the acceleration increasing towards the front of the ice shelf (see references above, cited also in the manuscript). The InSAR observations indicate stable conditions of the outlet glaciers, at least until 1999. It should be mentioned that the gates for retrieving the 1995-1999 ice fluxes correspond to the 2008/09 flux gates, and therefore were located in 1995-1999 several kilometres inland of the upper limit of the tidal deformation zone. We explained this now in the text.

Besides, we observe good agreement between GPS velocity measurements at two stations of British Antarctic Survey on Flask Glacier, 12 km and 16 km above the grounding, and our velocity analysis with TerraSAR-X. Velocities of 0.95 m/d, respectively 0.71 m/d, were measured by GPS at the two stations over an annual interval (9 Nov 2009 to 5 Nov 2010). The TerraSAR-X velocities at these stations, derived from an image pairspanning 2 July to 13 July 2010, are 0.91 m/d and 0.71 m/d. Because of the different time periods the GPS data cannot be used for direct validation of the satellite measurements. However, this agreement is another indication for rather stable velocities, as observed by TerraSAR-X during the 2009 to 2013 time period on Flask Glacier.

The retrieval of 2D velocities on grounded ice uses points on ice free surfaces near the glacier margins as reference for zero velocity. On floating ice points without horizontal motion are used, so that the observed reference signal corresponds to the tidal displacement. In our analysis for Larsen-B ice motion the reference points are located at the margins of the Seal Nunataks and at several points in small coves along the ice shelf margin of Jason Peninsula. The agreement of the tidal deformation at these points confirms that the vertical displacement is representative for the ice shelf. Uncertainty in tidal deformation plays mainly a role in the velocity retrievals from 1-day repeat pass ERS InSAR data. With the estimated uncertainty in tidal deformation at the reference points of 0.5 fringes (corresponding to 3.6 cm projected onto a horizontal surface) and an uncertainty of 0.2 fringes for a velocity point, the resulting total uncertainty for ERS InSAR derived horizontal motion is 3.9 cm. For grounded ice the estimated uncertainties are 0.2 fringes for the reference points and 0.2 fringes for the velocity point, resulting in total uncertainty of horizontal displacement of 2.0 cm.

**COMMENT:** 2/ As the authors describe, there has been a significant change in surface topography post ice shelf disintegration, and the DEM used in the interferometric analysis is therefore not representative. What is the introduced error in the InSAR analysis?

**REPLY:** The geocoding of the 2D velocity vector (retrieved in radar geometry) is performed for each displacement value independently. We use the 100 m Digital Elevation Model derived from the ASTER GDEM and available through the NSIDC database (Cook et al, 2012a), which is the most detailed available DEM for the total study area. This DEM is compiled from ASTER scenes from a

range of dates between 2000 and 2009 which are unspecified in the final product (Cook et al, 2012b). Part of this period is pre-collapse and part is post-collapse. In general absolute errors in the DEM lead to very small errors in velocity. In order to estimate the impact of a change in surface slope we estimate the introduced error for a hypothetical test case with 100 m surface lowering over 15 km, which is comparable to the amount of surface lowering observed on the lower terminus of large outlet glaciers. Even in this case the induced error in velocity is well below 1%.

*Cook, A. J., T. Murray, A. Luckman, D. G. Vaughan, and N. E. Barrand.: A new 100-m digital elevation model of the Antarctic Peninsula derived from ASTER Global DEM: Methods and accuracy assessment, Earth System Science Data Discussions, 5 (1), 365–403, doi:10.5194/essdd-5-365-2012, 2012a.*

*Cook, A. J., T. Murray, A. Luckman, D. G. Vaughan, and N. E. Barrand.: A new 100-m digital elevation model of the Antarctic Peninsula derived from ASTER Global DEM: Methods and accuracy assessment, Earth Syst. Sci. Data, 4, 129-142, doi:10.5194/essd-4-129-2012, 2012b.*

**COMMENT:** 3/ It would be good to show the velocity differences as a figure (the difference between Figure 2 left and right) to illustrate how far upglacier velocity acceleration was detected.

**REPLY:** We have added a third panel to Figure 2 showing the increase in velocities between the two epochs in percent. The figure illustrates the extent of the velocity changes due to the collapse. We also added extra info to the text.

**COMMENT:** Minor comments: 6272, Abstract, SCAR Inlet: I suggest capital letters for SCAR (often neglected in the literature, but it is an acronym for the Scientific Committee for Antarctic research, it has nothing to do with a scar) like in the heading for 3.2, but be consistent throughout the manuscript.

**REPLY:** All occurrences have been capitalized in the updated manuscript.

**COMMENT:** Change wording in last sentence, use e.g.: In 2013 their discharge was 38% (Flask Gl.) and 45 % (Leppard Gl.) higher than in 1995.

**REPLY:** Sentence is adjusted

**COMMENT:** 6272, 26: remove ‘calving’

**REPLY:** Deleted

**COMMENT:** 6273, 4/5: change wording, maybe: . . . and its interaction with grounded ice.

**REPLY:** Sentence is adjusted

**COMMENT:** 6: 'Larsen Ice Shelf' as it is a geographic name change to capital letters throughout the manuscript.

**REPLY:** All occurrences have been capitalized in the updated manuscript.

**COMMENT:** 18: wording. Maybe change to: “. . . tributary glaciers continued at almost the same rate over the period. . .”

**REPLY:** Sentence is adjusted

**COMMENT:** 21: use past tense in 'inferred'.

**REPLY:** Changed

**COMMENT:** 6274, 14: . . ., defined by the ASTER. . .

**REPLY:** Changed

**COMMENT:** 16/17: . . .vectors are provided in South polar stereographic . . .

**REPLY:** We prefer to keep it Antarctic polar stereographic projection

**COMMENT:** 6275, 5: . . .from the velocity vectors . . .

**REPLY:** Changed

**COMMENT:** 17/18: . . .with a typical accuracy of 0.1 fringes . . .

**REPLY:** Changed

**COMMENT:** 19: For a one day . . .

**REPLY:** Changed

**COMMENT:** 22: . . . uncertainty for the retrieval of the displacement is in the order . . .

**REPLY:** Changed

**COMMENT:** 6276, 3 (equation (1)): instead of an integral I suggest using a summation sign (with  $i=0,N$ ) and discrete step size with (greek) delta  $y$ , as the authors were summing up a finite number of pixels across the flux gate.

**REPLY:** Changed

**COMMENT:** 8-10: based on Paterson (1994), what are the assumptions made to come up with the value 0.95?

**REPLY:**

We use parameters specified by Hulbe et al. (2008) for estimating the sliding velocity of outlet glaciers to Larsen-B. The laminar flow approximation for ice deformation is applied, with flow-law exponent  $n = 3$  and the rate factor  $A = 2.0 \times 10^{-24} \text{ Pa}^{-3} \text{ s}^{-1}$ . The deformation velocity is computed by:

$$u_{def} = \frac{2A}{n+1} (f \rho g \sin \alpha)^n h^{n+1} \quad \text{and} \quad u_{mean} = u_{surf} - u_{def} \frac{1}{n+2}$$

Where  $\alpha$  is the surface slope. The value of the shape factor  $f$  (depending on glacier width and depth) is derived for a parabolic cross section according to Patterson (1994), Table 11.3. The results show moderate values of deformation velocities for all of the glaciers. The resulting  $u_{mean}$  - values for the glaciers with accurately know cross sections are: Starbuck:  $u_{mean} = 0.95 u_{surf}$ ; Flask (1995, 1999):  $u_{mean} = 0.95 u_{surf}$ ; Flask (2009):  $u_{mean} = 0.96 u_{surf}$ ; Crane (1995, 1999):  $u_{mean} = 0.96 u_{surf}$ . For flux computations we use for Crane (1995, 1999)  $u_{mean} = 0.96 u_{surf}$ , for the other glaciers  $u_{mean} = 0.95 u_{surf}$ . For gates at calving fronts we assume full sliding.

Reference: Hulbe, C.L., Scambos, T.A., Youngberg, T., and Lamb A.K.: Patterns of glacier response to disintegration of the Larsen B ice shelf, Antarctic Peninsula, *Global Planet. Change*, 63, 1–8, 2008.

**COMMENT:** 24: clarify if the 5% uncertainty is for the satellite measured surface velocity or the vertically averaged ice velocity. It is probably OK for the measured velocity, but because of additional assumptions likely too conservative for  $U_m$  in equation (1).

**REPLY:**

See reply to comment 8-10 above. According to these results the uncertainty in deriving  $u_{mean}$  from  $u_{surf}$  is quite small. The total error for the flux estimates is dominated by the assumptions on uncertainty of the cross sections (for which we are using conservative estimates of uncertainty).

**COMMENT:** 6277, 21 (wording): . . .break up of increasingly large areas. . .

**REPLY:** Changed

**COMMENT:** 6277 (27-28)-6278 (1-2): I do not see the connection between the mass turnover and sensitivity; the cited reference (Rott et al., 2011) is also unclear about this and inadequate. If this is true, why do the authors assume that the mass turnover of e.g. Mapple Glacier is different to e.g. Punchbowl Gl.? The catchment basins look very similar in size and distance from the plateau (Fig. 1).

**REPLY:** Specific surface mass balance and mass turnover of Mapple Glacier, Punchbowl Glacier, Starbuck Glacier are similar, as indicated by slow velocities at the frontal flux gates, and reduced surface accumulation compared to the glaciers originating at the main ice divide (discussed in the paper). The name of Punchbowl Glacier slipped in by mistake into p6278, line 7 and does not correspond to the glaciers with large mass turnover (Crane, Jorum, HG). Thanks for pointing this out. Corrected now.

**COMMENT:** 6278, 3-12: why is e.g. Punchbowl so different to Mapple Gl. ? See also previous comment.

**REPLY:** See reply to 6277 (27-28)-6278 (1-2).

**COMMENT:** 13-23: in this paragraph any observations of daily variations are missing, see also major comments.

**REPLY:**Further info is provided now. See response to Reviewer No. 1.

**COMMENT:** 6279, 27: I agree to exclude the velocities at the grounding line, but also for other additional reasons; see also major comments;

**REPLY:** Ok

**COMMENT:** 6280, 24: Because of the retreat. . .

**REPLY:** Changed

**COMMENT:** 6281, 8: . . . inland of the ice front. . .

**REPLY:** Changed

**COMMENT:** 10: Change ‘For June. . .’ to e.g. ‘Based on the June 2007 analysis . . .’. This sentence is not correct, as the flux is given for a whole year.

**REPLY:** Changed

**COMMENT:** 23: . . . difference compared to 1995.

**REPLY:** Changed

**COMMENT:** 6282, 1-3: Is the value of 0.78 Gt a<sup>-1</sup> for both glaciers? Change wording (see also comment in the abstract for the usage of ‘respectively’).

**REPLY:** Changed to: “In 1995 and 1999 the velocities in the centre are 1.31 m d<sup>-1</sup> (478 m a<sup>-1</sup>) and 1.36 m d<sup>-1</sup> (496 m a<sup>-1</sup>) respectively, resulting in an ice discharge of approximately 0.78 Gt a<sup>-1</sup> for both years.”

**COMMENT:** line 26 to 6283, 1-2: how far up glacier and how fast (implied by ‘moving upward’) was the acceleration detected? This could be nicely evaluated and illustrated by a figure showing the difference in velocities. See also major comments. Reword the sentences; e.g. ‘. . . caused flow acceleration moving upstream. Our analysis of new velocity data also shows that . . .’

**REPLY:** See response to major comment 3. Rewording done.

**COMMENT:** 9: ‘. . . for six periods between . . .’

**REPLY:** Changed

**COMMENT:** 15: ‘. . . the ratio between driving stress and lateral shear. . .’

**REPLY:** Changed

**COMMENT:** 6285, 15-19: Reword and shorten this sentence. It is especially unclear what is meant by ‘. . . shear zones vs. slowly moving ice. . .’.

**REPLY:** “vs. slowly moving ice” deleted

**COMMENT:** Figures: Figure 1: change 'coastline' to 'ice edge' or 'ice front'

**REPLY:** Changed to 'ice front'

**COMMENT:** Figure 3: increase font size

**REPLY:** The font size is increased

**COMMENT:** Figure 4: use different color (preferably white) for flux gate lines;

**REPLY:** Changed to white

**COMMENT:** Figure 5: What are the yellow arrows? Increase font size. Yellow sections of ICESat track hardly visible.

**REPLY:** Figure adjusted

**COMMENT:** Figure 6: move arrow up too the curve, increase font size;

**REPLY:** Figure adjusted

**COMMENT:** Figure 7: increase font size;

**REPLY:** Font increased

**COMMENT:** References: Marsh, O.J., W. Rack, D. Floricioiu, N.R. Golledge, and W. Lawson. (2013). Tidally induced velocity variations of the Beardmore Glacier, Antarctica, and their representation in satellite measurements of ice velocity. *The Cryosphere* 7: 1375-1384. doi:10.5194/tc-7-1375-2013.

See response to main comment 1

1 **Evolution of surface velocities and ice discharge of Larsen B outlet**  
2 **glaciers from 1995 to 2013**

3

4 J. Wuite<sup>1 \*</sup>, H. Rott<sup>1 ,2</sup>, M. Hetzenecker<sup>1</sup>, D. Floricioiu<sup>3</sup>, J. De Rydt<sup>4</sup>, G. H. Gudmundsson<sup>4</sup>,  
5 T. Nagler<sup>1</sup>, M. Kern<sup>5</sup>

6

7 [1] ENVEO IT GmbH, Innsbruck, Austria

8 [2] Institute for Meteorology and Geophysics, University of Innsbruck, Austria

9 [3] Institute for Remote Sensing Technology, German Aerospace Center, Oberpfaffenhofen,  
10 Germany

11 [4] British Antarctic Survey, Cambridge, UK

12 [5] ESA-ESTEC, Noordwijk, the Netherlands

13 \* Correspondence to: Jan.Wuite@enveo.at

14

15

16 **Abstract**

17

18 We use repeat-pass SAR data to produce detailed maps of surface motion covering the glaciers  
19 draining into the former ~~Larsen B ice shelf~~Larsen B Ice Shelf, Antarctic Peninsula, for different  
20 epochs between 1995 and 2013. We combine the velocity maps with estimates of ice thickness to  
21 analyze fluctuations of ice discharge. The collapse of the central and northern sections of the ice  
22 shelf in 2002 led to a near-immediate acceleration of tributary glaciers as well as of the remnant ice  
23 shelf in SearSCAR Inlet. Velocities of most of the glaciers discharging directly into the ocean  
24 remain to date well above the velocities of the pre-collapse period. The response of individual  
25 glaciers differs and velocities show significant temporal fluctuations, implying major variations in  
26 ice discharge ~~and mass balance~~ as well. Due to reduced velocity and ice thickness the ice discharge  
27 of Crane Glacier decreased from  $5.02 \text{ Gt a}^{-1}$  in 2007 to  $1.72 \text{ Gt a}^{-1}$  in 2013, whereas Hektor and  
28 Green glaciers continue to show large temporal fluctuations in response to successive stages of  
29 frontal retreat. The velocity on SearSCAR Inlet ice shelf increased two- to three fold since 1995,  
30 with the largest increase in the first years after the break-up of the main section of Larsen B. Flask  
31 and Leppard glaciers, the largest tributaries to SearSCAR Inlet ice shelf, accelerated. In 2013 their  
32 discharge was 38 %%, respectively and 46% respectively, higher than in 1995.

33

## 34 1. Introduction

35 Atmospheric warming and changes in ocean conditions during the past decades led to wide-spread  
36 retreat of ice shelves around the Antarctic Peninsula (API) (Cook and Vaughan, 2010). Progressive  
37 retreat culminated in the final disintegration of the Larsen A ice shelf in January 1995 and of the  
38 northern and central sections of the ~~Larsen B ice shelf~~ Larsen B Ice Shelf in March 2002 (Rott et al.,  
39 1996; Rack and Rott, 2004; Glasser and Scambos, 2008). The glaciers flowing from the Antarctic  
40 Peninsula plateau, previously feeding the ice shelves, became tidewater ~~calving~~-glaciers. Most of  
41 these glaciers ~~and~~-accelerated significantly, resulting in increased ice discharge (Rott et al., 2002;  
42 De Angelis and Skvarca, 2003; Rignot et al., 2004; Scambos et al., 2004). The response of these  
43 glaciers to ice-shelf disintegration is of particular interest not only for quantifying the contributions  
44 of API outlet glaciers to sea level rise, but also for studying processes of ice shelf retreat and its  
45 interactions with grounded ice ~~masses~~ (Vieli and Payne, 2005; Hulbe et al., 2008).

46 Investigations on retreat and acceleration of glaciers in the ~~Larsen ice shelf~~ Larsen Ice Shelf region  
47 so far focused mainly on the Larsen B embayment. Rignot et al. (2004) and Scambos et al. (2004)  
48 reported on acceleration of main glaciers draining into the Larsen B embayment, based on analysis  
49 of satellite images. Rott et al. (2011) derived velocities of nine Larsen B glaciers in pre-collapse  
50 state and in 2008 and 2009 from high-resolution radar images, and estimated calving fluxes and  
51 mass balance. Estimates of the mass balance of Larsen B glaciers in recent years have been derived  
52 from changes in surface topography. Shuman et al. (2011) and Scambos et al. (2011) tracked  
53 elevation changes over the period 2001 to 2009 using optical stereo imagery and laser altimetry of  
54 ICESat and of the airborne ATM sensor. Shuman et al. (2011) reported a combined mass loss of  $-8.4$   
55  $\text{Gt a}^{-1}$  for these glaciers for the period 2001 to 2006, excluding ice lost by frontal retreat. Berthier et  
56 al. (2012) explained that the mass loss of former Larsen B tributary glaciers continued at almost the  
57 same rate was going on unabated over the period 2002 to 2011, reporting a mass loss rate of  $-9.04$   
58  $\text{Gt a}^{-1}$  for the period 2006 to 2010/2011. Scambos et al. (2014) used satellite laser altimetry and  
59 satellite stereo-imagery to map ice elevation change and ~~inferred~~infer mass changes for 33 glacier  
60 basins of the northern API over the time span 2001-2010. They report a mass balance of  $-7.9 \text{ Gt a}^{-1}$   
61 for the tributaries to the Larsen B embayment and  $-1.4 \text{ Gt a}^{-1}$  for the tributaries to the remnant ice  
62 shelf in ~~Sear~~ SCAR Inlet.

63 These reports provided estimates of mass depletion for the Larsen B tributaries integrated over  
64 multi-year periods. Here we present new analysis of satellite data showing the spatial and temporal  
65 variability in velocities over the whole Larsen B area dating back to 1995. We have included new  
66 satellite data not used in any previous studies so far, and have also reprocessed satellite radar

67 images to generate fully consistent and comparable data sets on surface velocities. Our work  
68 includes both recent acquisitions by high resolution radar sensors as well as archived data, some of  
69 which have not been exploited until now. Velocity data and estimates of ice thickness are used to  
70 derive ice discharge at different epochs, showing significant temporal variability as well. The data  
71 sets provide a comprehensive basis for studying the dynamic response of the ice masses to the  
72 disintegration of Larsen B, including the glaciers that are draining now directly into the ocean as  
73 well as the remnant ice shelf in ~~Sea~~SCAR Inlet and its tributary glaciers.

74

## 75 **2. Data and ~~Methods~~methods**

76 We derived maps of ice flow velocities from repeat-pass Synthetic Aperture Radar (SAR) data of  
77 the satellite missions ERS-1, ERS-2, Envisat, TerraSAR-X (TSX), and ALOS, applying either offset  
78 tracking or SAR interferometry (InSAR). The source data were obtained from the archives at the  
79 European Space Agency (ESA) and the German Aerospace Center (DLR). We retrieved two-  
80 dimensional surface displacement in radar geometry which we projected onto the surface, defined  
81 by the ASTER based Antarctic Peninsula DEM (API-DEM) of Cook et al. (2012), in order to  
82 produce maps of surface velocities. The resulting maps of the surface velocity vector are provided  
83 in Antarctic polar stereographic projection resampled to a 50 m grid. The DEM is compiled from  
84 ASTER scenes from a range of dates between 2000 and 2009 which are unspecified in the final  
85 product (Cook et al, 2012). During this period various glaciers have been subject to major  
86 drawdown. The sensitivity analysis on the impact of possible DEM errors shows that even in  
87 extreme cases of surface lowering the induced error in geocoded velocity is below 1%.

88 The spatial resolution of the SAR images along the flight track and in radar line-of sight (LOS)  
89 ranges from ~~1-23.3~~ 1.23.3 m x ~~3-31.2~~ 3.31.2 m for TSX ~~and to~~ 5.6 m x 9.6 m for the Advanced Synthetic  
90 Aperture Radar (ASAR) of Envisat. The time span of the repeat pass image pairs ranges from one  
91 day for ERS-1/ERS-2 tandem images to 46 days for ALOS Phased Array L-band SAR (PALSAR)  
92 images. Because of temporal decorrelation of the phase of the backscatter signal the interferometric  
93 (InSAR) method could only be applied for ERS-1/ERS-2 tandem images, available ~~for~~ on several  
94 dates of the years 1995 to 1999. InSAR data of a single swath provides the surface displacement in  
95 LOS. We combined image pairs of ascending and descending orbits ~~image pairs~~ to derive 2-D  
96 velocity fields for the period late 1995 to early 1996, ~~based on two different LOS directions from~~  
97 ~~crossing orbits~~. Being well before the collapse of the ~~Larsen B ice shelf~~ Larsen B Ice Shelf, this  
98 period is of particular importance as reference for studying the impact of ice shelf disintegration on  
99 tributary glaciers. For 1999 ERS SAR data were available only from single view direction.  
100 Assuming unaltered flow direction since 1995/1996, we derived velocity maps in November 1999,

101 using the argument from the velocity ~~maps-vectors~~ of crossing orbits.

102 For retrieving maps of ice motion from the TSX SAR, Envisat ASAR and ALOS PALSAR we  
103 apply the offset tracking technique which is based on cross-correlation of templates in SAR  
104 amplitude images. Offset tracking delivers along track and LOS velocity components from a single  
105 image pair. It is less sensitive to displacement than InSAR, but this drawback is (at least partly)  
106 compensated by the longer time span between the repeat pass images (Rott, 2009). We used  
107 templates of 64 x 64 and 96 x 96 pixels size and applied sampling steps of 10 pixels for generating  
108 velocity maps. TSX images are our main data sources for velocity maps between 2007 and 2013,  
109 complemented by occasionally available ALOS PALSAR data. Envisat ASAR data are the basis of  
110 for velocity maps for 2003 to 2006 on large glaciers and on the ~~Sea~~SCAR Inlet ice shelf.

111 The uncertainty of retrieved velocities differs between the sensors. The ERS InSAR motion maps  
112 are based on InSAR pairs of good coherence, ~~with typical accuracy of 0.1 fringes. One fringe~~  
113 (phase cycle of  $2\pi$ ), ~~corresponds to 7.2 cm along to 4 mm in LOS and 10 mm~~ projected onto ~~the a~~  
114 horizontal surface. Assuming an uncertainty of 0.2 fringes for a point on the moving glacier surface  
115 and 0.2 for the zero velocity reference points on ice free surfaces, for ERS InSAR the~~For a one day~~  
116 ~~repeat pass this corresponds to  $\pm 0.01 \text{ m d}^{-1}$~~  uncertainty in surface velocity of grounded ice is  $\pm 0.02$   
117  $\text{m d}^{-1}$ . On floating ice control points without horizontal motion are used as reference, so that the  
118 observed signal corresponds to the tidal displacement. The phase differences between individual  
119 reference points, located around the Seal Nunataks and in inlets along Jason Peninsula, are less than  
120 0.5 fringes. Assuming an uncertainty of 0.2 fringes for the moving ice shelf and of 0.5 fringes for  
121 the reference points, the uncertainty in horizontal velocity of floating ice is  $\pm 0.04 \text{ m d}^{-1}$ .

122 For offset tracking the accuracy depends on the pixel size, the time interval, and the quality of  
123 features in order to obtain good correlation peaks. We excluded areas of low correlation, so that the  
124 uncertainty for ~~the~~ retrieval of displacement is ~~in~~ the order of 0.2 to 0.3 pixels. The resulting  
125 uncertainties in the magnitude of surface motion are  $\pm 0.05 \text{ m d}^{-1}$  for TSX SAR,  $\pm 0.08 \text{ m d}^{-1}$  for  
126 ALOS PALSAR and  $\pm 0.15 \text{ m d}^{-1}$  for Envisat ASAR.

127 The mass flux across a gate of width  $Y$  [m] near the calving front or grounding line is computed  
128 according to:

$$129 \quad F_Y = \rho_{ice} \int_{y=0}^{y=Y} [u_m(y) \sin \theta H(y)] dy \quad (1)$$

130 Where  $\rho_{ice}$  is the density of ice,  $u_m$  is the vertically averaged horizontal velocity,  $\theta$  is the angle  
131 between the velocity vector and the gate, ~~and~~  $H$  is the ice thickness, ~~and is the section width~~. We  
132 use a column-averaged ice density of  $900 \text{ kg m}^{-3}$  to convert ice volume into mass. ~~The surface~~

133 ~~velocity,  $u_s$ , is obtained from satellite data.~~ For calving glaciers full sliding is assumed across calving  
134 fronts, so that  $u_m$  corresponds to the surface velocity,  $u_s$ , obtained from satellite data. For glaciers  
135 discharging into the ice shelf we estimate the ice deformation at the flux gates applying the laminar  
136 flow approximation (Paterson, 1994) using a rate factor as derived by Hulbe et al. (2008) for outlet  
137 glaciers to Larsen-B. The results show moderate values of deformation velocities. For Crane  
138 Glacier the resulting vertically averaged velocity (pre-collapse) is  $u_m = 0.96 u_s$ , for other glaciers  $u_m$   
139  $= 0.95 u_s$ . ~~assume  $u_m = 0.95 u_s$ . This is based on an estimate for ice deformation in the lower section~~  
140 ~~of main outlet glaciers applying the laminar flow approximation (Paterson, 1994).~~

141 Ice thickness at the flux gates is obtained from various sources. For Flask and Starbuck glaciers  
142 radar sounding data are available (Farinotti et al., 2013; 2014). For Crane Glacier the cross section  
143 of the calving gate is deduced from bathymetric data (Zgur et al., 2007; Rott et al. 2011). For  
144 Leppard Glacier ice thickness data of Huss and Farinotti (2014) are used. For calving fluxes of  
145 Crane, Hektoria and Green glaciers the ice thickness in the centre of the flux gate is estimated from  
146 surface height above sea level assuming flotation. The central sections of these glacier fronts have  
147 been floating at least since 2007. The surface elevation near the calving front is obtained from laser  
148 ranging data of ICESat and the Airborne Thematic Mapper (ATM) (Shuman et al., 2011; Krabill  
149 and Thomas, 2013; 2014) and in 2011 and 2013 also from digital elevation data of TanDEM-X  
150 (Krieger et al., 2013). For uncertainty estimates of mass fluxes through the gates we assume  $\pm 10\%$   
151 error of the cross section area for Starbuck, Flask and Crane glaciers, and  $\pm 20\%$  for Hektoria,  
152 Green, Jorum and Leppard glaciers. For velocity across the gate we assume  $\pm 5\%$  uncertainty.

153

### 154 3. Evolution of glacier velocities

#### 155 3.1 Velocities and frontal retreat of glaciers draining into Larsen B embayment

156 The location of the glacier basins is shown in Fig. 1, and the areas of the basins for the region  
157 upstream of the 1995 grounding line and of the 2012 glacier fronts are specified in Table 1. The  
158 basin outlines inland were provided by A. Cook based on the ASTER derived Antarctic Peninsula  
159 DEM (API-DEM) (Cook et al., 2012). The positions of the grounding lines in 1995 are from the  
160 ERS InSAR analysis of Rack (2000). The update of glacier fronts and areas in 2012 is based on a  
161 Landsat image of 12 January 2012. Before 2002 all glaciers between the Seal Nunataks and Jason  
162 Peninsula drained into ~~Larsen B ice shelf~~Larsen B Ice Shelf. Since its collapse, in March 2002, they  
163 drain into a wide bay and in the remnant part of the ice shelf in ~~Sea~~SCAR Inlet. The area of the  
164 Larsen B tributary glaciers decreased by  $270 \text{ km}^2$  since 1995. The 2012 area refers to the ice front  
165 rather than the grounding line, so that the total loss in grounded ice extent is slightly higher because

166 frontal sections of some glaciers are floating.

167 The largest glaciers north of ~~Sea~~SCAR Inlet, where the ice shelf disappeared in 2002, are Hektor-  
168 Green-Evans (HGE) and Crane glaciers. Before the ice shelf breakup the frontal zone of HGE  
169 glaciers was formed by the confluence of the three glaciers, stretching across a wide bay. Following  
170 the ice shelf collapse, these frontal regions of HGE retreated quickly (Rack and Rott, 2004;  
171 Scambos et al., 2004), suggesting that they were lightly grounded and sensitive to changes in ice-  
172 shelf buttressing. The ice shelf collapse resulted in the progressive breakup of increasingly large  
173 ~~larger and larger~~ areas of grounded ice concomitant with acceleration of ice flow and dynamic  
174 thinning, amounting to a total retreat of 174 km<sup>2</sup> by January 2012. On Crane Glacier the loss of  
175 grounded ice has been smaller (35 km<sup>2</sup>) because the terminus is confined in a narrow fjord. Jorum  
176 Glacier lost 24 km<sup>2</sup> in grounded ice, ~~and~~ Punchbowl Glacier 12 km<sup>2</sup>, and Melville Glacier 4.1 km<sup>2</sup>.  
177 The frontal positions of Mapple and Pequod glaciers have been stationary, ~~while Melville Glacier~~  
178 ~~lost 4.1 km<sup>2</sup> in area. The frontal velocity of these three glaciers increased between 1999 and 2008~~  
179 ~~by a factor of 1.8, 2.0 and 2.8 respectively, but the mass turnover is rather modest so that these~~  
180 ~~glaciers have been less affected by the ice shelf breakup (Rott et al., 2011).~~

181 An overview map of surface velocities for the Larsen B region is shown in Fig. 2a for the year 1995  
182 based on ERS InSAR data. As already reported by Rott et al. (2011), the 1995 velocities of outlet  
183 glaciers to Larsen-B agree within a few percent with the velocities retrieved from 1999 InSAR data.  
184 There is no indication for a significant temporal trend in velocity on any of the glaciers. The  
185 velocities, derived from InSAR data on various dates in 1995 and 1999 differ by less than 5 % at  
186 any of the flux gates. ~~and in~~ Varying tidal deformation along the ice shelf margins, observed in the  
187 different interferograms, did not affect the ice motion at these flux gates which are located several  
188 kilometres inland of the 1995 - 1999 grounding zone.

189 -Fig. 2b ~~as is a~~ composite of several velocity-maps from TSX and ALOS PALSAR offset tracking  
190 analysis of the years 2008 to 2012. As the figures show, a major flow acceleration is observed for  
191 HGE, Jorum, ~~Punchbowl~~, and Crane glaciers. Flask and Leppard glaciers in ~~Sea~~SCAR Inlet also  
192 accelerated, but at a lower rate. In order to investigate the temporal evolution of velocities we  
193 extracted profiles along the central flow line of the main glaciers: Hektor, Green, Jorum, ~~and~~  
194 Crane, ~~Punchbowl and Melville glaciers~~, now terminating with calving fronts (Fig. 3), and Flask  
195 and Leppard which are still confined by the remnant part of Larsen B ~~ice~~ shelf. The location of  
196 the profiles is charted in Fig. 1. The map of velocity changes (Fig. 2c) and the longitudinal profiles  
197 show that the flow acceleration extends far upstream on the large glaciers, whereas on the smaller  
198 glaciers the acceleration has been modest and confined to the lower part of the tongues.

199 The velocity of Hektor and Green glaciers is presently still much higher than in 1995, but has

200 | been subject to strong variations since 2002 associated with glacier thinning and frontal retreat. The  
201 | velocity profiles (Fig. 3) show periods of acceleration followed by gradual deceleration.  
202 | ~~Superimposed on this general trend there is a smaller annual variation in velocities with higher~~  
203 | ~~velocities in summer. These annual variations in flow may possibly be related to changes in sea ice~~  
204 | ~~cover and associated buttressing.~~ In 2008 Hektoria and Green glaciers still had a common terminal  
205 | section, but the lower terminus was already heavily fractured (Fig. 4). In 2009 a major section along  
206 | the front broke away leading to another rise in velocities. In November 2009 the frontal velocity of  
207 | Hektoria and Green glaciers was about twice the velocity at the same point in October 2008. The  
208 | high velocities persisted until March 2012, after which significant slow-down and an interim  
209 | advance of the floating tongues was observed in 2013.

210 | Ice flow and calving fluxes of Crane and Jorum glaciers have been investigated by Rott et al. (2011)  
211 | based on ERS InSAR data of 1995 and 1999 and TerraSAR-X data of several dates between  
212 | October 2008 and November 2009. During 2008/09 the velocity was comparatively stable on both  
213 | glaciers (Rott et al., 2011). Our analysis of the extended TSX data set shows a strong deceleration  
214 | since 2007. The velocity in the centre of the flux gate 1 km upstream of the 2008 glacier front  
215 | decreased from  $6.8 \text{ m d}^{-1}$  in June 2007 to  $5.2 \text{ m d}^{-1}$  in November 2008 and October 2009, and to  $2.9$   
216 |  $\text{m d}^{-1}$  in November 2013. Between 2003 and 2007 the strong acceleration of Crane Glacier caused  
217 | dynamic thinning and subsidence on the order of 150 m on the lower terminus (Scambos et al.,  
218 | 2011). In spite of continued thinning, although with reduced rate, the position of the glacier front  
219 | has been rather stable since 2006. The shape of the glacier bedrock in form of a deep canyon,  
220 | inferred from bathymetric data, indicates that the central part of the lower terminus has been  
221 | ungrounded ~~since for~~ several years (Rott et al., 2011). This suggests that lateral drag plays a key  
222 | role in maintaining the frontal position since 2006. Also the velocity of the Jorum Glacier terminus  
223 | is still higher than before ice shelf collapse. As on Crane Glacier, the velocity decreased since 2007,  
224 | but at smaller percentage.

225 | The velocities of the glaciers that are originating east of the main ice divide are small, and the flow  
226 | acceleration has been modest. On Punchbowl Glacier the velocity at the central flow line near the  
227 | calving gate increased from  $0.20 \text{ m d}^{-1}$  in 1995 to  $0.50 \text{ m d}^{-1}$  in 2008 to 2012, on Melville Glacier  
228 | from  $0.25 \text{ m d}^{-1}$  in 1995 to  $0.40 \text{ m d}^{-1}$  in 2008 and  $0.70 \text{ m d}^{-1}$  in 2012 (Fig. 3). Whereas Punchbowl  
229 | and Melville Glaciers have been subject to frontal retreat, the frontal positions of Mapple and  
230 | Pequod glaciers have been stable. This is reflected in the observed velocities near the front. On both  
231 | glaciers a temporary acceleration is observed in 2007: on Pequod Glacier from  $0.29 \text{ m d}^{-1}$  in 1995 to  
232 |  $0.40 \text{ m d}^{-1}$  in 2007; on Mapple Glacier from  $0.16 \text{ m d}^{-1}$  in 1995 to  $0.21 \text{ m d}^{-1}$  in 2007. During the  
233 | period 2008 to 2012 the velocities returned to the pre-collapse values.

234 [The velocity variations of the outlet glaciers are clearly dominated by multi-annual trends triggered](#)  
235 [by ice shelf disintegration. On some of the glaciers seasonal variations in velocity by a few per cent](#)  
236 [are observed, but not in every year. Compared to the long term trend this signal is not significant.](#)

### 237 **3.2 Velocities of SCAR Inlet ice shelf and tributary glaciers**

238 The area of the ice shelf in ~~Sear~~SCAR Inlet decreased from 3463 km<sup>2</sup> on 18 March 2002 (Rack and  
239 Rott, 2004) to 1870 km<sup>2</sup> in January 2012. The velocities on the ice shelf section which is nourished  
240 by Flask and Leppard glaciers increased two- to three-fold since 1995/1999. This section is  
241 separated by distinct shear zones from the ice shelf sections along Jason Peninsula and the section  
242 downstream of Starbuck and Stubb glaciers (Fig. 5). Major rifts are apparent on the ice shelf in the  
243 ASAR image of 28 January 2004, indicating that the disintegration of the ~~central and northern main~~  
244 ~~sections of Larsen B ice shelf~~Larsen B Ice Shelf affected the stability of the remnant ice shelf  
245 ~~section~~ in ~~Sear~~SCAR Inlet rather soon. In June 2004 the velocities along the central flowlines  
246 downstream of Flask and Leppard glaciers had already doubled compared to the pre-collapse values  
247 (Fig. 6), another indication that the Larsen B disintegration event had a rather immediate impact on  
248 the stress field of ~~Sear~~SCAR Inlet ice shelf. In the profiles of 1995 and 1999, based on one-day  
249 InSAR repeat pass data, we exclude the tidal deformation zone because of ambiguity between  
250 horizontal motion and vertical displacement.

251 In spite of still being backed up by an ice shelf, both Flask and Leppard glaciers accelerated since  
252 1995/1999 (Fig. 6). Between 1995 and 1999 there are no apparent differences in velocity. On Flask  
253 Glacier the mean velocity in 2009 to 2013 at the flux gate, 6 km above the grounding line, is  
254 ~~3841~~ % higher than the velocity in 1995/1999. On Leppard Glacier the velocity at the flux gate, 4  
255 km above the grounding line, increased by 45 %. The signal of acceleration between the two  
256 periods extends more than 30 km up-glacier, with the velocity change decreasing with distance from  
257 the grounding zone. The acceleration of the glaciers is in line with substantial acceleration of  
258 ~~Sear~~SCAR Inlet ice shelf since 2002. The main speed-up happened before 2009. Between  
259 September 2009 and July 2013 the velocities have been rather stable. The smaller Rachel, Starbuck  
260 and Stubb glaciers do not show any significant change in velocities since 1995.

261

### 262 **4. Temporal variations of ice discharge**

263 Estimates of ice discharge of Crane, Jorum, Hektor, and Green glaciers in different years are  
264 presented in Table 2. The estimated discharge of Hektor and Green glaciers for 1995 amounts to  
265 1.19 Gt a<sup>-1</sup> using the same gate near the 2008 front as Rott et al. (2011) (Fig. 4). By February/March  
266 2004, two years after the collapse, the maximum velocity at this gate was 5.1 m d<sup>-1</sup> (1862 m a<sup>-1</sup>),

267 five times higher than in 1995. A transect on Hektoria Glacier, acquired by [the NASA ATM](#) in 2004  
268 (Krabill and Thomas, 2013), allows for an estimate of an ice thickness of 406 m under the  
269 assumption of flotation, resulting in a flux of  $4.74 \text{ Gt a}^{-1}$ . The estimate for 2008 by Rott et al. (2011)  
270 amounts to  $2.88 \text{ Gt a}^{-1}$ . At that time the two glaciers still formed a single calving front. The  
271 maximum velocity at the front was  $4.23 \text{ m d}^{-1}$  ( $1545 \text{ m a}^{-1}$ ) and the maximum ice thickness ~~of at~~ the  
272 (floating) calving gate, inferred from an ICESat profile, is estimated at 268 m. Because of [the](#)  
273 retreat of the terminus by 4 km between 2008 and 2011, the gates, ~~used here~~ for [the](#) 2010 and 2013  
274 fluxes, are shifted inland (Fig. 4). A transect of surface elevation on Hektoria Glacier was measured  
275 in 2011 by [the](#) ATM during the IceBridge campaign. The freeboard at the gate is 55 m, resulting in a  
276 maximum ice thickness of 450 m assuming freely floating ice. The corresponding numbers for the  
277 calving fluxes, with November 2010 velocities, are  $1.67 \text{ Gt a}^{-1}$  for Hektoria Glacier and  $1.99 \text{ Gt a}^{-1}$   
278 for Green Glacier, adding up to  $3.66 \text{ Gt a}^{-1}$  which is 27 % higher than the flux in 2008. By July  
279 2013 the combined flux decreased to  $3.05 \text{ Gt a}^{-1}$ . This illustrates the impact of velocity variations on  
280 calving fluxes, resulting in major fluctuations of glacier net mass balance within a few years.

281 For computing the ice flux for Crane Glacier, the same flux gate 1 km inland of [the ice](#) front  
282 ~~location~~ in 2008 and 2009 is used as by Rott et al. (2011) (Fig. 4). Because of slow down of ice  
283 flow (Fig. 3) and reduction in ice thickness, the calving flux of Crane Glacier decreased  
284 significantly during recent years. [Based on the June 2007 analysis](#) ~~For June 2007~~ the flux across the  
285 gate is estimated at  $5.02 \text{ Gt a}^{-1}$ , 4.4 times higher than the pre collapse calving flux of  $1.15 \text{ Gt a}^{-1}$ .  
286 Until November 2013 it decreases to  $1.72 \text{ Gt a}^{-1}$ , one third of the 2007 flux.

287 In 1995 the combined mass flux of Jorum Glacier across the 2008 calving gates of the two glacier  
288 branches amounted to  $0.35 \text{ Gt a}^{-1}$  (Rott et al., 2011). ~~In~~ For 2008 [the](#) elevation data from an ICESat  
289 transect close to the gates were used to estimate the maximum ice thickness. For estimating the ice  
290 thickness in 2012 we use surface elevation data of the TanDEM-X satellite mission, which show  
291 surface lowering by a few metres since 2008. The estimated calving flux for the two branches of  
292 Jorum Glacier decreased from  $0.61 \text{ Gt a}^{-1}$  in 2008 to  $0.45 \text{ Gt a}^{-1}$  in 2013.

293 For Starbuck, Flask and Leppard glaciers data on ice thickness are available from ice sounding  
294 measurements and ice flow modelling (Farinotti et al., 2013; 2014; Huss and Farinotti, 2014). On  
295 Starbuck Glacier the TSX ice motion data of 2009 and 2011 do not reveal any significant difference  
296 ~~versus compared to~~ 1995 (Fig. 7). Therefore the discharge has likely not changed significantly  
297 either. The flux through the cross section near the grounding line, with maximum velocity of  $0.34 \text{ m}$   
298  $\text{d}^{-1}$  ( $124 \text{ m a}^{-1}$ ), ~~yields a flux of~~ is estimated at  $0.67 \text{ Gt a}^{-1}$  (Table 3). The ice on Flask and Leppard  
299 glaciers is thicker and velocities are higher. For Flask Glacier the mass flux is derived for a gate  
300 along a transverse profile 4 km above the grounding line (Fig. 7). This corresponds to the position

301 of radio echo sounding profile 1, acquired by the BAS Polarimetric Airborne Survey Instrument in  
302 November 2011 (Farinotti et al., 2013). In the centre of the profile the ice thickness is 690 m. In  
303 1995 and 1999 the velocities in the centre are 1.31 m d<sup>-1</sup> (478 m a<sup>-1</sup>) and, respectively 1.36 m d<sup>-1</sup>  
304 (496 m a<sup>-1</sup>), respectively, the resulting in an ice discharge of across the gates is approximately 0.78  
305 Gt a<sup>-1</sup> for both years and 0.80 Gt a<sup>-1</sup> for the two years. On Flask Glacier the V velocities in 2009 to  
306 2013 range from 1.76 m d<sup>-1</sup> (642 m a<sup>-1</sup>) to 1.93 m d<sup>-1</sup> (704 m a<sup>-1</sup>), and the ice discharge ranges from  
307 1.08 Gt a<sup>-1</sup> to 1.23 Gt a<sup>-1</sup>, without a clear temporal trend. The discharge increased between 38-36 %  
308 and 57-56 % compared to 1995 and 1999. On Leppard Glacier the centre line velocity at the gate  
309 near the grounding line has increased from 1.00 m d<sup>-1</sup> (365 m a<sup>-1</sup>) in 1995 to 1.44 m d<sup>-1</sup> (526 m a<sup>-1</sup>)  
310 in 2009 and 1.48 m d<sup>-1</sup> (541 m a<sup>-1</sup>) in 2013. The flux increased from 1.22 Gt a<sup>-1</sup> in 1995 to 1.74 Gt a<sup>-1</sup>  
311 in 2011 and 1.78 Gt a<sup>-1</sup> in 2013, an increase of 43 % and 46 %, respectively. As for Flask Glacier,  
312 there is no significant difference between 2009 and 2013.

313 The flow acceleration and increased ice discharge results in dynamic thinning which is confirmed  
314 by ICESat laser altimeter measurements. For analysis of elevation change we selected dates with  
315 closely spaced ICESat repeat tracks: track 129 of 1 June 2004 and 27 November 2008 shifted by 31  
316 m on Leppard Glacier and 28 m on Flask Glacier; track 390 of 18 June 2004 and 19 March 2008  
317 shifted by 71 m on Leppard Glacier. We corrected for the shift by taking into account the surface  
318 slope derived from the API-DEM. For track No. 129, crossing Flask Glacier 0.5 km downstream  
319 and Leppard Glacier 5 km upstream of the Flux-flux gate, we obtain a mean annual rate of surface  
320 elevation change of -1.93 m a<sup>-1</sup> on Leppard Glacier and of -2.22 m a<sup>-1</sup> on Flask Glacier. For track  
321 No. 390 we obtain for a profile across Leppard Glacier 13 km upstream of the flux gate a mean  
322 annual rate of elevation change of -1.71 m a<sup>-1</sup>.

323

## 324 5. Discussion

325 In line with previous studies, our data shows a drastic increase in flow velocities of major tributary  
326 ies glaciers following the collapse of Larsen B ice shelf Larsen B Ice Shelf in early 2002. Reduced  
327 backstress and frontal retreat caused the signal of flow acceleration that moved propagated up-  
328 glacier. Beyond that, our analysis of new velocity data shows that some of the glaciers slowed down  
329 significantly during recent years. Strong acceleration and increase of calving flux is observed for  
330 HGE glaciers and Crane Glacier, downstream of which the seafloor map shows deep troughs  
331 (Lavoie et al., 2015).

332 Scambos et al. (2004) present data of ice motion in 2001, 2002, and 2003 along selected points of  
333 the central flow-lines of the glaciers Crane, Jorum, Hektor, and Green, derived by feature tracking

334 in Landsat images. They report for a point near the Hektoria Glacier front a velocity increase from 1  
335 m d<sup>-1</sup> in 2001 to 5 m d<sup>-1</sup> in early 2003. Rignot et al. (2004) derived velocities up to 6 m d<sup>-1</sup> in 2003  
336 near the front of Hektoria Glacier from Radarsat images which agrees with our analysis of ASAR  
337 data of December 2003. Scambos et al. (2011) derived velocities for six ~~periods dates~~ between April  
338 2002 and December 2009 for a point 6 km upstream of the Crane Glacier front, showing a  
339 maximum velocity of 5.3 m d<sup>-1</sup> in January 2006, similar to the value of 5.5 m d<sup>-1</sup> we derived for this  
340 point from TSX data of June 2007.

341 Subsequently, our analysis shows significant deceleration for Crane Glacier since mid-2007, yet  
342 over this time period the position of the ice front has remained comparatively stable. ~~This~~  
343 ~~suggests~~ Possibly this is due to a reduction ~~that in~~ the ratio ~~between~~ driving stress ~~and~~ ~~versus~~ lateral  
344 shear ~~due to traction at the valley walls decreased~~, in accordance with decreasing surface slope on  
345 the lower glacier terminus. Targeted ice-flow modelling is required to further address this issue.  
346 From June 2007 to November 2013 the calving flux of Crane Glacier decreased from 5.02 Gt a<sup>-1</sup> to  
347 1.72 Gt a<sup>-1</sup>. Under the assumption that the pre-collapse flux corresponds to the balance flux (Rott et  
348 al., 2011), the resulting rate of mass loss decreased from 3.87 Gt a<sup>-1</sup> to 0.57 Gt a<sup>-1</sup>. Based on  
349 differencing of DEMs from optical stereo imagery in combination with ICESat data, Scambos et al.  
350 (2014) ~~et al.~~ report for HG glaciers a mean loss rate of 2.24 Gt a<sup>-1</sup> for the period March 2003 to  
351 November 2008. This is 42 % lower than our estimate for June 2007 and 35% higher than our  
352 estimate for 2008/09. These large temporal variations emphasize the importance of using common  
353 epochs when comparing glacier contributions to sea level rise obtained by different methods.

354 Whereas on Crane Glacier a period of major flow acceleration during the first five years after ice  
355 shelf disintegration was followed by a steady gradual decrease in velocity, the flow behaviour of  
356 Hektoria and Green glaciers has been more variable. Periods with increased flow velocities and  
357 frontal retreat alternated with periods of comparatively stable front positions or short-term advance.  
358 ASTER and ICESat data show substantial elevation losses on lower Green Glacier amounting to  
359 about 100 m during the time span November 2001 to late 2008 (Shuman et al., 2011). Scambos et  
360 al. (2014) report a mean loss rate of 3.84 Gt a<sup>-1</sup> for the period March 2003 to November 2008 out of  
361 which 0.53 Gt a<sup>-1</sup> are attributed to the loss of ice mass above floating for the retreating glacier area.  
362 Our estimate of the calving flux across the 2008/09 gate, located about 4 km inland of the 2004 ice  
363 front, yields 4.74 Gt a<sup>-1</sup> for March 2004 and 2.88 Gt a<sup>-1</sup> for 2008/09. With the estimated balance flux  
364 of 1.19 Gt a<sup>-1</sup> (Rott et al., 2011), the resulting net balance for the glacier area above the 2008/09  
365 gate amounts to -3.55 Gt a<sup>-1</sup> based on velocities of March 2004. For 2008 the estimated net balance  
366 is -1.69 Gt a<sup>-1</sup>. The loss rate increased in 2010/2011, and decreased in 2013, with the discharge in  
367 Table 3 referring to gates shifted inland because of frontal retreat.

368 ~~The different patterns of frontal retreat and flow behaviour depend on glacier geometry, glacier size,~~  
369 ~~topography of the glacier bed, and mass turnover.~~ Hektoria, Green and Evans glaciers, forming a  
370 joined terminus in a wide bay in 2002, have been particularly vulnerable to stress perturbation after  
371 ice shelf collapse as evident from the frontal retreat. Successive phases of transition from weakly  
372 grounded to floating ice due to flow acceleration and thinning, associated with major calving  
373 events, have been maintaining high rates of mass depletion for HGE glaciers to date. Crane and  
374 Jorum glaciers, terminating in deep and narrow fjords, have been subject to acceleration and major  
375 mass depletion during the first five years after ice shelf collapse, but slowed down afterwards.  
376 Similar behaviour after retreat into narrow fjords is observed for Sjögren-Boydell glaciers in Prince-  
377 Gustav-Channel and for Dinsmoor-Bombardier-Edgeworth glaciers in the Larsen A embayment  
378 (Rott et al., 2014). The ratio of longitudinal stress to lateral shear stress is critical for glacier motion  
379 in narrow valleys (Hulbe et al. 2008). Decreasing ice thickness and surface slope affect driving  
380 stresses and causing deceleration in flow. However, considering the ongoing thinning of the  
381 terminus and the resulting decrease of lateral shear stress, it can be concluded that Crane and Jorum  
382 glaciers will still be subject to major retreat before reaching a new equilibrium state further inland.

383 The two main glaciers draining into SearSCAR Inlet ice shelf, Flask and Leppard glaciers, have  
384 also been affected by flow acceleration in recent years. GPS measurements at stakes on Larsen B  
385 located 50 km downstream of these glaciers showed flow acceleration ~~on~~ in the order of 10 %  
386 between 1994 and 1999 (Rack, 2000). This indicates that also the southern sections of Larsen B ice  
387 shelf Larsen B Ice Shelf had weakened mechanically previous to the disintegration event in 2002, as  
388 reported for the northern and central sections (Rack et al., 2000; Rack and Rott, 2004). Our analysis  
389 of substantial flow acceleration and development of rifts, evident in satellite data of 2004, imply  
390 implies that the break-up had a near immediate impact on the stress field of SearSCAR Inlet the ice  
391 shelf. Fricker and Padman (2012) report for two crossover points on SearSCAR Inlet ice shelf  
392 relatively constant elevation change of  $\sim -0.19 \text{ m a}^{-1}$  during 1992 to 2008. Our analysis ~~on~~ of the  
393 temporal evolution of ice shelf flow ~~points out~~ suggests that changes in the rheology and stress field  
394 might not have been continuous during this period. The m Main speed-up on SearSCAR Inlet ice  
395 shelf occurred during the first two years after the disintegration of the northern and central sections  
396 of Larsen B, whereas changes later on were more gradual. According to Given the spatial pattern of  
397 acceleration, with main speed-up in the ice shelf section nourished by outflow region downstream  
398 of Flask and Leppard glaciers, increased further weakening has to be expected for the ice in along  
399 the shear zones margins of this section versus slowly moving ice, and also as well as for the ice  
400 immediately downstream of the grounding zone. Numerical models of Larsen B ice shelf Larsen B  
401 Ice Shelf in pre-collapse state show a band of weak ice along the shear zone that separates the

402 outflow of Leppard Glacier from the slowly moving ice along Jason Peninsula (Rack et al., 2000;  
403 Vieli et al., 2006). The differential acceleration of flow and the formation of additional rifts, which  
404 are evident in ASAR ~~images~~ and TSX images ~~over the years 2004 to~~ ~~since 2014~~ ~~2004~~, indicate that  
405 ice in this zone ~~has is~~ further ~~weakened~~ ~~weakenings~~ ~~since 2002~~.

406 Whereas the ice at the flux gates of Leppard and Flask glaciers accelerated ~~from 1995 to~~ 2009 by  
407 44% and 38%, respectively, the velocity of Starbuck Glacier has been stable. This can on one hand  
408 be attributed to the bedrock topography, on the other hand to the rather modest mass turnover. The  
409 lower terminus of Starbuck Glacier is firmly grounded, with a broad sub-glacial ridge in the area of  
410 the grounding zone (Farinotti et al., 2014). Under the assumption of mass balance equilibrium,  
411 supported by the observed steady ice motion since 1995, a specific surface mass balance  $b_n = 230$   
412  $\text{kg m}^{-2}$  is inferred from the ice flux across the grounding line.

413 The stable velocity in 1995 and 1999 suggests that Flask Glacier has been close to equilibrium state  
414 in those years. Thus, assuming equilibrium condition, the 1995 mass flux of  $0.78 \text{ Gt a}^{-1}$  across the  
415 flux gate results in  $b_n = 779 \text{ kg m}^{-2}$ , 3.4 times higher than the specific mass balance on Starbuck  
416 Glacier. The large difference in  $b_n$  can be explained by the strong west-east decrease of  
417 accumulation (Turner et al., 2002). Flask Glacier flows down from the main ice divide of the  
418 peninsula, whereas ~~the upper boundary of~~ Starbuck Glacier ~~is located~~ ~~originates on a small ice~~  
419 plateau 25 km to the east, ~~separated~~ ~~off from~~ the main divide ~~with by~~ the deep trough of ~~the~~ Crane  
420 Glacier ~~tongue in between~~.

421 Flask and Leppard glaciers have responded to the changing stress conditions on the ice shelf in front  
422 by acceleration. The bedrock of Flask and Leppard glaciers ~~ascends~~ towards the grounding zone  
423 from depressions several kilometres upstream (Farinotti et al., 2013; Huss and Farinotti, 2014). The  
424 ~~elevation~~ height of the glacier surface above the bedrock suggests that the glaciers are still firmly  
425 grounded above ~~the transects which we use for retrieving the mass fluxes at the flux gates~~.  
426 Consequently, ~~changes~~ in the force balance of the grounding zone probably played a main role for  
427 initializing flow acceleration.

428 Scambos et al. (2014) report rates of mass change of  $+0.12 \text{ Gt a}^{-1}$  for Flask Glacier and  $-1.31 \text{ Gt a}^{-1}$   
429 for Leppard Glacier, based on differencing of optical stereo DEMs from November 2001 to  
430 November 2006 and ICESat data from 2003 to 2008. Our analysis over recent years does not show  
431 a contrasting behaviour for the two glaciers. Under assumption that the 1995 fluxes correspond to  
432 the balance fluxes, we obtain for different dates between 2009 to 2013 mass change rates of  $-0.30$   
433  $\text{Gt a}^{-1}$  to  $-0.44 \text{ Gt a}^{-1}$  for Flask Glacier and  $-0.52 \text{ Gt a}^{-1}$  to  $-0.56 \text{ Gt a}^{-1}$  for Leppard Glacier.

434

## 435 6. Conclusions

436 The collapse of the main section of ~~Larsen B ice shelf~~Larsen B Ice Shelf in March 2002 triggered a  
437 near immediate response of ~~its-most~~ tributary glaciers with increased velocities ~~for-most-glaciers~~  
438 maintained ~~until-the-present~~to date. Acceleration of ice flow is also observed on the remnant part of  
439 the ice shelf in ~~Sear~~SCAR Inlet ~~as-well-as-on~~and its ~~major-main~~ tributaries. ~~Later-on-the~~The  
440 behaviour of the individual glaciers varies, and velocities show significant fluctuations over time.  
441 Whereas, after an initial speed up, ~~the-velocity-of-both~~ Crane and Jorum glaciers ~~decreased-slowed~~  
442 down significantly since mid-2007, the Hektoria and Green glaciers continue to show widespread  
443 fluctuations in velocity and periods of major frontal retreat ~~intermittent-alternated~~ing by with more  
444 ~~stable-stationary~~ positions or short term frontal advance. ~~This-These-different-differences-in-the~~  
445 response ~~can-partly-be-explained-by~~are related to glacier geometry and ~~the-geographic~~  
446 ~~setting~~bedrock features. Crane and Jorum glaciers retreated into deep and narrow fjords while  
447 Hektoria and Green glaciers still calve into a wide bay. Temporal fluctuations of flow velocity are a  
448 main factor for fluctuations in ice discharge, being primarily a function of ice velocity and  
449 thickness, shows a similar fluctuating pattern over time as velocities, emphasizing the importance of  
450 ~~using~~ common epochs for reconciling glacier mass balance estimates derived by different methods  
451 (Shepherd et al., 2012).

452 Because of the combined effect of ~~reduced-velocity~~slow down and decrease in ice thickness, the ice  
453 discharge of Crane ~~and Jorum-glaciers~~Glacier decreased by 66 % between 2007 and 2013 and of  
454 Jorum Glacier by 26 %. Both glaciers are expected to retreat further inland before reaching a new  
455 equilibrium in spite of slow-down, concluding from ongoing thinning and the increase of floating  
456 ice area. Hektoria and Green glaciers maintained variable ~~and-but-consistently~~ high rates of mass  
457 depletion in recent years, as the calving front ~~alternates-alternated~~ between floating and weakly  
458 grounded phases.

459 ~~The-~~increase of flow velocity on ~~Sear~~SCAR Inlet ice shelf and its larger tributaries started soon  
460 after the 2002 Larsen B collapse event, but changes have been discontinuous with most of the  
461 increase in the first years followed by comparatively small variations in velocity since 2009~~reduced~~  
462 ~~acceleration-rates~~. On the smaller tributaries changes have been ~~more~~ modest or absent. The ~~ice~~  
463 ~~flow-velocity~~ on the ice shelf section downstream of Flask and Leppard glaciers, the largest  
464 tributaries, increased two- to three fold since 1995/1999. ~~The -and-the~~ velocity at the flux gates ~~near~~  
465 ~~of these glaciers~~the hinge line of Flask and Leppard Glaciers, its largest tributaries, increased- until  
466 2009 by ~~36-37~~ % and 44%, respectively, with minor fluctuations in velocity in later years. This  
467 suggests that the SCAR Inlet ice shelf and its main tributary glaciers may have temporarily adjusted  
468 to the loss of the backstress from the main Larsen B ice shelf. However, considering the sustained  
469 high flow velocities and the enhanced formation and extension of cracks along the shear margins of

470 [the central ice shelf section, this state will not be long-lasting. These are clear signs for flow](#)  
471 [instability that will very likely lead to a complete disintegration of SCAR Inlet Ice Shelf in the near](#)  
472 [future. The fast flowing sections nourished by Flask and Leppard glaciers are separated from the](#)  
473 [stagnant sections of the ice shelf by distinct shear zones. The enhanced development and widening](#)  
474 [of rifts and the sustained high velocities are clear signs of ongoing gradual breakup that if continued](#)  
475 [will lead to a complete disintegration of Sear SCAR Inlet in the near future.](#)

476

## 477 **Acknowledgements**

478 The authors would like to thank A. Cook (Univ. Swansea, UK) for providing outlines of glacier  
479 basins. The TerraSAR-X data and TanDEM-X data were made available by DLR through project  
480 HYD1864 and XTI\_GLAC0457, respectively. The ERS SAR and Envisat ASAR data were made  
481 available by ESA through Envisat AO project ID-308. The ICESat laser altimeter data were [obtained](#)  
482 [downloaded](#) from the NASA Distributed Active Archive Center, US National Snow and Ice Data  
483 Center (NSIDC), Boulder, Colorado. The work was supported by the European Space Agency, ESA  
484 Contract No.4000105776/12/NL/CBi.

485

486

487 **References**

- 488 Berthier, E., Scambos, T.A., and Shuman, C.A.: Mass loss of Larsen B tributary glaciers (Antarctic  
489 Peninsula) unabated since 2002, *Geophys. Res. Lett.*, 39, L13501, doi:10.1029/2012GL051755, 2012.  
490
- 491 Bindschadler, R., Vornberger, P., Fleming, A., Fox, A., Mullins, J., Binnie, D., Paulsen, S. J.,  
492 Granneman, B., and Gorodetzky, D.: The Landsat Image Mosaic of Antarctica, *Rem. Sens. Env.*,  
493 112(12), 4214–4226, doi:10.1016/j.rse.2008.07.006, 2008.
- 494 Cook, A. J. and Vaughan, D. G.: Overview of areal changes of the ice shelves on the Antarctic  
495 Peninsula over the past 50 years, *The Cryosphere*, 4, 77–98, 2010.
- 496 Cook, A. J., ~~T.~~Murray, ~~T.~~A. Luckman, ~~A.~~ D. G. Vaughan, ~~D. G.~~, and ~~N. E.~~ Barrand, ~~N. E.~~: A new  
497 100-m digital elevation model of the Antarctic Peninsula derived from ASTER Global DEM:  
498 Methods and accuracy assessment, *Earth Syst. Sci. Data*, 4, 129–142, doi:10.5194/essd-4-129-  
499 2012, 2012.
- 500 De Angelis, H. and Skvarca, P.: Glacier surge after ice shelf collapse. *Science*, 299 (5612), 1560–  
501 1562. doi:10.1126/science.1077987, 2003.
- 502 Farinotti, D., Corr, H. F. J., and Gudmundsson, G. H.: The ice thickness distribution of Flask  
503 Glacier, Antarctic Peninsula, determined by combining radio-echo soundings, surface velocity data  
504 and flow modelling, *Annals of Glaciology*, 54 (63), doi:10.3189/2013AoG63A603, 2013.
- 505 Farinotti, D., King, E. C., Albrecht, A., Huss, M., and Gudmundsson, G. H.: The bedrock  
506 topography of Starbuck Glacier, Antarctic Peninsula, as measured by ground based radio-echo  
507 soundings, *Ann. Glaciol.*, 55, 22–28, 2014.
- 508 Fricker, H. A. and Padman, L.: Thirty years of elevation change on Antarctic Peninsula ice shelves  
509 from multimission satellite radar altimetry, *J. Geophys. Res.*, 117, C02026,  
510 doi:10.1029/2011JC007126, 2012.
- 511 Glasser, N. F. and Scambos, T. A.: A structural glaciological analysis of the 2002 Larsen B ice-shelf  
512 collapse, *J. Glaciol.*, 54, 3–16, 2008.
- 513 Haran, T., Bohlander, J., Scambos, T. A., Painter, T., and Fahnestock, M.: MODIS Mosaic of  
514 Antarctica 2008-2009 (MOA2009) Image Map 2009. Boulder, Colorado USA: National Snow and  
515 Ice Data Center, <http://dx.doi.org/10.7265/N5KP8037>, 2014.
- 516 Hulbe, C. L., Scambos, T. A., Youngberg, T., and Lamb A. K.: Patterns of glacier response to  
517 disintegration of the ~~Larsen B ice shelf~~ Larsen B Ice Shelf, *Antarctic Peninsula, Global Planet.*  
518 *Change*, 63, 1–8, 2008.

519 | Huss, M. and Farinotti, D.: A high-resolution bedrock map for the Antarctic Peninsula, *The*  
520 | *Cryosphere*, 8, 1261–1273, doi:10.5194/tc-8-1261-2014, 2014.

521 | Jezek, K. C., Curlander, J. C., Carsey, F., Wales, C., and Barry, R.G.: RAMP AMM-1 SAR Image  
522 | Mosaic of Antarctica. Version 2. Boulder, Colorado USA: National Snow and Ice Data Center,  
523 | 2013.

524 | Krabill, W. B. and Thomas, R.: Pre-IceBridge ATM L2 Icessn Elevation, Slope, and Roughness.  
525 | Boulder, Colorado USA: NASA DAAC at the National Snow and Ice Data Center, 2013.

526 | Krabill, W. B. and Thomas, R.: IceBridge ATM L2 Icessn Elevation, Slope, and Roughness. Version  
527 | 2. Boulder, Colorado USA: NASA DAAC at the National Snow and Ice Data Center, 2014.

528 | Krieger, G., Zink, M., Bachmann, M., Bräutigam, B., Schulze, D., Martone, M., Rizzoli, P.,  
529 | Steinbrecher, U., Anthony, J. W., De Zan, F., Hajsek, I., Papathanassiou, K., Kugler, F., Rodriguez  
530 | Cassola, M., Younis, M., Baumgartner, S., Lopez Dekker, P., Prats, P., and Moreira, A.: TanDEM-X:  
531 | a radar interferometer with two formation flying satellites, *Acta Astronaut.*, 89, 83–98,  
532 | doi:10.1016/j.actaastro.2013.03.008, 2013.

533 | [Lavoie, C., Domack, E. W., Pettit, E. C., Scambos, T. A., Larter, R. D., Schenke, H.-W., Yoo, K. C.](#)  
534 | [Gutt, J., Wellner, J., Canals, M., Anderson, J. B., and Amblas D.: Paleo-ice flow directions of the](#)  
535 | [Northern Antarctic Peninsula ice sheet at LGM based upon a new synthesis of seabed imagery. \*The\*](#)  
536 | [Cryosphere](#), 8, 613-629, 2015.

537 | Paterson, W. S. B.: The physics of glaciers, Third Edition, Oxford, etc., Elsevier, 1994.

538 | Rack, W.: Dynamic behaviour and disintegration of the northern Larsen Ice Shelf, Antarctic  
539 | Peninsula, Ph.D. thesis, University of Innsbruck, Austria, 166 pp., 2000.

540 | Rack, W., Doake, C. S. M., Rott, H., Siegel, A., and Skvarca, P.: Interferometric analysis of the  
541 | deformation pattern of the Northern ~~Larsen ice shelf~~Larsen Ice Shelf, Antarctic Peninsula,  
542 | compared to field measurements and numerical modeling, *Ann. Glaciol.*, 31, 205–210, 2000.

543 | Rack, W. and Rott, H.: Pattern of retreat and disintegration of ~~Larsen B ice shelf~~Larsen B Ice Shelf,  
544 | Antarctic Peninsula, *Ann. Glaciol.*, 39, 505-510, 2004.

545 | Rignot, E., Casassa, G., Gogineni, P., Rivera, A., and Thomas, R.: Accelerated ice discharges from  
546 | the Antarctic Peninsula following the collapse of the ~~Larsen B ice shelf~~Larsen B Ice Shelf.  
547 | *Geophys. Res. Lett.*, 31, L18401, doi:10.1029/2004GL020697, 2004.

548 | Rott, H.: Advances in interferometric synthetic aperture radar (InSAR) in earth system science,  
549 | *Progress in Phys. Geogr.*, 33(6), 769-791, doi: 10.1177/0309133309350263, 2009.

550 Rott, H., Skvarca, P., and Nagler, T.: Rapid collapse of Northern Larsen Ice Shelf, Antarctica,  
551 Science, 271, 788–792, 1996.

552 Rott, H., Rack, W., Skvarca, P., and De Angelis, H.: Northern Larsen Ice Shelf, Antarctica: Further  
553 retreat after collapse, Ann. Glaciol., 34, 277–282, 2002.

554 Rott, H., Müller, F., Nagler, T., and Floricioiu, D.: The imbalance of glaciers after disintegration of  
555 Larsen B Ice Shelf, Antarctic Peninsula, The Cryosphere, 5 (1): 125–134. doi:10.5194/tc-5-125-  
556 2011, 2011.

557 Rott, H., Floricioiu, D., Wuite, J., Scheiblauer, S., Nagler, T., and Kern, M.: Mass changes of outlet  
558 glaciers along the Nordenskjöld Coast, northern Antarctic Peninsula, based on TanDEM-X satellite  
559 measurements, Geophys. Res. Lett., 41, doi:10.1002/2014GL061613.

560 Scambos, T. A., Bohlander, J. A., Shuman, C. A., and Skvarca, P.: Glacier acceleration and thinning  
561 after ice shelf collapse in the Larsen B embayment, Antarctica, Geophys. Res. Lett., 31, L18402,  
562 doi:10.1029/2004GL020670, 2004.

563 Scambos, T. A., Berthier, E., and Shuman, C. A.: The triggering of subglacial lake drainage during  
564 rapid glacier drawdown: Crane Glacier, Antarctic Peninsula, Ann. Glaciol., 52(59), 74–82, 2011.

565 Scambos, T. A., Berthier, E., Haran, T., Shuman, C. A., Cook, A. J., Ligtenberg, S. R. M., and  
566 Bohlander, J.: Detailed ice loss pattern in the northern Antarctic Peninsula: widespread decline  
567 driven by ice front retreats, The Cryosphere, 8, 2135–2145, doi:10.5194/tc-8-2135-2014, 2014.

568 Shepherd, A., Ivins, E. R., [Geruo, A. G.](#), Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H.,  
569 Bromwich, D. H., Forsberg, R., Galin, N., Horwath, M., Jacobs, S., Joughin, I., King, M. A.,  
570 Lenaerts, J. T., Li, J., Ligtenberg, S. R., Luckman, A., Luthcke, S. B., McMillan, M., Meister, R.,  
571 Milne, G., Mouginot, J., Muir, A., Nicolas, J. P., Paden, J., Payne, A. J., Pritchard, H., Rignot, E.,  
572 Rott, H., Sørensen, L. S., Scambos, T. A., Scheuchl, B., Schrama, E. J., Smith, B., Sundal, A. V.,  
573 van Angelen, J. H., van de Berg, W. J., van den Broeke, M. R., Vaughan, D. G., Velicogna, I., Wahr,  
574 J., Whitehouse, P. L., Wingham, D. J., Yi, D., Young, D., and Zwally, H. J.: A reconciled estimate of  
575 ice-sheet mass balance, Science, 338, 1183–1189, doi:10.1126/science.1228102, 2012.

576 Shuman, C. A., Berthier, E., and Scambos, T. A.: 2001–2009 elevation and mass losses in the  
577 Larsen A and B embayments, Antarctic Peninsula, J. Glaciol., 57, 737–754, 2011.

578 Turner, J., Lachlan-Cope, T. A., Marshall, G. J., Morris, E. M., Mulvaney, R., and Winter, W.:  
579 Spatial variability of Antarctic Peninsula net surface mass balance, J. Geophys. Res., 107(D13),  
580 4173, doi:10.1029/2001JD000755, 2002.

581 [Vieli, A. and Payne, A. J.: Assessing the ability of numerical ice sheet models to simulate grounding](#)

582 | [line migration, J. Geophys. Res.-Earth, 110, F01003, doi:10.1029/2004JF000202, 2005.](#)

583 | Vieli, A., Payne, A. J., Du, Z., and Shepherd, A.: Numerical modelling and data assimilation of the

584 | ~~Larsen B ice shelf~~[Larsen B Ice Shelf](#), Antarctic Peninsula, Phil. Trans. R. Soc. A, 364, 1815–1839,

585 | doi:10.1098/rsta.2006.1800, 006, 2006.

586 | Zgur, F., Rebesco, M., Domack, E. W., Leventer, A., Brachfeld, S., and Willmott, V.: Geophysical

587 | survey of the thick, expanded sedimentary fill of the new-born Crane fjord (former Larsen B\_Ice

588 | Shelf, Antarctica): in Antarctica: A Keystone in a Changing World – Online Proceedings\_of the 10th

589 | ISAES, edited by: Cooper, A., Raymond, C., and the 10th ISAES Editorial Team,\_USGS Open-File

590 | Report 2007–1047, Extended Abstract 141, p. 4 p., 2007.

591

592 **Tables**

593 **Table 1.** Area of glacier basins (in km<sup>2</sup>) shown in Fig. 1 above the October 1995 grounding line and  
 594 updated for glacier fronts on 12 January 2012, and change of glacier area 1995 to 2012. *\_NC* – no  
 595 significant change of front position or grounding line. The areas of glacier basins include rock  
 596 outcrops and mountain slopes.

Nr.	Glacier	Area [km <sup>2</sup> ]	Area [km <sup>2</sup> ]	ΔArea
		10/1995	01/2012	1995-2012
1	Hektoria Glacier Headland	118.13	100.64	-17.49
2	HGE	1588.97	1415.05	-173.92
3	Evans Glacier Headland	124.38	119.49	-4.89
4	Punchbowl	129.56	117.82	-11.74
5	Jorum	484.09	460.36	-23.73
6	Crane	1354.29	1319.74	-34.55
7	Mapple	154.97	155.43	+0.46
8	Melville	295.26	291.18	-4.08
9	Pequod	151.06	150.64	-0.42
10	Rachel	51.27	51.27	NC
11	Starbuck	300.69	300.69	NC
12	Stubb	109.92	109.92	NC
13	Flask	1144.84	1144.84	NC
14	Leppard	1877.08	1877.08	NC
<i>Sum</i>		<i>7884.51</i>	<i>7614.15</i>	<i>-270.36</i>

597

598

599 **Table 2.** Drainage area above the flux gate, velocities at the centre of the flux gate ( $V_c$ ), discharge  
600 across the flux gates, and difference of discharge versus 1995; for glaciers draining into Larsen B  
601 embayment. \* From Rott et al. (2011).  
602

Glacier	Gate	Area km <sup>2</sup>	Date YYYY-MM	$V_c$ (m a <sup>-1</sup> )	Discharge (Gt a <sup>-1</sup> )	$\Delta$ 1995-Date 2 (Gt a <sup>-1</sup> )
Crane	C1	1235	1995/99*	548	<del>1.449</del> <u>15</u> <del>±0.128</del> <u>13</u>	
			2007-06	2464	<del>5.018</del> <u>02</u> <del>±0.561</del>	-3. <del>869</del> <u>87</u>
			2008/09*	1882	<del>2.919</del> <u>92</u> <del>±0.326</del> <u>33</u>	-1.770
			2010-04	1650	<del>2.437</del> <u>44</u> <del>±0.272</del>	-1. <del>288</del> <u>29</u>
			2011-01	1329	<del>2.209</del> <u>21</u> <del>±0.247</del> <u>25</u>	-1.060
			2012-08	1292	<del>2.145</del> <u>15</u> <del>±0.240</del>	<del>-1.00</del> <u>-0.996</u>
			2013-11	1059	<del>1.724</del> ±0.193	-0.57 <del>558</del>
Jorum	J1	382	1995/99*	475	<del>0.346</del> <u>35</u> <del>±0.071</del>	
			2008/09*	865	<del>0.534</del> ±0.110	-0.18 <del>819</del>
			2012-04	759	<del>0.388</del> <u>39</u> <del>±0.080</del>	-0.042
	J2	52	1995/99*	68	<del>0.042</del> <del>±0.009</del> <u>02</u>	
			2008/09*	146	<del>0.082</del> <del>±0.017</del> <u>02</u>	-0.040
			2012-04	153	<del>0.064</del> ±0.013	-0.022
Hektoria -Green	H1 & G1	1188	1995/99*	387	<del>1.186</del> <u>19</u> <del>±0.246</del> <u>25</u>	
			2004-03	1862	<del>4.741</del> <del>±0.978</del> <u>98</u>	-3.555
			2008/09*	1545	<del>2.878</del> <u>88</u> <del>±0.593</del>	-1.692
Hektoria	H2	341	2010-11	822	<del>1.670</del> ±0.344	
			2013-07	741	<del>1.490</del> <del>±0.307</del> <u>31</u>	
Green	G2	618	2010-11	1278	<del>1.993</del> ±0.411	
			2013-07	1095	<del>1.560</del> ±0.313	


603

604

605 **Table 3.** Drainage area above the flux gate, velocities at the centre of the flux gate ( $V_c$ ), discharge  
606 across the flux gates, and difference of discharge versus 1995; for glaciers draining into Scar SCAR  
607 Inlet ice shelf.  
608

Glacier	Area km <sup>2</sup>	Date YYYY-MM	$V_c$ (m a <sup>-1</sup> )	Discharge (Gt a <sup>-1</sup> )	$\Delta$ 1995-Date 2 (Gt a <sup>-1</sup> )
Starbuck	296	1995-10	124	<del>0.06707</del> $\pm 0.00701$	
		2009-09	125	<del>0.06907</del> $\pm 0.00801$	
		2011-01	124	<del>0.06807</del> $\pm 0.00801$	
Flask	1003	1995-10	478	<del>0.781</del> $\pm 0.08709$	
		1999-11	496	<del>0.78801</del> $\pm 0.08709$	
		2009-09	661	<del>1.111</del> $\pm 0.124$	-0.330
		2011-01	704	<del>1.22623</del> $\pm 0.13714$	-0.44545
		2012-02	690	<del>1.154</del> $\pm 0.12913$	-0.373
Leppard	1822	1995-10	365	<del>1.223</del> $\pm 0.252$	
		2009-10	526	<del>1.73974</del> $\pm 0.35836$	-0.51652
		2013-07	541	<del>1.77778</del> $\pm 0.36637$	-0.5654

609  
610  
611




613

614

615 **Fig. 1.** Overview map of glacier basins in the Larsen B region. Background: 2009 MODIS mosaic  
 616 (Haran et al., 2014). Names and size of basins in Table 1. Glacier boundaries inland are based on  
 617 the ASTER GDEM (courtesy A. Cook). Coastlines and grounding line derived from ERS-1 and  
 618 Landsat images. SN - Seal Nunataks, SI - ~~Sear~~SCAR Inlet, JP - Jason Peninsula. Red lines show  
 619 location of velocity profiles in Figs. 3 and 6.

620


621



622

623

624 **Fig. 2.** Maps of glacier surface velocity in the Larsen B region. Left: based on ERS InSAR data of  
 625 October/November 1995 (background RAMP mosaic; Jezek et al., 2013). Centre: Right: based on  
 626 TSX and PALSAR offset tracking 2008 – 2012. Right: Velocity increase 2008-2012 versus 1995.  
 627 B(background LIMA mosaic (; Bindschadler et al., 2008). The dashed line shows the position of the  
 628 1995 grounding line. Front positions are from October 1995 (left) and January 2012 (center and  
 629 right).



630

631

632 **Fig. 3.** Surface velocities along the central flow line of Hektoria, Green, Jorum, and Crane, Punchbowl and Melville glaciers and their frontal positions at different dates (month/year). Vertical  
 633 lines show positions of calving front (cf). The vertical bars show uncertainties in velocity for  
 634 TerraSAR-X (T, 2007 to 2013; PALSAR (P) Nov. 2009; ASAR (A) 2003, 2004; ERS (E) 1995,  
 635 1999.

637

639


640

641


642

643

644




**Fig. 4.** Section of TerraSAR-X amplitude images. Left and Center: HGE glaciers on 23 March 2008 and 25 March 2012, with flux gates on Hektor (H) and Green (G) glaciers and location of the glacier front on 24 December 2004 (dotted blue line) and 25 March 2012 (dotted green line). Right: TSX image of Crane (C ) and Jorum (J) glaciers on 22 April 2012 with flux gates.



645

646

647 **Fig. 5.** Envisat ASAR image of ~~Sea~~**SCAR** Inlet ice shelf and tributary glaciers, 28 January 2004.  
 648 Red lines show the flux gates for Flask (F), Leppard (L) and Starbuck (S) glaciers. The broken red  
 649 lines delimit the outflow downstream of Flask and Leppard glaciers. The yellow sections of the  
 650 ICESat tracks are used for deriving surface elevation change on Leppard and Flask glaciers. The  
 651 arrows point to major rifts.




652

653

654 **Fig. 6.** Surface velocities along the central flowline of Flask (top left) and Leppard (bottom left)  
 655 glaciers and downstream of these glaciers on **ScarSCAR** Inlet ice shelf (right, top and bottom). The  
 656 arrow shows the location of the flux gate. Position of flowline profiles [shown](#) in Fig.1 and of flux  
 657 gates in Fig. 5. [The vertical bars show uncertainties in velocity for TerraSAR-X \(T\) 2007 to 2013;](#)  
 658 [PALSAR \(P\) Nov. 2009; ASAR \(A\) 2003, 2004; ERS \(E\) 1995, 1999.](#)

659



660

661

662 **Fig. 7.** Ice thickness and surface velocity across the flux gate of Starbuck Glacier (top) and Flask  
 663 Glacier (bottom). Ice thickness from Farinotti et al. (2013; 2014).

664