

**Response to reviewer Quincey:**

Thank you to Duncan Quincey for the very positive and constructive review! We have done our best to address the comments that you have raised, and I reply to them individually below:

Voice: Sections 3 4 were re-written to include active voice rather than passive voice.

P3262 lines 5-7: The terms have been so re-ordered.

P3263 lines 8-9: The definition has been removed.

P3263 lines 20-21: An extra paragraph has been added, and we will look out for it in later typesetting, too.

P3264 line 8: 'of' changed to 'on'

P3266 line 16: based moved to after the reference

P3267 line 14: "airborne thematic mapper" is capitalized and "lidar" changed to "LiDAR"

P3271 line 6: comma changed to semi-colon

P3272 line 13: included the word "elevation" to finish the sentence

Section 5: Introductory paragraph removed, and 4 subheadings were included: Retrieval Performance Factors, Revisiting Lake Depth Retrievals, Supraglacial Lakes in the Hydrological System, and Sensitivity Analysis.

P3275 line 23: 'an' changed to 'and'

P3279 lines 1-5: This third paragraph attempts to move on from method development to applications, and as such rather than integrate it into the previous two paragraphs, it was expanded. The paragraph as been amended as follows to better address what we found out for SK region and the implications for other regions: "The recommended depth retrieval method was applied to all available Landsat 8 imagery for summer 2014 for the Sermeq Kujalleq (Jakobshavn) region of west Greenland. Seasonal and regional trends in lake depth (deepening and then shallowing), evolution (proceeding inland/up-glacier and northwards through the summer), and distribution (~300 m to ~2100 m a.s.l.) were observed. At most, lakes contain a similar magnitude of water to supraglacial streams, but this may not be true for other parts of Greenland. Both elevation (and relatedly, accumulation / melt forcing) and surface topography play a role in lake formation and extent, behavior that we expect to be modified but observable in other regions. Further work moving forward will need to contextualize Landsat data with other remote sensing imagery, fieldwork, and model outputs."

**Response to reviewer Hamilton:**

Thank you to Gordon Hamilton for providing a positive, thoughtful, and constructive review of our work. We have done our best to reply to your comments, as detailed below:

Major Comment: Thank you for this suggestion to bring more focus to the paper. While we agree that methods and applications are sometimes better divided, in our case we think that it is important to demonstrate the utility of the method and provide an example of its use. Providing an application also places the method more in discussion with the research

literature. Without this, we believe that there would be questions about efficacy of application and results. One year of data does not tell us anything about longer term changes, and so we do not think it merits a separate publication at this time. Provided Landsat 8 keeps functioning well, we would hope to return to the dataset and method in 5 years' time to look at a more meaningful time series for a separate work.

P3259 L3: Deleted.

P3259 L21: replaced "glacial" with "ice sheet's"

P3260 L4: Descriptions of positive feedbacks joined into one sentence: "supraglacial lakes contribute to multiple positive feedback processes, including ice shelf disintegration in Antarctica (Banwell et al., 2013; Glasser and Scambos, 2008) and melt-albedo interactions (Leeson et al., 2015)."

P3260 L10: Reference changed to 2011

P3260 L17: added "WorldView-3" (because WorldView-1 didn't have this capability)

P3261 L21: replaced "a supraglacial" with "the same" in order to clarify

P3262 L12: Thank you for this clarification. That sentence has been removed.

P3262 L18: Changed to "lake bottoms"

P3262 L25: Changed position of the word "parameter"

P3263 L3: Phrase deleted

P3263 L5: replaced "assumption" with "approximation for  $A_d$ "

P3263 L6: replaced "in an application of a form of" with "we use spectral mixture analysis to"

P3264 L15: The Legleiter relationship had only three field sites and tested rather than location, the setting, i.e., more testing river vs. creek vs. lake. As they only tested one lake, we make no claim here about location.

P3264 L15: deleted based

P3265 L5: Deleted "In situ concurrent and bathymetric"

P3266 L7: Shortened as suggested

P3266 L7: Written out as "four", replaced "further" with "farther." All numbers below 11 used in a non-technical context were replaced with written words.

P3266 L9: "in 2014" moved to the end of the sentence

P3266 L15: "based" moved after parentheses

P3267 L1: Sentence added: "Similar validation for ASTER has been carried out with airborne LiDAR from before lake drainage (Georgiou et al., 2009)."

P3267 L2: Written with a superscript as "meters per pixel"

P3267 L14: Changed to LiDAR

P3267 L21: This is from work in prep, and so “preliminary studies” can’t be cited – so deleted.

P3267 L26: “as” changed to “because”

P3268 L2: “errors” changed to “blunders”

P3268 L6: clarified to be in total

P3268 L10: Edited as suggested

P3268 L17: replaced “success” with “performance”

P3268 L19: added in “m” where appropriate

P3268 L21: replaced “Landsat 7” with “ETM+” for clarity

P3269 L17: Sentenced replaced as suggested

P3269 L25: “day” not “Day”

P3270 L4: replaced “returned by” with “derived with”

P3270 L5: Sentence deleted.

P3270 L7: This section includes BOTH regions. The first sentence has been amended to clarify this: “We difference all overlapping areas of DEM and Landsat-derived lake depths in both case study regions.”

P3270 L17: Replaced “very good” with “close”

P3270 L22: deleted “ice flow”

P3270 L23: Replaced the first phrase with “Despite meter scale uncertainties (1.6m) at the pixel level...”

P3271 L5: added “across the entire study area”

P3271 L5: Deleted the first part of the sentence “A histogram of all lake depths shows that”

P3272 L20: First paragraph has been removed.

P3273 L11: changed to “Indeed, other work (Moussavi et al., in review) ...”

P3273 L16: re-written as suggested

P3273 L23: removed “values”

P3274 L1: this “m” was removed by TCD. Re-added in text.

P3274 L29: Removed text regarding “significantly overestimated” – the 30% is what falls out when you follow through the overestimated g with an average reflectance estimate.

P3275 L2: replaced “believed” with “concluded”

P3275 L5: Removed final two sentences of the paragraph.

P3275 L12: Reference added.

P3275 L13: deleted as suggested

P3275 L14: moved reference to earlier section and added supplementary material to make it easier to find these values

P3275 L17: The first paragraph in this group was discussed. In order to better focus the discussion (as suggested by reviewer Quincey), four subheadings have been added: Retrieval Performance Factors, Revisiting Lake Depth Retrievals, Supraglacial Lakes in the Hydrological System, and Sensitivity Analysis.

P3276 L29: This radiative transfer modelling addresses atmospheric variability, not the lake-based HYDROLIGHT model that you used, so the two aren't directly related, if I understand correctly.

P3277 L4 changed "variability" to "varies" and added "is" later in the sentence

P3277 L6: deleted "on"

P3277 L6: Parentheticals were removed and the one run-of sentence was replaced with "Much of this error appears largely random for a given point in time and space. Thus, while it decreases confidence in individual lake depth retrievals, averaged water volume retrieval should not be biased."

P3278 L24: Deleted "in the discussion"

**Final note: not changes to figures necessary. Original figures all sufficient.**

# 1 Estimating Supraglacial Lake Depth in Western Greenland 2 Using Landsat 8 and Comparison with Other Multispectral 3 Methods

4  
5 **A. Pope**<sup>1,2,3</sup>, **T. A. Scambos**<sup>1,2</sup>, **M. Moussavi**<sup>2,4</sup>, **M. Tedesco**<sup>5</sup>, **M. Willis**<sup>6,7</sup>, **D.**  
6 **Shean**<sup>3</sup>, and **S. Grigsby**<sup>2</sup>

7 [1]{National Snow and Ice Data Center, University of Colorado, Boulder, Boulder, Colorado}

8 [2]{Cooperative Institute for Research in Earth Sciences, University of Colorado, Boulder,  
9 Boulder, Colorado}

10 [3]{Polar Science Center, Applied Physics Lab, University of Washington, Seattle,  
11 Washington}

12 [4]{Earth Science and Observation Center, University of Colorado, Boulder, Boulder,  
13 Colorado}

14 [5]{The City College, of New York, CUNY, New York City, New York}

15 [6]{Earth and Atmospheric Sciences, Cornell University, Ithaca, New York}

16 [7]{Geological Sciences, University of North Carolina, Chapel Hill, North Carolina}

17 Correspondence to: A. Pope (allen.pope@post.harvard.edu)

## 18 19 **Abstract**

20 Liquid water stored on the surface of ice sheets and glaciers impacts surface mass balance, ice  
21 dynamics and heat transport. Multispectral remote sensing can be used to detect supraglacial  
22 lakes and estimate their depth and area. In this study, we use in situ spectral and bathymetric  
23 data to assess lake depth retrieval using the recently launched Landsat 8 Operational Land  
24 Imager (OLI). We also extend our analysis to other multispectral sensors to evaluate their  
25 performance with similar methods. Digital elevation models derived from WorldView stereo  
26 imagery (pre-lake filling and post-drainage) are used to validate spectrally derived depths,  
27 combined with a lake edge determination from imagery. The optimal supraglacial lake depth  
28 retrieval is a physically based single-band model applied to two OLI bands independently (red

Allen Pope 21/10/2015 07:55

**Deleted:** Supraglacial lakes, therefore, play a significant role in ice sheet behavior as a surface expression of the glacial hydrological system.

1 and panchromatic) that are then averaged together. When OLI- and WorldView-derived  
2 depths are differenced, they yield a mean and standard deviation of  $0.0 \pm 1.6$  m. This method  
3 is then applied to OLI data for the Sermeq Kujalleq (Jakobshavn Isbrae) region of Greenland  
4 to study the spatial and intra-seasonal variability of supraglacial lakes during summer 2014.  
5 We also give coefficients for estimating supraglacial lake depth using a similar method as  
6 OLI with other multispectral sensors.

7

## 8 1 Introduction & Rationale

9 Supraglacial lakes in Greenland play a crucial role in the [ice sheet's](#) hydrological system.  
10 Together with supraglacial streams (Smith et al., 2015), lakes temporarily store large  
11 quantities of meltwater which can promote the opening of conduits to the bed through  
12 hydrofracture (Das et al., 2008; Phillips et al., 2013; Selmes et al., 2011; Tedesco et al., 2013)  
13 [and thus influence ice dynamics](#) (Joughin et al., 2013; Parizek and Alley, 2004; Sundal et al.,  
14 2011; Zwally et al., 2002). [Supraglacial lakes are influence surface heat fluxes by storing](#)  
15 [latent heat near the surface of the ice sheet](#) (Koenig et al., 2015). Finally, [supraglacial lakes](#)  
16 [contribute to multiple positive feedback processes, including ice shelf disintegration in](#)  
17 [Antarctica](#) (Banwell et al., 2013; Glasser and Scambos, 2008) [and melt-albedo interactions](#)  
18 [\(Leeson et al., 2015\)](#).

19 Several multispectral remote sensing tools and methods exist for both classifying (Johansson  
20 and Brown, 2013; Leeson et al., 2013; Sundal et al., 2011) and estimating the depth of  
21 supraglacial lakes (Sneed and Hamilton, 2007) [in Greenland](#). MODIS (the MODerate  
22 Resolution Imaging Spectroradiometer) is able to provide large spatial coverage (2,330 km  
23 swath width), moderate resolution (~250 m) images [of Greenland](#) twice per day (e.g., Box and  
24 Ski, 2007; Fitzpatrick et al., 2013). ASTER (the Advanced Spaceborne Thermal Emission and  
25 Reflection Radiometer, e.g. Sneed and Hamilton, 2007) and Landsat (e.g. Banwell et al.,  
26 2014; Morriss et al., 2013) have higher spatial resolution (10-30 m) but lower spatial coverage  
27 and fewer acquisitions (16 day repeat). Commercial sensors, such as [WorldView-2 and](#)  
28 [Worldview-3](#), provide high resolution multispectral measurements (~2 m) that can be used to  
29 image [small water features such as streams over smaller areas](#) (17 km wide swath), at both  
30 high temporal and spatial resolution (Chu, 2014; Legleiter et al., 2014; Smith et al., 2015).  
31 However, commercial imagery is collected largely 'on demand' and cloud cover can still be a

Allen Pope 21/10/2015 07:56
Deleted: glacial
Unknown
Field Code Changed
Allen Pope 26/10/2015 16:17
Deleted: and thus influence ice dynamics
Unknown
Field Code Changed
Allen Pope 26/10/2015 16:17
Deleted: Surface runoff has been increasing in Greenland over the past years (Seo et al., 2015), and supraglacial lakes have been shown to establish hydrological connections to the glacier bed and speed up flow, both through observational tools (Joughin et al., 2013) and models (Parizek and Alley, 2004).
Allen Pope 21/10/2015 08:00
Deleted: In Antarctica, supraglacial lakes can contribute to ice shelf disintegration, possibly through positive feedback mechanisms
Unknown
Field Code Changed
Allen Pope 21/10/2015 08:01
Deleted: Supraglacial lakes are also important for surface heat fluxes by storing latent heat near the surface of the ice sheet. In addition, lakes reduce surface albedo, <a href="#">k... [1]</a>
Unknown
Field Code Changed
Unknown
Field Code Changed
Allen Pope 21/10/2015 08:03
Deleted: (Sneed and Hamilton, 2011)
Unknown
Field Code Changed
Unknown
Field Code Changed
Unknown
Field Code Changed
David Shean 21/10/2015 11:02
Deleted: the
David Shean 21/10/2015 11:02
Deleted:
Allen Pope 21/10/2015 08:04
Deleted: -2
David Shean 21/10/2015 11:02
Deleted: series
Allen Pope 26/10/2015 16:31
Deleted: relatively
Allen Pope 26/10/2015 16:31
Deleted: during the daylight period
Unknown
Field Code Changed

1 | confounding factor. [Here we provide the first regional scale validation of supraglacial lake](#)  
2 | [depth estimation methods](#) with all of the above multispectral sensors.

3 | Lake depth retrieval is based upon the [understanding that deep water absorbs more energy](#)  
4 | [than shallow water](#) and therefore will have lower reflectance [of solar radiation](#). Some methods  
5 | use one band for a reflectance-depth relationship, while others use a ratio of reflectances from  
6 | two different spectral bands (see Sect. 2). [Satellite retrieval of supraglacial lake depth is](#)  
7 | [confounded by difficulty measuring the true reflectance of dark/deep lakes, assumptions](#)  
8 | inherent in the method about minimal quantities of suspended and dissolved matter in lake  
9 | water, [the requirement for](#) a smooth (i.e., not wind-roughened) lake surface, and  
10 | homogeneous and low-slope lake bottoms (Sneed and Hamilton, 2011). [In this study we](#)  
11 | [assume that it is possible to apply](#) locally calibrated coefficients to [broad areas](#) (e.g., Legleiter  
12 | et al., 2014), and can ignore minor variations in effects of atmospheric path radiance.

13 | With the successful launch of Landsat 8 in 2013 (Irons et al., 2012; Roy et al., 2014), a new  
14 | and improved multispectral sensor is available for lake depth estimation. The Landsat 8  
15 | Operational Land Imager (OLI) has enhanced radiometric resolution (12-bit vs. 8-bit), a  
16 | higher signal to noise ratio, and an expanded dynamic range compared to Landsat 7's  
17 | Enhanced Thematic Mapper Plus (ETM+). While published studies have largely used red and  
18 | green wavelengths, OLI's two additional bands (coastal, 0.433-0.453  $\mu\text{m}$ ; and cirrus, 1.360-  
19 | 1.390  $\mu\text{m}$ ) and narrower multispectral and panchromatic bands relative to ETM+ will provide  
20 | more spectral information and more unique (i.e., less auto-correlated) reflectance values,  
21 | respectively. These properties lead to improvements for [lake depth retrieval](#) methods based on  
22 | band ratios. [Furthermore](#), an increased scene collection rate by Landsat 8 will lead to more  
23 | opportunities to observe ice sheets and their supraglacial lakes.

24 | In this paper we investigate refinements on retrieval methods for supraglacial lake depth from  
25 | Landsat 8 imagery. We use in situ spectral measurements from a supraglacial lake in  
26 | Greenland to emulate satellite reflectance, and compare them with depth data from [the same](#)  
27 | lake to test several techniques to extract lake depth. [We then apply the best methods to](#)  
28 | Landsat 8 OLI imagery for case study areas in Northwest Greenland and the Sermeq Kujalleq  
29 | (Jakobshavn Isbrae) area. We validate depth estimates using digital elevation models (DEMs)  
30 | derived from stereo sub-meter imagery. We discuss best practices for deriving lake depths  
31 | using Landsat 8 and the implications of these conclusions for other multispectral sensors.

- Allen Pope 26/10/2015 16:32  
**Deleted:** For
- Allen Pope 26/10/2015 16:32  
**Deleted:** , a large scale validation has not been undertaken to date
- Allen Pope 26/10/2015 16:33  
**Deleted:** intuitive
- Allen Pope 26/10/2015 16:33  
**Deleted:** er
- Allen Pope 26/10/2015 16:33  
**Deleted:** will
- Allen Pope 26/10/2015 16:33  
**Deleted:** values
- Allen Pope 26/10/2015 16:33  
**Deleted:** However, s
- Allen Pope 26/10/2015 16:34  
**Deleted:** faces several limitations, including
- Allen Pope 26/10/2015 16:34  
**Deleted:** We
- Allen Pope 26/10/2015 16:34  
**Deleted:** further
- Allen Pope 26/10/2015 16:34  
**Deleted:** we can extrapolate
- Allen Pope 26/10/2015 16:35  
**Deleted:** new
- Unknown  
**Field Code Changed**
- Unknown  
**Field Code Changed**
- Allen Pope 26/10/2015 16:37  
**Deleted:** as
- Allen Pope 26/10/2015 16:37  
**Deleted:** Finally
- Allen Pope 26/10/2015 16:37  
**Deleted:** ,
- Allen Pope 26/10/2015 16:37  
**Deleted:** then we
- Allen Pope 21/10/2015 08:05  
**Deleted:** a
- Allen Pope 21/10/2015 08:05  
**Deleted:** supraglacial
- Allen Pope 26/10/2015 16:38  
**Deleted:** Based on this analysis, w

1 Analysis of 2014 imagery yields information about supraglacial lake size, distribution, and  
2 seasonal behavior.

## 3 2 Methods

### 4 2.1 Physically Based Lake Depth

5 The depth of a supraglacial lake can be approximated as (after Philpot, 1989):

$$6 \quad z = [\ln(A_d - R_\infty) - \ln(R_{lake} - R_\infty)] / g \quad (1)$$

7 where  $z$  is lake depth in meters,  $A_d$  is the lake bottom albedo,  $R_\infty$  is the reflectance of optically  
8 deep water,  $R_{lake}$  is the reflectance of a lake pixel, and  $g$  is related to the losses in upward and  
9 downward travel through the water column (units:  $m^{-1}$ ). This method has been used  
10 successfully in both Greenland and Antarctica (e.g., Banwell et al., 2014; Sneed and  
11 Hamilton, 2007). The method is based upon a description of the processes that take place as  
12 light enters, passes through, and exits a lake, it is physically based and therefore easy to adjust  
13 if measurements of lake water and lake bed properties are available. However, this method  
14 assumes that lake water has little to no dissolved or suspended matter and would be severely  
15 impacted by surface waves (wind-driven ripples, choppy waves, etc.). Additionally, it requires  
16 that the lake bottoms have low slopes and a homogeneous albedo (Sneed and Hamilton,  
17 2011). While most of these assumptions hold for supraglacial lakes in Greenland (Sneed and  
18 Hamilton, 2011), lake bottoms are known to be too inhomogeneous to support the approach  
19 generally. In addition, optically deep water (i.e., deep lakes or ocean where the upwelling  
20 radiance originates from the water column without any bottom signal contribution) is not  
21 always available in inland Landsat scenes. The effects of these shortcomings on supraglacial  
22 lake depth retrievals have not been quantified.

23 In this study, for application to Landsat 8 imagery,  $R_\infty$  was obtained from dark ocean or lake  
24 water in the scene, following Sneed and Hamilton (2007, 2011). If no coast was available in  
25 the scene containing the lake,  $R_\infty$  was obtained from another scene further along the path  
26 (with an implicit assumption of similar atmospheric conditions). The parameter  $g$  was  
27 calculated following earlier studies (Smith and Baker, 1981; Sneed and Hamilton, 2007), but  
28 with an updated absorption coefficient from Pope and Fry (1997, Table 3); for more details,  
29 see the Supplementary Material.

Unknown  
Field Code Changed

Allen Pope 26/10/2015 16:38  
Deleted: (units: m)

Allen Pope 6/9/2015 15:35  
Deleted:  $R_\infty$  is the reflectance of optically deep water,  $A_d$  is the lake bottom albedo,

Allen Pope 26/10/2015 16:38  
Deleted: several studies in

Unknown  
Field Code Changed

Allen Pope 26/10/2015 16:39  
Deleted: Since it

Allen Pope 21/10/2015 08:07  
Deleted: it requires optically deep water (i.e., no detectable reflected component from the lake bed) to be present in the images. It further

Allen Pope 21/10/2015 08:08  
Deleted: bed

Allen Pope 21/10/2015 08:08  
Deleted: surfaces

Allen Pope 26/10/2015 16:43  
Formatted: English (UK)

Unknown  
Field Code Changed

Allen Pope 21/10/2015 08:09  
Deleted: parameter

Unknown  
Field Code Changed

Unknown  
Field Code Changed

$A_d$  was obtained from the reflectance immediately outside identified lake areas. However, in order to test this approximation for  $A_d$ , we also solve for lake bottom albedo rather than assuming it to be the same as the surrounding ice. We use spectral mixture analysis (Lillesand et al., 2007) to define a fractional coverage of ice ( $r_i$ ) and cryoconite ( $r_c = 1 - r_i$ ) in each lake bottom pixel. We then use these coefficients and end-members of ice and cryoconite reflectances to calculate the lake bottom albedo ( $A_d$ ). To create a determinable equation after introducing this new unknown ( $r_i$ ), we use reflectances from two OLI spectral bands (indicated with subscripts 1 and 2, below), and derive end-member reflectances for ice ( $R_{i1}$  or  $R_{i2}$ ) and cryoconite ( $R_{c1}$  or  $R_{c2}$ ) using glacier reflectance spectra from Pope and Rees (2014b) in conjunction with OLI spectral response functions in both bands (Barsi et al., 2014). We input these parameters into Equation 1 and then combine the expressions by equating lake depth, thus obtaining:

$$\left[ \frac{r_i(R_{i1} - R_{c1}) + R_{c1} - R_{\infty 1}}{R_{w1} - R_{\infty 1}} \right] g_2 = \left[ \frac{r_i(R_{i2} - R_{c2}) + R_{c2} - R_{\infty 2}}{R_{w2} - R_{\infty 2}} \right] g_1 \quad (2)$$

After Eq. 2 is solved for  $r_i$ , the bottom albedo for one OLI spectral band can be calculated and subsequently used to compute lake depth:

$$A_{d1} = r_i R_{i1} + (1 - r_i) R_{c1} \quad (3)$$

$$z = \frac{\log(R_{lake1} - R_{\infty 1}) - \log(A_{d1} - R_{\infty 1})}{-g_1} \quad (4)$$

where  $R_{lake1}$  is water leaving reflectance (as in Eq. 1) for the first band in the pair used and  $z$  is lake depth.

[[Table 1]]

## 2.2 Empirically Derived Lake Depth

The second method we consider uses spectral band ratios has been used to derive water depth in shallow marine settings (e.g., Dierssen et al., 2003) and alluvial rivers (e.g., Legleiter and Overstreet, 2012), and has been also adapted for use on the Greenland ice sheet (Legleiter et al., 2014). While the physically based method above is highly dependent on  $A_d$  and  $g$ , earlier studies show that the band ratio method is expected to be more robust to variations in these parameters (Legleiter et al., 2009; Stumpf et al., 2003). This is because the method relies on relative behavior in two different wavelengths, as opposed to absolute optical behavior.

Allen Pope 21/10/2015 08:10  
 Deleted: As the spatially closest available estimate for lake bottom albedo,

Allen Pope 21/10/2015 08:11  
 Deleted: assumption

Allen Pope 21/10/2015 08:12  
 Deleted: In an application of a form of spectral mixture analysis

Unknown  
 Field Code Changed

Allen Pope 21/10/2015 08:12  
 Deleted: , we

Allen Pope 6/9/2015 15:36  
 Deleted: (Cryoconite is accumulated small surface debris; Wientjes et al., 2011.)

Unknown  
 Field Code Changed

Allen Pope 6/9/2015 15:37  
 Deleted: f

1 This band ratio method employs an empirically derived quadratic formula to relate lake  
2 depths to the ratio of the reflectance of two spectral bands ( $R_1$  and  $R_2$ ):

$$3 \quad z = a + bX + cX^2 \quad (5)$$

$$4 \quad X = \ln(R_1/R_2) \quad (6)$$

5 This empirical / band ratio method requires the derivation of calibrated coefficients (i.e.  $a$ ,  $b$ ,  
6 and  $c$ ), and coefficients vary depending on which sensors and bands are used (Legleiter et al.,  
7 2014). We calculate these coefficients using a known set of reflectances and depths (from in  
8 situ measurements, see Sect. 3.1 & 4.1).

### 9 3 Data

10 We use three datasets in this study: in situ reflectance spectra and lake depth, Landsat 8  
11 multispectral imagery, and DEMs derived from stereo WorldView imagery. We use in situ  
12 data to test different lake retrieval methods for a range of spectral bands. Then, we calculate  
13 lake depth with a range of the most promising methods using OLI imagery. We then use  
14 WorldView DEMs to validate the OLI-derived lake depths. The detailed workflow of  
15 software (including MATLAB and [shell scripts that call GDAL utilities](#)) used for data  
16 analysis and presentation in this study will be fully described and documented in a subsequent  
17 paper (Pope, in review).

#### 18 3.1 In Situ Data

19 Tedesco and Steiner used a small remote-controlled boat with a payload including a compact  
20 spectroradiometer and a small sonar to collect in situ, coincident lake-bottom reflectance and  
21 depth over one lake west Greenland in the summer of 2010 (Tedesco et al., 2015; 2011). We  
22 use 2226 unique sample points from that study to evaluate the performance of the remote  
23 sensing methods described above. Field spectra are convolved to account for the spectral  
24 response of the spaceborne sensors as follows:

$$25 \quad r_{nb} = \frac{\int_0^\infty r(\lambda)R(\lambda)d\lambda}{\int_0^\infty R(\lambda)d\lambda} \quad (7)$$

26 where  $r_{nb}$  is the narrowband reflectance,  $r(\lambda)$  is the spectral reflectance,  $R(\lambda)$  is the relative  
27 spectral response (Barsi et al., 2014), and  $\lambda$  is the wavelength. In order to emulate sensor  
28 dynamic range and radiometric resolution, we impose minimum and maximum reflectances  
29 and round reflectance values to the appropriate precision (i.e., 8-bit or 12-bit; see Pope and

Allen Pope 21/10/2015 08:17  
Deleted: based  
Unknown  
Field Code Changed

Allen Pope 21/10/2015 10:07  
Deleted: T  
Allen Pope 21/10/2015 10:07  
Deleted: are used

Allen Pope 21/10/2015 17:20  
Deleted: code

Allen Pope 21/10/2015 08:18  
Deleted: In situ concurrent and bathymetric  
measurements were collected using

Allen Pope 21/10/2015 10:10  
Deleted: (Tedesco and Steiner, 2011)  
Unknown  
Field Code Changed

Unknown  
Field Code Changed

Unknown  
Field Code Changed

1 Rees, 2014a). We then regress the convolved reflectances and in situ depth measurements to  
2 test the goodness of fit of the physically based relationship presented in Eq. 1 and the  
3 empirical method described in Eqs. 5 and 6.

### 4 3.2 Landsat 8 Imagery

5 Landsat 8 launched on 11 February 2013 and became operational on 30 May 2013 (Roy et al.,  
6 2014). OLI collects spectral data gridded at 30 m spatial resolution (15 m for panchromatic  
7 data). We calculate top-of-atmosphere (TOA) reflectance using calibration coefficients  
8 provided in the image metadata and a solar elevation cosine correction (USGS, 2013). Based  
9 on a sensitivity analysis of path radiance to water vapor and ozone using an atmospheric  
10 radiative transfer model (see Sect. 5), we did not atmospherically correct the images.

11 We choose two study areas for applying OLI imagery (see Fig 1). One site located in  
12 northwest Greenland (including Sverdrup Gletsjer, Dietrichson Gletsjer, Sermersuaq, and  
13 Kjer Gletsjer, on Melville Bay; 56.2966-58.7186°W, 74.9685-75.7808°N) is an area with a  
14 high concentration of lakes and was imaged four times by Landsat 8 throughout summer  
15 2013. A larger region farther to the south is examined using all available Landsat 8 scenes  
16 collected over the Sermeq Kujalleq (Jakobshavn Isbrae) region in West Greenland in 2014.  
17 For a list of all OLI scenes used in this study, see Table S2.

18 [[Fig. 1]]

19 Using the calculated TOA reflectances, we define supraglacial lake extent using the ratio  
20 between the blue and red bands (Banwell et al., 2014; Box and Ski, 2007). However, since  
21 OLI bands are slightly different from those of past sensors, we could not use published  
22 thresholds for extent. We set the threshold for this ratio at 1.5 (vs. 1.05-1.25 for ETM+ in  
23 Banwell et al., 2014) based upon visual comparison with the imagery. We then visually  
24 inspected and manually adjusted the threshold mask to remove coastal water areas (i.e., not on  
25 the ice sheet) and clouds. Although Leeson et al. (2013) describe such thresholding as too  
26 coarse for low resolution imagery (i.e. MODIS), they do acknowledge its utility for higher  
27 resolution imagery (i.e. ASTER, Landsat, etc.). We remove regions four pixels or smaller (i.e.  
28 small lakes likely comprised solely of mixed pixels) or less than two pixels wide (i.e. linear  
29 features likely to be channels, not lakes) from the lake mask.

30 We interpolate the lake mask using a nearest neighbor algorithm, in order to apply the  
31 physically based method to the higher resolution panchromatic band. Where both

Allen Pope 21/10/2015 10:11  
Deleted: was

Unknown  
Field Code Changed

Allen Pope 21/10/2015 10:11  
Deleted: T

Allen Pope 21/10/2015 10:11  
Deleted: is calculated

Unknown  
Field Code Changed

Allen Pope 21/10/2015 10:12  
Deleted: were not atmospherically corrected

Allen Pope 21/10/2015 10:12  
Deleted: T

Allen Pope 21/10/2015 10:12  
Deleted: are chosen

Allen Pope 21/10/2015 08:19  
Deleted: examined in 2013 as it as

Allen Pope 21/10/2015 08:19  
Deleted: 4

Allen Pope 21/10/2015 08:19  
Deleted: the

Allen Pope 21/10/2015 08:20  
Deleted: u

Allen Pope 21/10/2015 08:21  
Deleted: in 2014

Unknown  
Field Code Changed

Allen Pope 6/9/2015 15:38  
Deleted: based

Unknown  
Field Code Changed

Unknown  
Field Code Changed

Allen Pope 21/10/2015 10:12  
Deleted: R

Allen Pope 21/10/2015 09:57  
Deleted: 4

Allen Pope 21/10/2015 09:57  
Deleted: 2

Allen Pope 21/10/2015 10:12  
Deleted: were removed

1 panchromatic and spectral bands were used together, we bilinearly interpolate the  
2 panchromatic image to 30 m resolution.

### 3 3.3 WorldView DEMs

4 We use submeter (~0.5 m / pixel) stereo imagery from DigitalGlobe's WorldView-1 and  
5 WorldView-2 to create DEMs of lake areas both before filling and after drainage. [Similar  
6 validation for ASTER has been carried out with airborne LiDAR from before lake drainage  
7 \(Georgiou et al., 2009\), and for estimating lake drainage volumes \(Stevens et al., 2015\). We  
8 generate the high resolution WorldView DEMs using the open source NASA Ames Stereo  
9 Pipeline tool \(Moratto et al., 2010; Shean et al., 2015\). For both the Sermeq Kujalleq  
10 \(Jakobshavn\) and northwest sites, we use DEMs from six different days, for a total of 12  
11 DEMs \(see Table S2\).](#)

12 WorldView-1 image data have a geolocation accuracy of better than 4.0 m horizontal 90%  
13 circular error of probability and WorldView-2 better than 3.5 m (DigitalGlobe, 2014). Thus,  
14 the imagery and DEMs are more precisely positioned than the 15-30 m Landsat 8 pixels.

15 The vertical accuracy of the derived DEM products is less than 5.0 m 90% vertical error of  
16 probability with submeter relative vertical precision (Mitchell, 2010). Differencing a  
17 WorldView DEM with an [Airborne Topographic Mapper LiDAR](#) profile over a pronounced  
18 basin in northeast Greenland provided a standard deviation over the spread of elevations of  
19 0.25 m. Considered conservatively, differencing one WorldView DEM with a second DEM  
20 collected one year later provided a standard deviation of 0.58 m for the elevation differences  
21 (Willis et al., 2015). [Stacks of 13 and 17 overlapping WorldView-1 and WorldView-2 DEMs  
22 over Summit Station and Tracy Glacier, Greenland provide absolute vertical accuracy  
23 estimates of ~2.0-3.0 m relative to airborne LiDAR measurements \(~10 cm accuracy\). After  
24 removing absolute horizontal and vertical offsets from all DEMs, the relative vertical  
25 accuracy \(1-sigma\) for the stack was ~15-30 cm \(Shean et al., 2015\).](#)

26 We resample the DEMs to the same grid as Landsat imagery using cubic interpolation. The  
27 Landsat and WorldView acquisitions are from different dates, and although lake basins do  
28 ablate during the summer, this should not have significant impact on the results presented  
29 here, [because most supraglacial lakes in Greenland remain fixed over bedrock-controlled  
30 surface depressions \(Lampkin and VanderBerg, 2011\). Using the lake mask, we identify a  
31 shoreline for a given date \(see Sect. 3.2\), which is then used to derive lake depth. We remove](#)

Allen Pope 21/10/2015 08:22  
**Deleted: v**

Allen Pope 21/10/2015 10:13  
**Deleted: T**

Allen Pope 21/10/2015 09:29  
**Deleted: se**

Allen Pope 21/10/2015 10:13  
**Deleted: are generated**

Unknown  
**Field Code Changed** ... [2]

Allen Pope 21/10/2015 17:26  
**Deleted: (Shean et al., 2015; Moratto...** [3]

Allen Pope 21/10/2015 09:58  
**Deleted: 6**

Allen Pope 21/10/2015 10:14  
**Deleted: were used**

Unknown  
**Field Code Changed** ... [4]

Unknown  
**Field Code Changed** ... [5]

Allen Pope 6/9/2015 15:44  
**Deleted: a**

Allen Pope 6/9/2015 15:44  
**Deleted: t**

Allen Pope 6/9/2015 15:44  
**Deleted: m**

Allen Pope 6/9/2015 15:44  
**Deleted: lidar**

David Shean 21/10/2015 11:14  
**Deleted: A stack of 13**

Unknown  
**Field Code Changed** ... [6]

David Shean 21/10/2015 11:14  
**Deleted: (~17x17 km)**

David Shean 21/10/2015 11:14  
**Deleted: ,**

David Shean 21/10/2015 11:15  
**Deleted: s**

David Shean 21/10/2015 11:15  
**Deleted: (±1.2 m)**

Allen Pope 21/10/2015 08:24  
**Deleted: , according to preliminary studies**

David Shean 21/10/2015 11:16  
**Deleted: the**

David Shean 21/10/2015 11:17  
**Deleted: per-pixel standard deviation...** [7]

Allen Pope 21/10/2015 17:26  
**Deleted: (Shean et al., 2015)**

Allen Pope 21/10/2015 08:25  
**Deleted: as**

Unknown  
**Field Code Changed** ... [8]

1 | outliers of impossibly shallow (i.e. negative depth) or deep (>65 m) values as blunders in the  
2 | DEM. In addition, we remove lakes having a standard deviation in lake elevation along the  
3 | shoreline of larger than 1.5 m. These steps also mitigated any potential bias caused by  
4 | temporal offset between DEM and spectral depth measurements.

5 | After filtering, over 250,000 pixels (30 m) in total remained for spectral lake depth validation  
6 | over six days in 2013 and six days in 2014.

## 7 | 4 Results

### 8 | 4.1 In Situ Results

9 | The results (Table 1) of depth-reflectance regressions for all methods are shown in Fig. 2. We  
10 | base the bands tested here using in situ data upon those identified in the literature (e.g., Box  
11 | and Ski, 2007; Sneed and Hamilton, 2007; Tedesco and Steiner, 2011), as well as the OLI's  
12 | new coastal band and the significantly narrowed panchromatic band (0.500-0.680  $\mu\text{m}$ , at 15m  
13 | spatial resolution). ETM+ high and low gain results are virtually indistinguishable, and so  
14 | only low gain results are shown here. For each regression, we use the correlation coefficient  
15 | ( $r$ ) and the root mean square error (RMSE, relative to sonar depths) to assess the performance  
16 | of each method. The results of the physically based method show that the OLI blue and  
17 | coastal bands do not perform well relative to other bands (RMSE of 3.10 m and 11.03 m,  
18 | respectively;  $r$  of 0.29 and 0.05, respectively). The OLI Band 3 (green, 0.525-0.600  $\mu\text{m}$ ; 0.78  
19 | m,  $r = 0.78$ ) performs as well as legacy ETM+'s Band 2 (green, 0.525-0.605  $\mu\text{m}$ ; 0.77 m,  $r =$   
20 | 0.79). Finally, both OLI Band 4 (red, 0.640-0.670  $\mu\text{m}$ ) and Band 8 (panchromatic, 0.500-  
21 | 0.680  $\mu\text{m}$ ) bands outperform their analogous ETM+ bands (RMSE of 0.28 m and 0.63 m,  
22 | respectively;  $r$  of 0.96 and 0.84, respectively).

23 | [[Fig. 2]]

24 | Red light attenuates more strongly in water than green or blue light. So, for the same lake  
25 | depth, there will be a larger (and easier to measure) change in net reflectance for red  
26 | wavelengths than shorter wavelengths. However, the rapid attenuation of red light means that  
27 | only shallower lakes may be measured in this band. The maximum in situ lake depth  
28 | measurement is  $\sim 5$  m, well within the red light limit, but deeper lakes may exist in the overall  
29 | study area. We address this issue below by using many Landsat scenes and WorldView  
30 | DEMs.

Allen Pope 21/10/2015 08:26  
Deleted: errors

Allen Pope 26/10/2015 16:51  
Deleted: all

David Shean 21/10/2015 11:29  
Deleted: in total from

Allen Pope 21/10/2015 09:58  
Deleted: 6

Allen Pope 21/10/2015 09:58  
Deleted: 6

David Shean 21/10/2015 11:29  
Deleted: remained for spectral lake depth validation

Allen Pope 21/10/2015 08:27  
Deleted: displayed

Allen Pope 21/10/2015 08:27  
Deleted: Table 1 and

Allen Pope 21/10/2015 10:16  
Deleted: T

Allen Pope 21/10/2015 10:16  
Deleted: were based

Unknown  
Field Code Changed

Allen Pope 21/10/2015 08:28  
Deleted: success

Allen Pope 21/10/2015 08:30  
Deleted: Landsat 7

1 We investigate the two-band physically based method (where  $A_d$  was calculated) with a range  
2 of emulated OLI bands (see Table 1). We find similarly high correlation coefficients ( $r =$   
3 0.94) to the regression method. Nevertheless, only the combination of blue and green bands  
4 had an RMSE below 1 m. This method appears to slightly overestimate lake depths. We  
5 investigate the reasons for this with the Landsat and WorldView data below.

6 Applying the empirical method using field data (see Table 1, Fig. 2) indicates that the more  
7 continuous bands of the ETM+ outperform the narrower (less spectrally auto-correlated)  
8 bands of the OLI when estimating lake depths. However, the addition of the coastal band  
9 should allow the OLI to still perform quite well ( $r > 0.92$ ,  $RMSE < 0.38$ ), in particular when  
10 paired with the green or panchromatic bands.

11 Our analysis shows [that supraglacial lake depth retrievals using Landsat 8 are as good as or](#)  
12 [better than Landsat 7 retrievals](#). We identify the best methods for OLI (identified with  
13 asterisks in Table 1) based on the highest correlation coefficients and lowest RMSEs. We then  
14 apply these methods to Landsat 8 data and validate them with WorldView stereo DEMs.

Allen Pope 21/10/2015 08:34  
**Deleted:** the potential of using Landsat 8 to retrieve supraglacial lake depth as well as or better than with Landsat 7

## 15 4.2 2013 Northwest Greenland Results

16 In the northwest Greenland study area, we identified 694 lakes on 2 July 2013 (day of year  
17 183) with a total area of 27.2 km<sup>2</sup>, 1259 lakes totaling 43.7 km<sup>2</sup> on 18 July 2013 (day 199),  
18 955 lakes totaling 38.8 km<sup>2</sup> on 3 August 2013 (day 215), and 274 lakes totaling 8.6 km<sup>2</sup> on 19  
19 August 2013 (day 231). [We calculate lake depths with all previously discussed methods, as](#)  
20 [well as an average between the two best single-band depth estimates.](#) Total lake volume in the  
21 study area increased in early July, stayed almost constant as lake growth areas moved higher  
22 in elevation over the following three weeks, and then decreased again toward the end of  
23 August as cooler conditions prevailed (see Fig. 3). While all methods show the same pattern  
24 of surface water storage, the total water volumes [derived with](#) the different methods differ by  
25 over a factor of 2.

Allen Pope 21/10/2015 08:36  
**Deleted:** D

Allen Pope 21/10/2015 08:36  
**Deleted:** D

Allen Pope 21/10/2015 08:36  
**Deleted:** D

Allen Pope 21/10/2015 10:17  
**Deleted:** Lake

Allen Pope 21/10/2015 10:17  
**Deleted:** were calculated

Allen Pope 21/10/2015 08:34  
**Deleted:** returned

Allen Pope 21/10/2015 08:35  
**Deleted:** by

Allen Pope 21/10/2015 08:37

**Deleted:** We investigate which method agrees best with the WorldView DEMs, providing some confidence in the retrievals of that method.

26 [[Fig. 3]]

## 27 4.3 Comparison with DEMs

28 We difference [all](#) overlapping areas of DEM and Landsat-derived lake depths [in both case](#)  
29 [study regions](#). The statistics of this comparison are shown in Fig. 4. As seen in the northwest  
30 Greenland case study, the methods are divided into two groups. Landsat-derived depths using

1 band 3, bands 2 & 3, a ratio of bands 1 and 3, and a ratio of bands 1 and 8 all considerably  
2 overestimate lake depth relative to the DEMs. However, the physically based single band  
3 method for the red band (OLI Band 4) only slightly underestimates lake depth ( $-0.1 \pm 1.7$  m),  
4 while the panchromatic band (OLI Band 8) slightly overestimates lake depth ( $0.1 \pm 1.4$  m).

5 Combining these two best-performing bands, the resulting spectral and DEM-derived lake  
6 depths are in close agreement, showing a difference of  $0.0 \pm 1.6$  m. We infer that the optimal  
7 method for estimating supraglacial lake depth with Landsat 8 is to take an average of the  
8 physically based (see Eqn. 1) depths as derived from the red and panchromatic channels (bold  
9 in Table 1). It is likely that the spread in depths is the result of a combination of factors  
10 including temporal offset between DEM and spectral data collection, image coregistration,  
11 and atmospheric effects, as well as uncertainties inherent in the lake depth retrievals. Despite  
12 meter scale uncertainties (1.6m) at the pixel level, the mean lake depth derived from these  
13 methods agrees well.

14 [[Fig. 4]]

#### 15 4.4 2014 Sermeq Kujalleq (Jakobshavn) Area Results

16 We apply the lake depth algorithm (i.e., average of single band depths from OLI red &  
17 panchromatic bands) to 34 Landsat 8 scenes from the summer of 2014 over the Sermeq  
18 Kujalleq (Jakobshavn) area (see Figs. 1 & 5). The total meltwater storage in supraglacial lakes  
19 peaked near three cubic kilometers, across the entire study area in mid-July 2014. There are  
20 many shallow lakes (0.3 to 1.5 m depth) and many lakes with depths of 2.5 to 4 m. Few lakes  
21 exceed 5.5 m depth (see Fig. 6a). The preponderance of shallow lake pixels reflects the fact  
22 that the observed lakes have low surface slopes at their edges.

23 [[Fig 5]]

24 The Sermeq Kujalleq (Jakobshavn) dataset provides a timeseries that shows lake growth and  
25 drainage / freezing (see Fig. 5a). There are many factors that contribute to lake growth and  
26 drainage, including temperature, insolation, albedo, topography, and ice dynamics. These  
27 complex drivers are related to the more easily quantified mean elevation and latitude of each  
28 scene. For example, isolating the coastal scenes shows the delayed onset of melt and earlier  
29 shutdown in the north compared to the south (see Fig. 5b).

Allen Pope 21/10/2015 08:39

Deleted: very good

Allen Pope 21/10/2015 08:48

Deleted: ice flow,

Allen Pope 21/10/2015 08:49

Deleted: Therefore,

Allen Pope 21/10/2015 08:49

Deleted: while having a fairly large (1.6 m) uncertainty for individual pixels

Allen Pope 21/10/2015 10:18

Deleted: T

Allen Pope 21/10/2015 10:18

Deleted: was applied

Allen Pope 21/10/2015 09:57

Deleted: 3 km<sup>3</sup>

Allen Pope 21/10/2015 08:51

Deleted: A histogram of all lake depths shows that t

Allen Pope 6/9/2015 15:39

Deleted: , the distribution has many

Allen Pope 26/10/2015 16:51

Deleted: at

Allen Pope 26/10/2015 16:51

Deleted: , with f

Allen Pope 26/10/2015 16:52

Deleted: ing

1 To further refine our investigation of geographic factors associated with lake depth over the  
2 summer season, we examine single swaths of Landsat imagery through time. Path 008 (in the  
3 WRS-2 reference scheme, Irons et al., 2012) which transects the lower Sermeq Kujalleq  
4 (Jakobshavn Isbrae) shows a strong influence of both elevation and latitude in rates of lake  
5 growth and water storage (Fig. 5c). Isolating Path 006, on the other hand, conflates the effects  
6 of elevation and altitude on surface meltwater storage, but because we have more temporal  
7 coverage (see Fig. 5d) we see the decline of total lake volume as summer progresses toward  
8 autumn. Again, higher latitude and elevation delay melt onset (i.e. Path 006, Row 012). For  
9 006/013 and 006/014, it is likely that the reduced ice sheet area within 006/014 is the  
10 explanation for the reduced meltwater volume. Rates of increase and decay of lake volume are  
11 similar for this pair.

12 The distribution of lake depths (by pixel) with elevation is shown in Fig. 6b. Lakes are  
13 distributed from ~300 m to ~2100 m elevation. Maximum lake depths occur at about 1200 m  
14 a.s.l. At lower elevations, lake depths recorded by our method vary significantly, likely due to  
15 rapid lake growth and drainage across a range of dates at lower elevations, versus the higher  
16 elevation maximum depths mostly derived from a Landsat scene on July 30 2014. From 1200  
17 m to 2100 m, measured lake depths decline steadily with less variation. This likely reflects a  
18 combination of factors, including the variations in induced surface topography of the ice sheet  
19 as it flows over undulating bedrock (Lampkin and VanderBerg, 2011). At higher elevations,  
20 slow flow leads to low-amplitude ice surface topography as well as less available meltwater.  
21 In addition, while lakes are less likely to variably fill and drain at higher elevations, there was  
22 also reduced imagery available from ~July 30 2014 onwards. Therefore, the more consistent  
23 maximum depths at higher elevations are a combination of incomplete temporal coverage and  
24 [elevation](#). Further down, more melt and higher amplitude topography from faster ice flow  
25 facilitate lake formation. However, below 1200 m, increased ablation begins to reduce this  
26 topography. In addition, the volume of melt available will determine whether depressions are  
27 large enough to hold lakes or instead drain via connecting supraglacial channels. The melt  
28 volume and therefore the relationship between lakes and channels will thus vary both  
29 seasonally and with elevation as well (Lampkin and VanderBerg, 2014).

30 [[Fig 6]]

Unknown  
Field Code Changed

Unknown  
Field Code Changed

## 5 Discussion

### 5.1 Retrieval Performance Factors

The depths returned by the empirical (band ratio) method considerably overestimate lake depths relative to the WorldView DEMs. The method is entirely dependent upon the calibration of the input parameters (i.e.,  $a$ ,  $b$ , and  $c$ ). The parameters used in this study are in turn based solely upon extrapolation from in situ measurements at a single lake. Therefore, it is possible that the lake used for calibration is not representative of lakes in our study region. Legleiter et al. (2014) note that the coefficients for the empirical method may be scale-dependent, and values calculated from field data may not be appropriate for the 30 m pixels of Landsat 8. Indeed, other work (Moussavi et al., in review) both calibrates and validates spectrally derived depths with WorldView DEMs to show that the band-ratio/empirical method and single-band/physically based method perform similarly well. The use of a ratio of coastal and green reflectances performed well for lake depth retrieval using WorldView-2 imagery (Legleiter et al., 2014). Therefore the band ratio method may, with better parameters, produce results consistent with the physically-based single-band approaches.

Nevertheless, the physically based depth retrievals show a large spread in total water volume returns. Physically based depth retrievals rely on accurate bottom albedos ( $A_d$ ) and water absorption coefficient ( $g$ ). While  $A_d$  is derived from the imagery,  $g$  is always calculated for each spectral band based on laboratory measurements and is therefore consistent across all OLI scenes. Comparison of laboratory-measured  $g$  values with  $g$  derived from in situ data (see Table 1) shows that when the laboratory-measured  $g$  is higher than the one obtained from regressing in situ data, lake depths are overestimated and vice versa. For example, OLI Band 3 (green) shows a 70% difference in directly measured and regressed  $g$ , and it overestimates lake depths by a mean of  $2.4 \pm 2.1$  m relative to WorldView DEMs. By contrast, Band 4 (red) and Band 8 (panchromatic) have very small differences between measured and regressed  $g$  (-0.06% and 0.06%, respectively) and yield accurate lake depth estimates ( $-0.1 \pm 1.7$  m and  $0.1 \pm 1.4$  m, respectively) relative to WorldView DEMs.

Water absorption properties also vary with wavelength. For example, poor performance in blue and coastal bands is related to very low absorption. Red wavelengths attenuate relatively quickly in water, and this is described by a relatively high  $g$  ( $0.7507 \text{ m}^{-1}$ ) compared to green ( $0.1413 \text{ m}^{-1}$ ). This high  $g$  value for red light makes it less sensitive to errors in  $g$  than green

Allen Pope 21/10/2015 09:51

**Deleted:** In the above results section, we presented data showing promising lake depth retrieval techniques, used Landsat imagery and WorldView DEMs to test these methods, and applied imagery towards an understanding of the supraglacial hydrology of a section of the Greenland ice sheet. In discussion below, we consider factors which control the success and/or failure of both lake depth retrieval methods (i.e. physically based and empirically derived) as well as the merits of particular wavelengths of light for lake depth estimation. We then apply this understanding to assess earlier lake depth retrievals in the literature that used ETM+ and ASTER data. We then include a sensitivity analysis of atmospheric corrections of OLI imagery and finish with a discussion of limitations to OLI lake depth retrievals. .

Allen Pope 21/10/2015 09:49

**Formatted:** Heading 2

Unknown

**Field Code Changed**

Allen Pope 21/10/2015 09:07

**Deleted:** preliminary

Allen Pope 21/10/2015 09:33

**Deleted:** (a separate study led by co-author Moussavi)

Unknown

**Field Code Changed**

Allen Pope 21/10/2015 09:08

**Deleted:** perform similarly to

Allen Pope 21/10/2015 09:08

**Deleted:** values

Allen Pope 21/10/2015 09:09

**Deleted:** m

wavelengths. Lake depth estimates using a red channel are also less sensitive to  $A_d$  than with a green channel (Tedesco and Steiner, 2011), again due to the high absorption for longer wavelengths. Ultimately, as long as the sensor radiometry is able to measure the return from deep-water pixels, longer wavelengths (i.e., red) can return generally more accurate lake depths because they are less sensitive to the input parameters.

## 5.2 Revisiting Lake Depths Retrievals

To evaluate other studies in the literature and compare them to our results, we applied the same methods we use (i.e., lab-measured absorption/scattering parameters and appropriate spectral response functions) to calculate  $g$  values for Landsat 7 ETM+ bands (see Table 1). Tedesco and Steiner (2011) studied the accuracy of ETM+'s green band for lake depth estimation. They tested different multipliers of the diffuse attenuation coefficient for downwelling light to get the water absorption coefficient  $g$ . They showed that for Landsat 7 ETM+'s green band, sonar and spectral depths correlated better when a larger multiplier was used. This is broadly consistent with the 70% offset between observed and theoretical values that we observe (Table 1). They also find that this offset "cannot be easily explained, aside from a possible chlorophyll concentration in the water, currently considered to be unlikely." Morriss et al. (2013) used ETM+'s red band and extracted a higher value of  $g$  ( $0.86 \text{ m}^{-1}$ ); this is very close to the regressed value we observe of  $0.83 \text{ m}^{-1}$  (see Table 1), and so we expect their depth estimates to be slightly overestimated.

Banwell et al. (2014) and Arnold et al. (2014) also used Landsat 7's green band with a  $g$  of  $0.1954 \text{ m}^{-1}$ , ~40% percent higher than our regressed value of  $0.14 \text{ m}^{-1}$ , leading to depths overestimated by ~30%. Because the comparisons of Greenland and Antarctic lakes (Banwell et al., 2014) are based on relative depths, their conclusions are likely still valid. Arnold et al. (2014) concluded that their model under-predicted water depths, which could in reality mean that their model is behaving correctly but their validation data (i.e. Landsat lake depths) were biased.

Using the same process as for Landsat sensors, we calculated  $g$  values for ASTER, MODIS, and WorldView-2 bands (see Table S1). Sneed and Hamilton (2007, 2011) used ASTER's green band for lake depth estimation ( $g = 0.1180 \text{ m}^{-1}$ ). This is ~20% smaller than the regressed value of  $0.15 \text{ m}^{-1}$  (see Table S1). They will therefore have likely underestimated lake depth (Sneed and Hamilton, 2007).

Allen Pope 21/10/2015 09:50  
Formatted: Heading 2

Unknown  
Field Code Changed

Unknown  
Field Code Changed

Unknown  
Field Code Changed

Allen Pope 21/10/2015 09:12  
Deleted: probably

Allen Pope 21/10/2015 09:12  
Deleted: significantly

Allen Pope 21/10/2015 09:12  
Deleted: lake depths, estimated to be

Unknown  
Field Code Changed

Unknown  
Field Code Changed

Allen Pope 21/10/2015 09:13  
Deleted: believed

Allen Pope 21/10/2015 09:14  
Deleted: Shallower lake depths would also likely reduce the identified threshold for initiation of hydrofracture - if indeed lake volume initiates drainage as opposed to other basal hydrological factors (i.e., reduction of local compressional stresses, Stevens et al., 2014).

Unknown  
Field Code Changed

1 In all three studies, the regressed  $g$  values are much closer to the updated lab-based  $g$  values  
2 ([see Sect. 2.1 & Supplementary Material](#)) than those used previously used in the literature.  
3 Adoption of the new  $g$  values presented here in Tables 1 and S1 would therefore likely lead to  
4 improved lake depth estimates.

### 5 **5.3 Sensitivity Analysis**

6 [For all sensors, wavelengths, and input parameters, an important consideration for](#)  
7 [reflectance-derived lake depth is the atmospheric correction used to prepare the multispectral](#)  
8 [imagery. All imagery is processed to TOA reflectance, which means that there is some](#)  
9 [extraneous path radiance remnant in the data. Therefore, TOA values will slightly](#)  
10 [overestimate the true reflectance. This offset will not be the same between bands, and will](#)  
11 [influence the retrieved lake depths as discussed below.](#)

12 [The single band physically based model requires that the reflectance of optically deep water](#)  
13 [be derived for each scene separately. Effectively, this shifts the exponential decay curve of](#)  
14 [light in lake water but does not change its shape. Therefore, as long as path radiance is](#)  
15 [assumed to be homogeneous across the 185-km wide Landsat scene, TOA reflectance is](#)  
16 [sufficient for lake depth estimation. To test this assumption, the MODTRAN radiative](#)  
17 [transfer model \(Berk et al., 2005\) was used to simulate path radiance on a day for which](#)  
18 [Landsat data were used in northwest Greenland \(18 July 2013\) to investigate variations](#)  
19 [associated with variable water vapor and ozone across a Landsat scene. According to MODIS](#)  
20 [retrievals \(accurate to 30 DU; Borbas et al., 2011\), ozone variability within a Landsat scene is](#)  
21 [on the order of approximately  \$\pm 50\$  DU, which translates to a path radiance of  \$\pm 1.6\%\$  in the red](#)  
22 [channel. For lake depth, this can propagate to a  \$\sim 20\%\$  error in lake depth. Much of this error](#)  
23 [appears largely random for a given point in time and space. Thus, while it decreases](#)  
24 [confidence in individual lake depth retrievals, averaged water volume retrieval should not be](#)  
25 [biased. For water vapor there was a 0.3% change in path radiance between the minimum and](#)  
26 [maximum Landsat scene values, making it a small contributor to overall error. Between days,](#)  
27 [however, path radiance effects due to water vapor may vary by an order of magnitude more.](#)

28 [For the multiple band methods, the differential change in path radiance has larger effects.](#)  
29 [Sensitivity tests showed that a 3% change in path radiance for one or both bands changed](#)  
30 [water volumes on the order of 10-30%. Therefore, a more rigorous atmospheric correction is](#)  
31 [necessary in order to apply multi-band lake depth algorithms. Still, for the study here, because](#)

Allen Pope 21/10/2015 09:15

**Deleted:** discussed in the previous paragraphs

Allen Pope 21/10/2015 09:16

**Deleted:** (see Sect. 2.1)

validation is conducted across 12 non-consecutive days in both spring and autumn, we do not expect atmospheric conditions to bias our conclusions.

There are additional limitations to our method. As discussed above, OLI lake depth estimates (average single-band estimates from red and panchromatic bands) are robust for regional averages but not single pixels. In addition, the threshold used to identify lake extent may need to be adjusted for different regions and scenes (e.g. Banwell et al., 2014; Box and Ski, 2007). Lake depth retrievals are also sensitive to variations in ice albedo, as well as to the presence of ice lids on the surface of supraglacial lakes, which can be common in both early and late summer. Cloud cover and Landsat's 16-day revisit time also limit the conclusions that can be drawn from OLI lake depths. Many studies have used daily MODIS data to identify and track supraglacial lakes (e.g. Liang et al., 2012; Selmes et al., 2011; Sundal et al., 2011). Fusing the higher temporal resolution of MODIS (or additional sensors such as ESA's upcoming Sentinel-2) and higher spatial resolution of Landsat, along with more in situ calibration and validation data, should lead to unique insights to supraglacial water storage.

#### 5.4 Supraglacial Lakes in the Hydrological System

Both supraglacial lakes and channels can contribute significantly to regional water storage and transport (Smith et al., 2015). If the water stored in supraglacial lakes in row 12 of path 008 in mid-July were spread across the whole 25,246 km<sup>2</sup> of ice in the scene, it would have an average depth of almost three centimeters. In other scenes, calculations provide average depths of 0.5 to 1.5 cm. Our maximum observed value is almost as high as the volume in supraglacial streams measured by Smith et al. (2015), reinforcing the potentially daily turnover of a well-connected surface system they observed. Indeed, Tedesco et al. (2012) observe bare ice melt rates next to supraglacial lakes in west Greenland of ~2.5-3 cm per day, similar to those observed by van den Broeke et al. (2011). This implies that lakes are storing on the order of one day's worth of melt (or less), indicating daily or subdaily residence times, depending on connectivity.

Lakes, therefore, provide only a transient role in water storage. So, what is their importance to the ice sheet? Lakes have already (Howat et al., 2013) and are projected to continue to advance inland under a warming climate, with a minimal effect on overall ice sheet albedo but the potential to increase water transfer to the bed in areas without efficient drainage networks, therefore speeding up ice flow (Leeson et al., 2015). Indeed, episodic transport of water to the

Allen Pope 21/10/2015 09:50  
Formatted: Heading 2

Allen Pope 21/10/2015 09:19  
**Deleted:** In addition, historical Landsat imagery has been used to study the spatial evolution of melt ponds over the past decade (Howat et al., 2013); no revision of these conclusions are relevant, but improved confidence in lake depth estimates now warrants investigation of possible changes in surface meltwater storage in supraglacial lakes using the historical record.

Unknown  
**Field Code Changed**

Allen Pope 21/10/2015 09:58  
**Deleted:** 3

Allen Pope 21/10/2015 09:58  
**Deleted:** m

Unknown  
**Field Code Changed**

Unknown  
**Field Code Changed**

Unknown  
**Field Code Changed**

1 bed is expected to have a larger effect than continuously increased fluxes which set up an  
2 efficient drainage system (Flowers, 2015). In addition to episodically increased velocities, the  
3 enhanced hydrofracture propagated by the water stored in supraglacial lakes may play a key  
4 role in heat transport to the glacier bed, contributing another mechanism to increased flow.  
5 While the limit for fracture propagation on the Greenland ice sheet is at ~1600 m elevation  
6 (Poinar et al., 2015), with current maximum lake depths ~1200 m, this still leaves  
7 considerable room for lakes to exert their influence on the ice sheet. Thus, although only  
8 transiently storing water, supraglacial lakes still have an important role to play in the evolving  
9 Greenland supraglacial hydrological system.

Unknown  
Field Code Changed

Unknown  
Field Code Changed

## 10 | 6 Conclusion

11 Examination of the evolution of water storage on the surface of ice sheets and glaciers is  
12 important for understanding mass balance, dynamics, and heat transport throughout the ice  
13 mass. In this study, in situ data were used to test the capability of Landsat 8's OLI to estimate  
14 supraglacial lake depth. Promising methods were applied to two sets of Landsat observations.  
15 Patterns of water storage were similar from the two methods, but a factor of two difference  
16 was calculated for the total water volume. WorldView DEMs were used to assess which of  
17 the methods was most accurate. The best method identified for Landsat 8 OLI was an average  
18 of the depth derived from single-band physically-based retrievals of Band 4 (red) and Band 8  
19 (panchromatic); the mean difference between spectrally-derived and DEM-derived lake  
20 depths is only  $0.0 \pm 1.6$  m, showing no bias but some spread. Therefore, this method is  
21 recommended for future lake depth retrievals with OLI, especially for regional studies. This is  
22 the first time supraglacial lake depths have been validated across multiple dates and regions.

Allen Pope 26/10/2015 16:53  
**Deleted:** For all sensors, wavelengths, and input parameters, an important consideration for reflectance-derived lake depth is the atmospheric correction used to prepare the multispectral imagery. All imagery is processed to TOA reflectance, which means that there is some extraneous path radiance remnant in the data. Therefore, TOA values will slightly overestimate the true reflectance. This offset will not be the same between bands, and will influence the retrieved lake depths as discussed below. - ... [9]

23 Discrepancies between spectrally- and DEM-derived depths appear to be explained by  
24 differences between lab-measured and in situ-derived water absorption coefficients ( $g$ ). The  
25 success of other sensors and bands in deriving supraglacial lake depth can thus be inferred  
26 from these  $g$  values. With this insight, multispectral lake depth estimates in the literature were  
27 revisited. Lake extent studies can now be expanded to include lake volume with higher  
28 confidence. Updated  $g$  values are provided (see Tables 1 and S1), but further in situ data  
29 collection and satellite-based studies are needed to build more robust methods.

Allen Pope 21/10/2015 09:24  
**Deleted:** in the discussion

30 The recommended depth retrieval method was applied to all available Landsat 8 imagery for  
31 summer 2014 for the Sermeq Kujalleq (Jakobshavn) region of west Greenland. Seasonal and  
32 regional trends in lake depth ([deepening and then shallowing](#)), evolution ([proceeding](#)

1 [inland/up-glacier and northwards through the summer](#)), and distribution (~300 m to ~2100 m  
2 [a.s.l.](#)) were observed. [At most, lakes contain a similar magnitude of water to supraglacial](#)  
3 [streams, but this may not be true for other parts of Greenland. Both elevation \(and relatedly,](#)  
4 [accumulation / melt forcing\) and surface topography play a role in lake formation and extent,](#)  
5 [behavior that we expect to be modified but observable in other regions.](#) Further work moving  
6 forward will need to contextualize Landsat data with other remote sensing imagery,  
7 fieldwork, and model outputs.

## 8 **Acknowledgements**

9 A. Pope and T. Scambos were supported by USGS contract G12PC00066. D. Shean was  
10 supported by NASA grant NNX12AN36H for this work. An award from the NASA NH  
11 Space Grant supported a collaboration visit between A. Pope and M. Willis. WorldView  
12 imagery was provided by the Polar Geospatial Center at the University of Minnesota, which is  
13 supported by grant ANT-1043681 from the US National Science Foundation. [M. Willis was](#)  
14 [supported by NSF ARC-1111882.](#) We thank the University of North Carolina at Chapel Hill  
15 Research Computing group for providing computational resources that have contributed to  
16 these research results. Non-commercial software tools used in this study included Plot.ly,  
17 ImageJ, and QGIS. [Publication of this article was funded by the University of Colorado](#)  
18 [Boulder Libraries Open Access Fund.](#)

## 20 **References**

- 21 [Arnold, N. S., Banwell, A. F. and Willis, I. C.: High-resolution modelling of the seasonal](#)  
22 [evolution of surface water storage on the Greenland Ice Sheet, The Cryosphere, 8\(4\), 1149–](#)  
23 [1160, doi:10.5194/tc-8-1149-2014, 2014.](#)
- 24 [Banwell, A. F., MacAyeal, D. R. and Sergienko, O. V.: Breakup of the Larsen B Ice Shelf](#)  
25 [triggered by chain reaction drainage of supraglacial lakes, Geophys. Res. Lett., 40\(22\), 5872–](#)  
26 [5876, doi:10.1002/2013GL057694, 2013.](#)
- 27 [Banwell, A. F., Caballero, M., Arnold, N. S., Glasser, N., Cathles, L. M. and MacAyeal, D.](#)  
28 [R.: Supraglacial lakes on the Larsen B Ice Shelf, Antarctica, and at Paakitsoq, W. Greenland:](#)  
29 [a comparative study, Ann. Glaciol., 55\(66\), 1–8, 2014.](#)
- 30 [Barsi, J. A., Lee, K., Kvaran, G., Markham, B. L. and Pedelty, J. A.: The Spectral Response](#)  
31 [of the Landsat-8 Operational Land Imager, Remote Sens., 6\(10\), 10232–10251,](#)  
32 [doi:10.3390/rs61010232, 2014.](#)
- 33 [Berk, A., Anderson, G. P., Acharya, P. K., Bernstein, L. S., Muratov, L., Lee, J., Fox, M.,](#)  
34 [Adler-Golden, S. M., Chetwynd, J. H., Hoke, M. L., Lockwood, R. B., Gardner, J. A., Cooley,](#)

Allen Pope 26/10/2015 16:54

**Moved down [1]:** Publication of this article was funded by the University of Colorado Boulder Libraries Open Access Fund.

Allen Pope 26/10/2015 16:54

**Moved (insertion) [1]**

Allen Pope 26/10/2015 16:54

**Deleted:**

Allen Pope 26/10/2015 16:17

**Formatted:** Bibliography, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Unknown

**Field Code Changed**

Allen Pope 21/10/2015 07:59

**Deleted:** Arnold, N. S., Banwell, A. F. and Willis, I. C.: High-resolution modelling of the seasonal evolution of surface water storage on the Greenland Ice Sheet, The Cryosphere, 8(4), 1149–1160, doi:10.5194/tc-8-1149-2014, 2014. ... [10]

Allen Pope 26/10/2015 16:17

**Formatted:** Font color: Auto

- 1 [T. W., Borel, C. C. and Lewis, P. E.: MODTRAN 5: a reformulated atmospheric band model](#)  
2 [with auxiliary species and practical multiple scattering options: update, edited by S. S. Shen](#)  
3 [and P. E. Lewis, pp. 662–667., 2005.](#)
- 4 [Borbas, E. E., Seemann, S. W., Kern, A., Moy, L., Li, J., Gumley, L. and Menzel, W. P.:](#)  
5 [MODIS atmospheric profile retrieval algorithm theoretical basis document., \[online\]](#)  
6 [Available from: \[http://modis-atmos.gsfc.nasa.gov/docs/MOD07\\\_atbd\\\_v7\\\_April2011.pdf\]\(http://modis-atmos.gsfc.nasa.gov/docs/MOD07\_atbd\_v7\_April2011.pdf\),](#)  
7 [2011.](#)
- 8 [Box, J. E. and Ski, K.: Remote sounding of Greenland supraglacial melt lakes: implications](#)  
9 [for subglacial hydraulics, \*J. Glaciol.\*, 53\(181\), 257–265, 2007.](#)
- 10 [Van den Broeke, M. R., Smeets, C. J. P. P. and van de Wal, R. S. W.: The seasonal cycle and](#)  
11 [interannual variability of surface energy balance and melt in the ablation zone of the west](#)  
12 [Greenland ice sheet, \*The Cryosphere\*, 5\(2\), 377–390, doi:10.5194/tc-5-377-2011, 2011.](#)
- 13 [Chu, V. W.: Greenland ice sheet hydrology A review, \*Prog. Phys. Geogr.\*, 38\(1\), 19–54,](#)  
14 [doi:10.1177/0309133313507075, 2014.](#)
- 15 [Das, S. B., Joughin, I., Behn, M. D., Howat, I. M., King, M. A., Lizarralde, D. and Bhatia, M.](#)  
16 [P.: Fracture Propagation to the Base of the Greenland Ice Sheet During Supraglacial Lake](#)  
17 [Drainage, \*Science\*, 320\(5877\), 778–781, doi:10.1126/science.1153360, 2008.](#)
- 18 [Dierssen, H. M., Zimmerman, R. C., Leathers, R. A., Downes, T. V. and Davis, C. O.: Ocean](#)  
19 [color remote sensing of seagrass and bathymetry in the Bahamas Banks by high-resolution](#)  
20 [airborne imagery, \*Limnol. Oceanogr.\*, 48\(1part2\), 444–455,](#)  
21 [doi:10.4319/lo.2003.48.1\\_part\\_2.0444, 2003.](#)
- 22 [DigitalGlobe: Geolocation Accuracy of WorldView Products, \[online\] Available from:](#)  
23 [https://www.digitalglobe.com/sites/default/files/WorldView\\_Geolocation\\_Accuracy.pdf](https://www.digitalglobe.com/sites/default/files/WorldView_Geolocation_Accuracy.pdf),  
24 [2014.](#)
- 25 [Fitzpatrick, A. A. W., Hubbard, A. L., Box, J. E., Quincey, D. J., van As, D., Mikkelsen, A. P.](#)  
26 [B., Doyle, S. H., Dow, C. F., Hasholt, B. and Jones, G. A.: A decade of supraglacial lake](#)  
27 [volume estimates across a land-terminating margin of the Greenland Ice Sheet, \*Cryosphere\*](#)  
28 [Discuss., 7\(2\), 1383–1414, doi:10.5194/tcd-7-1383-2013, 2013.](#)
- 29 [Flowers, G. E.: Modelling water flow under glaciers and ice sheets, \*Proc. R. Soc. Lond. Math.\*](#)  
30 [Phys. Eng. Sci., 471\(2176\), 20140907, doi:10.1098/rspa.2014.0907, 2015.](#)
- 31 [Georgiou, S., Shepherd, A., McMillan, M. and Nienow, P.: Seasonal evolution of supraglacial](#)  
32 [lake volume from ASTER imagery, \*Ann. Glaciol.\*, 50\(52\), 95–100, 2009.](#)
- 33 [Glasser, N. and Scambos, T.: A structural glaciological analysis of the 2002 Larsen B ice](#)  
34 [shelf collapse, \*J. Glaciol.\*, 54\(184\), 3–16, 2008.](#)
- 35 [Howat, I. M., de la Peña, S., van Angelen, J. H., Lenaerts, J. T. M. and van den Broeke, M.](#)  
36 [R.: Brief Communication “Expansion of meltwater lakes on the Greenland Ice Sheet,” \*The\*](#)  
37 [Cryosphere, 7\(1\), 201–204, doi:10.5194/tc-7-201-2013, 2013.](#)

1 [Howat, I. M., Negrete, A. and Smith, B. E.: The Greenland Ice Mapping Project \(GIMP\) land](#)  
2 [classification and surface elevation data sets, \*The Cryosphere\*, 8\(4\), 1509–1518,](#)  
3 [doi:10.5194/tc-8-1509-2014, 2014.](#)

4 [Irons, J. R., Dwyer, J. L. and Barsi, J. A.: The next Landsat satellite: The Landsat Data](#)  
5 [Continuity Mission, \*Remote Sens. Environ.\*, 122, 11–21, doi:10.1016/j.rse.2011.08.026, 2012.](#)

6 [Johansson, A. M. and Brown, I. A.: Adaptive Classification of Supra-Glacial Lakes on the](#)  
7 [West Greenland Ice Sheet, \*IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.\*, 6\(4\), 1998–](#)  
8 [2007, doi:10.1109/JSTARS.2012.2233722, 2013.](#)

9 [Joughin, I., Das, S. B., Flowers, G. E., Behn, M. D., Alley, R. B., King, M. A., Smith, B. E.,](#)  
10 [Bamber, J. L., van den Broeke, M. R. and van Angelen, J. H.: Influence of ice-sheet geometry](#)  
11 [and supraglacial lakes on seasonal ice-flow variability, \*The Cryosphere\*, 7\(4\), 1185–1192,](#)  
12 [doi:10.5194/tc-7-1185-2013, 2013.](#)

13 [Koenig, L. S., Lampkin, D. J., Montgomery, L. N., Hamilton, S. L., Turrin, J. B., Joseph, C.](#)  
14 [A., Moutsafa, S. E., Panzer, B., Casey, K. A., Paden, J. D., Leuschen, C. and Gogineni, P.:](#)  
15 [Wintertime storage of water in buried supraglacial lakes across the Greenland Ice Sheet, \*The\*](#)  
16 [Cryosphere, 9\(4\), 1333–1342, doi:10.5194/tc-9-1333-2015, 2015.](#)

17 [Lampkin, D. J. and VanderBerg, J.: A preliminary investigation of the influence of basal and](#)  
18 [surface topography on supraglacial lake distribution near Jakobshavn Isbrae, western](#)  
19 [Greenland, \*Hydrol. Process.\*, 25\(21\), 3347–3355, doi:10.1002/hyp.8170, 2011.](#)

20 [Lampkin, D. J. and VanderBerg, J.: Supraglacial melt channel networks in the Jakobshavn](#)  
21 [Isbræ region during the 2007 melt season, \*Hydrol. Process.\*, 28\(25\), 6038–6053,](#)  
22 [doi:10.1002/hyp.10085, 2014.](#)

23 [Leeson, A., Shepherd, A., Sundal, A., Johansson, A. M., Selmes, N., Briggs, K. H., Hogg, A.](#)  
24 [E. and Fettweis, X.: A comparison of supraglacial lake observations derived from MODIS](#)  
25 [imagery at the western margin of the Greenland ice sheet, \*J. Glaciol.\*, 59\(218\), 1179–1188,](#)  
26 [2013.](#)

27 [Leeson, A. A., Shepherd, A., Briggs, K., Howat, I., Fettweis, X., Morlighem, M. and Rignot,](#)  
28 [E.: Supraglacial lakes on the Greenland ice sheet advance inland under warming climate, \*Nat.\*](#)  
29 [Clim. Change, 5\(1\), 51–55, doi:10.1038/nclimate2463, 2015.](#)

30 [Legleiter, C. J. and Overstreet, B. T.: Mapping gravel bed river bathymetry from space, \*J.\*](#)  
31 [Geophys. Res. Earth Surf., 117\(F4\), F04024, doi:10.1029/2012JF002539, 2012.](#)

32 [Legleiter, C. J., Roberts, D. A. and Lawrence, R. L.: Spectrally based remote sensing of river](#)  
33 [bathymetry, \*Earth Surf. Process. Landf.\*, 34\(8\), 1039–1059, doi:10.1002/esp.1787, 2009.](#)

34 [Legleiter, C. J., Tedesco, M., Smith, L. C., Behar, A. E. and Overstreet, B. T.: Mapping the](#)  
35 [bathymetry of supraglacial lakes and streams on the Greenland ice sheet using field](#)  
36 [measurements and high-resolution satellite images, \*The Cryosphere\*, 8\(1\), 215–228,](#)  
37 [doi:10.5194/tc-8-215-2014, 2014.](#)

38 [Liang, Y.-L., Colgan, W., Lv, Q., Steffen, K., Abdalati, W., Stroeve, J., Gallaher, D. and](#)  
39 [Bayou, N.: A decadal investigation of supraglacial lakes in West Greenland using a fully](#)

1 [automatic detection and tracking algorithm, Remote Sens. Environ., 123, 127–138,](#)  
2 [doi:10.1016/j.rse.2012.03.020, 2012.](#)

3 [Lillesand, T., Kiefer, R. W. and Chipman, J.: Remote Sensing and Image Interpretation, 6](#)  
4 [edition., Wiley, Hoboken, NJ., 2007.](#)

5 [Mitchell, G.: PhotoSat WorldView-2 Stereo Satellite DEM Comparison to a LiDAR DEM](#)  
6 [over the Garlock Fault in Southeast California, 2010.](#)

7 [Moratto, Z. M., Broxton, M. J., Beyer, R. A., Lundy, M. and Husmann, K.: Ames Stereo](#)  
8 [Pipeline, NASA's Open Source Automated Stereogrammetry Software, vol. 41, p. 2364,](#)  
9 [\[online\] Available from: <http://adsabs.harvard.edu/abs/2010LPI...41.2364M> \(Accessed 5](#)  
10 [March 2014\), 2010.](#)

11 [Morriss, B. F., Hawley, R. L., Chipman, J. W., Andrews, L. C., Catania, G. A., Hoffman, M.](#)  
12 [J., Lüthi, M. P. and Neumann, T. A.: A ten-year record of supraglacial lake evolution and](#)  
13 [rapid drainage in West Greenland using an automated processing algorithm for multispectral](#)  
14 [imagery, The Cryosphere, 7\(6\), 1869–1877, doi:10.5194/tc-7-1869-2013, 2013.](#)

15 [Moussavi, M., Abdalati, W., Pope, A., Scambos, T., Tedesco, M., Macferrin, M. and Grigsby,](#)  
16 [S.: Derivation and validation of supraglacial lake volumes on the Greenland Ice Sheet from](#)  
17 [high-resolution satellite imagery, Remote Sens. Environ., in review.](#)

18 [Parizek, B. R. and Alley, R. B.: Implications of increased Greenland surface melt under](#)  
19 [global-warming scenarios: ice-sheet simulations, Quat. Sci. Rev., 23\(9–10\), 1013–1027,](#)  
20 [doi:10.1016/j.quascirev.2003.12.024, 2004.](#)

21 [Phillips, T., Rajaram, H., Colgan, W., Steffen, K. and Abdalati, W.: Evaluation of cryo-](#)  
22 [hydrologic warming as an explanation for increased ice velocities in the wet snow zone,](#)  
23 [Sermeq Avannarleq, West Greenland, J. Geophys. Res. Earth Surf., 118\(3\), 1241–1256,](#)  
24 [doi:10.1002/jgrf.20079, 2013.](#)

25 [Philpot, W. D.: Bathymetric mapping with passive multispectral imagery, Appl. Opt., 28\(8\),](#)  
26 [1569–1578, 1989.](#)

27 [Poinar, K., Joughin, I., Das, S. B., Behn, M. D., Lenaerts, J. T. M. and van den Broeke, M. R.:](#)  
28 [Limits to future expansion of surface-melt-enhanced ice flow into the interior of western](#)  
29 [Greenland, Geophys. Res. Lett., 42\(6\), 2015GL063192, doi:10.1002/2015GL063192, 2015.](#)

30 [Pope, A.: Reproducibly Estimating and Evaluating Supraglacial Lake Depth With Landsat 8](#)  
31 [and Other Multispectral Sensors, Earth Space Sci., in review.](#)

32 [Pope, A. and Rees, W. G.: Impact of spatial, spectral, and radiometric properties of](#)  
33 [multispectral imagers on glacier surface classification, Remote Sens. Environ., 141, 1–13,](#)  
34 [doi:10.1016/j.rse.2013.08.028, 2014a.](#)

35 [Pope, A. and Rees, W. G.: Using in situ Spectra to Explore Landsat Classification of Glacier](#)  
36 [Surfaces, J. Appl. Earth Obs. Geoinformation, 27A, 42–52, doi:10.1016/j.jag.2013.08.007,](#)  
37 [2014b.](#)

- 1 [Pope, R. M. and Fry, E. S.: Absorption spectrum ~380–700 nm! of pure water. II. Integrating](#)  
2 [cavity measurements, \*Appl. Opt.\*, 36\(33\), 8710–8723, 1997.](#)
- 3 [Roy, D. P., Wulder, M. A., Loveland, T. R., C.E., W., Allen, R. G., Anderson, M. C., Helder,](#)  
4 [D., Irons, J. R., Johnson, D. M., Kennedy, R., Scambos, T. A., Schaaf, C. B., Schott, J. R.,](#)  
5 [Sheng, Y., Vermote, E. F., Belward, A. S., Bindschadler, R., Cohen, W. B., Gao, F., Hipple,](#)  
6 [J. D., Hostert, P., Huntington, J., Justice, C. O., Kilic, A., Kovalskyy, V., Lee, Z. P.,](#)  
7 [Lymburner, L., Masek, J. G., McCorkel, J., Shuai, Y., Trezza, R., Vogelmann, J., Wynne, R.](#)  
8 [H. and Zhu, Z.: Landsat-8: Science and product vision for terrestrial global change research,](#)  
9 [Remote Sens. Environ., 145, 154–172, doi:10.1016/j.rse.2014.02.001, 2014.](#)
- 10 [Selmes, N., Murray, T. and James, T. D.: Fast draining lakes on the Greenland Ice Sheet,](#)  
11 [Geophys. Res. Lett., 38\(15\), L15501, doi:10.1029/2011GL047872, 2011.](#)
- 12 [Seo, K.-W., Waliser, D. E., Lee, C.-K., Tian, B., Scambos, T., Kim, B.-M., van Angelen, J. H.](#)  
13 [and van den Broeke, M. R.: Accelerated Mass Loss from Greenland Ice Sheet: Links to](#)  
14 [Atmospheric Circulation in the North Atlantic, \*Glob. Planet. Change,\*](#)  
15 [doi:10.1016/j.gloplacha.2015.02.006, 2015.](#)
- 16 [Shean, D. E., Alexandrov, O., Moratto, Z. M., Smith, B. E., Joughin, I. R., Porter, C. C. and](#)  
17 [Morin, P. J.: An automated, open-source pipeline for mass production of digital elevation](#)  
18 [models \(DEMs\) from very-high-resolution commercial stereo satellite imagery, \*ISPRS J.\*](#)  
19 [Photogramm. Remote Sens., in press, 2015.](#)
- 20 [Smith, L. C., Chu, V. W., Yang, K., Gleason, C. J., Pitcher, L. H., Rennermalm, A. K.,](#)  
21 [Legleiter, C. J., Behar, A. E., Overstreet, B. T., Moustafa, S. E., Tedesco, M., Forster, R. R.,](#)  
22 [LeWinter, A. L., Finnegan, D. C., Sheng, Y. and Balog, J.: Efficient meltwater drainage](#)  
23 [through supraglacial streams and rivers on the southwest Greenland ice sheet, \*Proc. Natl.\*](#)  
24 [Acad. Sci., 112\(4\), 1001–1006, doi:10.1073/pnas.1413024112, 2015.](#)
- 25 [Smith, R. C. and Baker, K. S.: Optical properties of the clearest natural waters \(200–800 nm\),](#)  
26 [Appl. Opt., 20\(2\), 177–184, 1981.](#)
- 27 [Sneed, W. A. and Hamilton, G.: Evolution of melt pond volume on the surface of the](#)  
28 [Greenland Ice Sheet, \*Geophys. Res. Lett.\*, 34, L03501, doi:10.1029/2006GL028697, 2007.](#)
- 29 [Sneed, W. A. and Hamilton, G. S.: Validation of a method for determining the depth of](#)  
30 [glacial melt ponds using satellite imagery, \*Ann. Glaciol.\*, 52\(59\), 15–22, 2011.](#)
- 31 [Stevens, L. A., Behn, M. D., McGuire, J. J., Das, S. B., Joughin, I., Herring, T., Shean, D. E.](#)  
32 [and King, M. A.: Greenland supraglacial lake drainages triggered by hydrologically induced](#)  
33 [basal slip, \*Nature\*, 522\(7554\), 73–76, doi:10.1038/nature14480, 2015.](#)
- 34 [Stumpf, R. P., Holderied, K. and Sinclair, M.: Determination of water depth with high-](#)  
35 [resolution satellite imagery over variable bottom types, \*Limnol. Oceanogr.\*, 48\(1part2\), 547–](#)  
36 [556, doi:10.4319/lo.2003.48.1\\_part\\_2.0547, 2003.](#)
- 37 [Sundal, A. V., Shepherd, A., Nienow, P., Hanna, E., Palmer, S. and Huybrechts, P.: Melt-](#)  
38 [induced speed-up of Greenland ice sheet offset by efficient subglacial drainage, \*Nature\*,](#)  
39 [469\(7331\), 521–524, doi:10.1038/nature09740, 2011.](#)

1 [Tedesco, M. and Steiner, N.: In-situ multispectral and bathymetric measurements over a](#)  
2 [supraglacial lake in western Greenland using a remotely controlled watercraft, \*The\*](#)  
3 [Cryosphere](#), 5(2), 445–452, 2011.

4 [Tedesco, M., Lüthje, M., Steffen, K., Steiner, N., Fettweis, X., Willis, I., Bayou, N. and](#)  
5 [Banwell, A.: Measurement and modeling of ablation of the bottom of supraglacial lakes in](#)  
6 [western Greenland, \*Geophys. Res. Lett.\*](#), 39, L02502, doi:201210.1029/2011GL049882, 2012.

7 [Tedesco, M., Willis, I. C., Hoffman, M. J., Banwell, A. F., Alexander, P. and Arnold, N. S.:](#)  
8 [Ice dynamic response to two modes of surface lake drainage on the Greenland ice sheet,](#)  
9 [Environ. Res. Lett.](#), 8(3), 034007, doi:10.1088/1748-9326/8/3/034007, 2013.

10 [Tedesco, M., Steiner, N. and Pope, A.: In situ spectral reflectance and depth of a supraglacial](#)  
11 [lake in Greenland, UCAR-NCAR CISL ACADIS](#), doi:10.5065/D6FQ9TN2, 2015.

12 [USGS: Using the USGS Landsat 8 Product, \[online\] Available from:](#)  
13 [http://landsat.usgs.gov/Landsat8\\_Using\\_Product.php](http://landsat.usgs.gov/Landsat8_Using_Product.php) (Accessed 4 April 2015), 2013.

14 [Willis, M. J., Herried, B. G., Bevis, M. G. and Bell, R. E.: Recharge of a subglacial lake by](#)  
15 [surface meltwater in northeast Greenland, \*Nature\*](#), 518(7538), 223–227,  
16 [doi:10.1038/nature14116](#), 2015.

17 [Zwally, H. J., Abdalati, W., Herring, T., Larson, K. M., Saba, J. L. and Steffen, K.: Surface](#)  
18 [Melt-Induced Acceleration of Greenland Ice-Sheet Flow, \*Science\*](#), 297, 218–222, 2002.

19  
20

1 Table 1. Laboratory-based and in situ-derived water absorption coefficients for lake depth  
 2 estimation using the physically based method ( $g$ , see Eqn. 1) and empirical method ( $a$ ,  $b$ , and  
 3  $c$ , see Eqns. 5-6). Regression statistics (correlation coefficient and root mean squared error)  
 4 for lake depth estimates using field spectra convolved to emulate multispectral bands are also  
 5 included. Asterisks indicate the methods applied to OLI data in this paper. Bold text indicates  
 6 recommended bands for lake depth estimation with OLI. See Table S1 for results from other  
 7 multispectral sensors.

8


Satellite & Band	Lab-based $g$ ( $m^{-1}$ )	Regressed $g$ ( $m^{-1}$ )	$r$	RMSE (m)
OLI 1 (coastal)	0.0178	0.0093	0.0494	11.03
OLI 2 (blue)	0.0341	0.025	0.2886	3.10
*OLI 3 (green)	0.1413	0.01	0.7842	0.78
<b>*OLI 4 (red)</b>	<b>0.7507</b>	<b>0.80</b>	<b>0.9624</b>	<b>0.28</b>
<b>*OLI 8 (panchromatic)</b>	<b>0.3817</b>	<b>0.36</b>	<b>0.8422</b>	<b>0.63</b>
ETM+ 1 (blue) Gain H	0.0334	0.03	0.2626	3.34
ETM+ 1 (blue) Gain L	0.0334	0.03	0.2625	3.34
ETM+ 2 (green) Gain H	0.1665	0.15	0.7892	0.77
ETM+ 2 (green) Gain L	0.1665	0.14	0.7890	0.77
ETM+ 3 (red) Gain H	0.8049	0.83	0.9548	0.31
ETM+ 3 (red) Gain L	0.8049	0.83	0.9412	0.37
OLI 1 & 2 (coastal & blue)	-	-	0.7871	2.57
OLI 1 & 3 (coastal & green)	-	-	0.9208	1.10
OLI 1 & 4 (coastal & red)	-	-	0.8987	1.34
*OLI 2 & 3 (blue & green)	-	-	0.9401	0.88

OLI 2 & 3 (blue & red)	-	-	0.8885	1.41
OLI 3 & 4 (green & red)	-	-	0.6063	1.74

1

Satellite & Bands	<i>a</i>	<i>b</i>	<i>c</i>	<i>r</i>	RMSE (m)
OLI 3 & 4 (green & red)	-13.8398	40.0344	-23.4057	0.4537	0.89
OLI 2 & 4 (blue & red)	3.4414	-9.0500	7.8243	0.8610	0.51
OLI 1 & 2 (coastal & blue)	0.9750	18.1837	145.7811	0.8031	0.59
*OLI 1 & 3 (coastal & green)	0.1488	5.0370	5.0473	0.9228	0.38
OLI 1 & 4 (coastal & red)	4.8374	-11.2317	8.2001	0.8964	0.44
*OLI 1 & 8 (coastal & pan)	1.6240	-5.9696	12.4983	0.9473	0.32
ETM+ 2 & 3 (green & red) L	1.4794	-3.2173	2.8860	0.8855	0.46
ETM+ 2 & 3 (green & red) H	2.3102	-4.4616	3.2802	0.8970	0.44
ETM+ 1 & 3 (blue & red) L	4.0925	-5.3290	2.4296	0.9655	0.26
ETM+ 1 & 3 (blue & red) H	4.2825	-5.4754	2.4225	0.9694	0.24


2



1  
2  
3  
4  
5  
6  
7  
8  
9  
10

Figure 1. Regional map showing the two study regions for lake depth estimation using Landsat 8 OLI imagery. The northwest Greenland study region is identified with a single box indicating a subszene area. The Sermeq Kujalleq (Jakobshavn) study region shows WRS-2 path/row outlines for Landsat scenes color-coded and dashed to indicate the mean latitude and average elevation of ice within the scenes (see Sect. 4.4 and Table S2). The background is elevation from the Greenland Ice Mapping Project (GIMP) DEM, courtesy BPRC Glacier Dynamics Research Group, Ohio State University (Howat et al., 2014).


Unknown  
Field Code Changed



1  
2  
3  
4  
5  
6  
7

Figure 2. Regression plots for in situ measured reflectance spectra used to emulate Landsat OLI and ETM+ reflectance and sonar-measured depths, including OLI single band (a), ETM+ low gain single band (b), OLI coastal and panchromatic (c), and OLI coastal and green (d). Statistics for all regressions are reported in Table 1.

1




2

3


4 Figure 3. Total water volume stored in supraglacial lakes in the northwest Greenland study  
5 region for the summer of 2014. Based on analysis, “Band Average 4 & 8” is likely to be the  
6 most accurate (see Fig. 4).

7



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12

Figure 4. Statistics for the difference in supraglacial lake depth from physically-based and empirical methods derived from OLI imagery and WorldView DEMs, including mean/standard deviation (solid lines) and median/quartiles (dotted lines). An average of the Band 4 and Band 8 methods is used for our mapping (Figures 5 and 6). The method showing the least bias and lowest errors is an average of Band 4 (red) and Band 8 (panchromatic) single band physically based retrievals, with a mean offset of  $0.0 \pm 1.6$  m (as indicated by the bar at the bottom of the diagram). Discrepancies in lake depth estimation for physically based retrievals can be traced to differences between lab-measured and in situ-regressed water absorption coefficients (see Table 1).




1

2

3 Figure 5. Total water stored in supraglacial lakes over the 2014 summer using single Landsat  
 4 8 scenes covering the Sermeq Kujalleq (Jakobshavn) region (see Fig. 1, Table S1). All scenes  
 5 are shown together in (a). (b) shows only the low elevation, coastal scenes, demonstrating  
 6 delayed lake formation at higher latitudes. (c) shows both elevation and latitude effects in  
 7 driving supraglacial water storage for scenes in WRS-2 path 8. (d) shows latitude and  
 8 elevation effects for scenes in WRS-2 path 6. All sub-figures are on the same grid as part (a).

9



1

2

3 Figure 6. Statistics of lake depth and elevation distribution for all Sermeq Kujalleq  
 4 (Jakobshavn) region 2014 Landsat OLI imagery (see Table S2). A histogram of lake depths  
 5 (a) shows many shallow lakes, many lakes reaching depths of ~3 m, and a maximum lake  
 6 depth of ~6 m. Lakes depth by pixel distribution with elevation is shown in (b), the  
 7 hypsometry of maximum supraglacial lake depths on this region of the Greenland Ice Sheet as  
 8 determined from one, m bins of the GIMP DEM (Howat et al., 2014).

Allen Pope 21/10/2015 09:57  
 Deleted: 1  
 Unknown  
 Field Code Changed

<b>Page 2: [1] Deleted</b>	<b>Allen Pope</b>	<b>21/10/2015 08:01</b>
----------------------------	-------------------	-------------------------

Supraglacial lakes are also important for surface heat fluxes by storing latent heat near the surface of the ice sheet. In addition, lakes reduce surface albedo, hence further promoting melt through a positive feedback mechanism (Leeson et al., 2015).

<b>Page 8: [2] Change</b>	<b>Unknown</b>
---------------------------	----------------

Field Code Changed

<b>Page 8: [3] Deleted</b>	<b>Allen Pope</b>	<b>21/10/2015 17:26</b>
----------------------------	-------------------	-------------------------

(Shean et al., 2015; Moratto et al., 2010)

<b>Page 8: [4] Change</b>	<b>Unknown</b>
---------------------------	----------------

Field Code Changed

<b>Page 8: [5] Change</b>	<b>Unknown</b>
---------------------------	----------------

Field Code Changed

<b>Page 8: [6] Change</b>	<b>Unknown</b>
---------------------------	----------------

Field Code Changed

<b>Page 8: [7] Deleted</b>	<b>David Shean</b>	<b>21/10/2015 11:17</b>
----------------------------	--------------------	-------------------------

per-pixel standard deviation of elevations for the stack are

<b>Page 8: [8] Change</b>	<b>Unknown</b>
---------------------------	----------------

Field Code Changed

<b>Page 17: [9] Deleted</b>	<b>Allen Pope</b>	<b>26/10/2015 16:53</b>
-----------------------------	-------------------	-------------------------

For all sensors, wavelengths, and input parameters, an important consideration for reflectance-derived lake depth is the atmospheric correction used to prepare the multispectral imagery. All imagery is processed to TOA reflectance, which means that there is some extraneous path radiance remnant in the data. Therefore, TOA values will slightly overestimate the true reflectance. This offset will not be the same between bands, and will influence the retrieved lake depths as discussed below.

The single band physically based model requires that the reflectance of optically deep water be derived for each scene separately. Effectively, this shifts the exponential decay curve of light in lake water but does not change its shape. Therefore, as long as path

radiance is assumed to be homogeneous across the 185-km wide Landsat scene, TOA reflectance is sufficient for lake depth estimation. To test this assumption, the MODTRAN radiative transfer model (Berk et al., 2005) was used to simulate path radiance on a day for which Landsat data were used in northwest Greenland (18 July 2013) to investigate variations associated with variable water vapor and ozone across a Landsat scene. According to MODIS retrievals (accurate to 30 DU; Borbas et al., 2011), ozone varies within a Landsat scene on the order of approximately  $\pm 50$  DU, which translates to a path radiance of  $\pm 1.6\%$  in the red channel. For lake depth, this can propagate on to a  $\sim 20\%$  error in lake depth. However, because much of this error appears largely random (for a given point in time and space, although coherent patterns on a given day), although it decreases confidence in individual lake depth retrievals (and likely contributes to the  $\pm 1.6$  m spread seen with WorldView validation of lake depths), averaged water volume retrieval should not be biased. For water vapor there was a 0.3% change in path radiance between the minimum and maximum Landsat scene values, making it a small contributor to overall error. Between days, however, path radiance effects due to water vapor may vary by an order of magnitude more.

For the multiple band methods, the differential change in path radiance has larger effects. Sensitivity tests showed that a 3% change in path radiance for one or both bands changed water volumes on the order of 10-30%. Therefore, a more rigorous atmospheric correction is necessary in order to apply multi-band lake depth algorithms. Still, for the study here, because validation is conducted across 12 non-consecutive days in both spring and autumn, we do not expect atmospheric conditions to bias our conclusions.

There are additional limitations to our method. As discussed above, OLI lake depth estimates (average single-band estimates from red and panchromatic bands) are robust for regional averages but not single pixels. In addition, the threshold used to identify lake extent may need to be adjusted for different regions and scenes (e.g. Banwell et al., 2014; Box and Ski, 2007). Lake depth retrievals are also sensitive to variations in ice albedo, as well as to the presence of ice lids on the surface of supraglacial lakes, which can be common in both early and late summer. Cloud cover and Landsat's 16-day revisit time also limit the conclusions that can be drawn from OLI lake depths. Many studies have used daily MODIS data to identify and track supraglacial lakes (e.g. Liang et al., 2012;

Selmes et al., 2011; Sundal et al., 2011). Fusing the higher temporal resolution of MODIS (or additional sensors such as ESA's upcoming Sentinel-2) and higher spatial resolution of Landsat, along with more in situ calibration and validation data, should lead to unique insights to supraglacial water storage.

Arnold, N. S., Banwell, A. F. and Willis, I. C.: High-resolution modelling of the seasonal evolution of surface water storage on the Greenland Ice Sheet, *The Cryosphere*, 8(4), 1149–1160, doi:10.5194/tc-8-1149-2014, 2014.

Banwell, A. F., MacAyeal, D. R. and Sergienko, O. V.: Breakup of the Larsen B Ice Shelf triggered by chain reaction drainage of supraglacial lakes, *Geophys. Res. Lett.*, 40(22), 5872–5876, doi:10.1002/2013GL057694, 2013.

Banwell, A. F., Caballero, M., Arnold, N. S., Glasser, N., Cathles, L. M. and MacAyeal, D. R.: Supraglacial lakes on 1 the Larsen B Ice Shelf, Antarctica, and at Paakitsoq, W. Greenland: a comparative study, *Ann. Glaciol.*, 55(66), 1–8, 2014.

Barsi, J. A., Lee, K., Kvaran, G., Markham, B. L. and Pedelty, J. A.: The Spectral Response of the Landsat-8 Operational Land Imager, *Remote Sens.*, 6(10), 10232–10251, doi:10.3390/rs61010232, 2014.

Berk, A., Anderson, G. P., Acharya, P. K., Bernstein, L. S., Muratov, L., Lee, J., Fox, M., Adler-Golden, S. M., Chetwynd, J. H., Hoke, M. L., Lockwood, R. B., Gardner, J. A., Cooley, T. W., Borel, C. C. and Lewis, P. E.: MODTRAN 5: a reformulated atmospheric band model with auxiliary species and practical multiple scattering options: update, edited by S. S. Shen and P. E. Lewis, pp. 662–667., 2005.

Borbas, E. E., Seemann, S. W., Kern, A., Moy, L., Li, J., Gumley, L. and Menzel, W. P.: MODIS atmospheric profile retrieval algorithm theoretical basis document., [online] Available from: [http://modis-atmos.gsfc.nasa.gov/\\_docs/MOD07\\_atbd\\_v7\\_April2011.pdf](http://modis-atmos.gsfc.nasa.gov/_docs/MOD07_atbd_v7_April2011.pdf), 2011.

Box, J. E. and Ski, K.: Remote sounding of Greenland supraglacial melt lakes: implications for subglacial hydraulics, *J. Glaciol.*, 53(181), 257–265, 2007.

Van den Broeke, M. R., Smeets, C. J. P. P. and van de Wal, R. S. W.: The seasonal cycle and interannual variability of surface energy balance and melt in the ablation zone of the west Greenland ice sheet, *The Cryosphere*, 5(2), 377–390, doi:10.5194/tc-5-377-2011, 2011.

Chu, V. W.: Greenland ice sheet hydrology A review, *Prog. Phys. Geogr.*, 38(1), 19–54, doi:10.1177/0309133313507075, 2014.

Das, S. B., Joughin, I., Behn, M. D., Howat, I. M., King, M. A., Lizarralde, D. and Bhatia, M. P.: Fracture Propagation to the Base of the Greenland Ice Sheet During

Supraglacial Lake Drainage, *Science*, 320(5877), 778–781, doi:10.1126/science.1153360, 2008.

Dierssen, H. M., Zimmerman, R. C., Leathers, R. A., Downes, T. V. and Davis, C. O.: Ocean color remote sensing of seagrass and bathymetry in the Bahamas Banks by high-resolution airborne imagery, *Limnol. Oceanogr.*, 48(1part2), 444–455, doi:10.4319/lo.2003.48.1\_part\_2.0444, 2003.

DigitalGlobe: Geolocation Accuracy of WorldView Products, [online] Available from: [https://www.digitalglobe.com/sites/default/files/WorldView\\_Geolocation\\_Accuracy.pdf](https://www.digitalglobe.com/sites/default/files/WorldView_Geolocation_Accuracy.pdf), 2014.

Fitzpatrick, A. A. W., Hubbard, A. L., Box, J. E., Quincey, D. J., van As, D., Mikkelsen, A. P. B., Doyle, S. H., Dow, C. F., Hasholt, B. and Jones, G. A.: A decade of supraglacial lake volume estimates across a land-terminating margin of the Greenland Ice Sheet, *Cryosphere Discuss.*, 7(2), 1383–1414, doi:10.5194/tcd-7-1383-2013, 2013.

Flowers, G. E.: Modelling water flow under glaciers and ice sheets, *Proc. R. Soc. Lond. Math. Phys. Eng. Sci.*, 471(2176), 20140907, doi:10.1098/rspa.2014.0907, 2015.

Glasser, N. and Scambos, T.: A structural glaciological analysis of the 2002 Larsen B ice shelf collapse, *J. Glaciol.*, 54(184), 3–16, 2008.

Howat, I. M., de la Peña, S., van Angelen, J. H., Lenaerts, J. T. M. and van den Broeke, M. R.: Brief Communication “Expansion of meltwater lakes on the Greenland Ice Sheet,” *The Cryosphere*, 7(1), 201–204, doi:10.5194/tc-7-201-2013, 2013.

Howat, I. M., Negrete, A. and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets, *The Cryosphere*, 8(4), 1509–1518, doi:10.5194/tc-8-1509-2014, 2014.

Irons, J. R., Dwyer, J. L. and Barsi, J. A.: The next Landsat satellite: The Landsat Data Continuity Mission, *Remote Sens. Environ.*, 122, 11–21, doi:10.1016/j.rse.2011.08.026, 2012.

Johansson, A. M. and Brown, I. A.: Adaptive Classification of Supra-Glacial Lakes on the West Greenland Ice Sheet, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 6(4), 1998–2007, doi:10.1109/JSTARS.2012.2233722, 2013.

Joughin, I., Das, S. B., Flowers, G. E., Behn, M. D., Alley, R. B., King, M. A., Smith, B. E., Bamber, J. L., van den Broeke, M. R. and van Angelen, J. H.: Influence of ice-sheet geometry and supraglacial lakes on seasonal ice-flow variability, *The Cryosphere*, 7(4), 1185–1192, doi:10.5194/tc-7-1185-2013, 2013.

Lampkin, D. J. and VanderBerg, J.: A preliminary investigation of the influence of basal and surface topography on supraglacial lake distribution near Jakobshavn Isbrae, western Greenland, *Hydrol. Process.*, 25(21), 3347–3355, doi:10.1002/hyp.8170, 2011.

Lampkin, D. J. and VanderBerg, J.: Supraglacial melt channel networks in the Jakobshavn Isbræ region during the 2007 melt season, *Hydrol. Process.*, 28(25), 6038–6053, doi:10.1002/hyp.10085, 2014.

Leeson, A., Shepherd, A., Sundal, A., Johansson, A. M., Selmes, N., Briggs, K. H., Hogg, A. E. and Fettweis, X.: A comparison of supraglacial lake observations derived from MODIS imagery at the western margin of the Greenland ice sheet, *J. Glaciol.*, 59(218), 1179–1188, 2013.

Leeson, A. A., Shepherd, A., Briggs, K., Howat, I., Fettweis, X., Morlighem, M. and Rignot, E.: Supraglacial lakes on the Greenland ice sheet advance inland under warming climate, *Nat. Clim. Change*, 5(1), 51–55, doi:10.1038/nclimate2463, 2015.

Legleiter, C. J. and Overstreet, B. T.: Mapping gravel bed river bathymetry from space, *J. Geophys. Res. Earth Surf.*, 117(F4), F04024, doi:10.1029/2012JF002539, 2012.

Legleiter, C. J., Roberts, D. A. and Lawrence, R. L.: Spectrally based remote sensing of river bathymetry, *Earth Surf. Process. Landf.*, 34(8), 1039–1059, doi:10.1002/esp.1787, 2009.

Legleiter, C. J., Tedesco, M., Smith, L. C., Behar, A. E. and Overstreet, B. T.: Mapping the bathymetry of supraglacial lakes and streams on the Greenland ice sheet using field measurements and high-resolution satellite images, *The Cryosphere*, 8(1), 215–228, doi:10.5194/tc-8-215-2014, 2014.

Liang, Y.-L., Colgan, W., Lv, Q., Steffen, K., Abdalati, W., Stroeve, J., Gallaher, D. and Bayou, N.: A decadal investigation of supraglacial lakes in West Greenland using a fully automatic detection and tracking algorithm, *Remote Sens. Environ.*, 123, 127–138, doi:10.1016/j.rse.2012.03.020, 2012.

Lillesand, T., Kiefer, R. W. and Chipman, J.: *Remote Sensing and Image Interpretation*, 6 edition., Wiley, Hoboken, NJ., 2007.

Mitchell, G.: *PhotoSat WorldView-2 Stereo Satellite DEM Comparison to a LiDAR DEM over the Garlock Fault in Southeast California*, 2010.

Moratto, Z. M., Broxton, M. J., Beyer, R. A., Lundy, M. and Husmann, K.: Ames Stereo Pipeline, NASA's Open Source Automated Stereogrammetry Software, vol. 41, p. 2364. [online] Available from: <http://adsabs.harvard.edu/abs/2010LPI....41.2364M> (Accessed 5 March 2014), 2010.

Morriss, B. F., Hawley, R. L., Chipman, J. W., Andrews, L. C., Catania, G. A., Hoffman, M. J., Lüthi, M. P. and Neumann, T. A.: A ten-year record of supraglacial lake evolution and rapid drainage in West Greenland using an automated processing algorithm for multispectral imagery, *The Cryosphere*, 7(6), 1869–1877, doi:10.5194/tc-7-1869-2013, 2013.

Parizek, B. R. and Alley, R. B.: Implications of increased Greenland surface melt under global-warming scenarios: ice-sheet simulations, *Quat. Sci. Rev.*, 23(9–10), 1013–1027, doi:10.1016/j.quascirev.2003.12.024, 2004.

Phillips, T., Rajaram, H., Colgan, W., Steffen, K. and Abdalati, W.: Evaluation of cryo-hydrologic warming as an explanation for increased ice velocities in the wet snow zone, Sermeq Avannarleq, West Greenland, *J. Geophys. Res. Earth Surf.*, 118(3), 1241–1256, doi:10.1002/jgrf.20079, 2013.

Philpot, W. D.: Bathymetric mapping with passive multispectral imagery, *Appl. Opt.*, 28(8), 1569–1578, 1989.

Poinar, K., Joughin, I., Das, S. B., Behn, M. D., Lenaerts, J. T. M. and van den Broeke, M. R.: Limits to future expansion of surface-melt-enhanced ice flow into the interior of western Greenland, *Geophys. Res. Lett.*, 2015GL063192, doi:10.1002/2015GL063192, 2015.

Pope, A. and Rees, W. G.: Impact of spatial, spectral, and radiometric properties of multispectral imagers on glacier surface classification, *Remote Sens. Environ.*, 141, 1–13, doi:10.1016/j.rse.2013.08.028, 2014a.

Pope, A. and Rees, W. G.: Using in situ Spectra to Explore Landsat Classification of Glacier Surfaces, *J. Appl. Earth Obs. Geoinformation*, 27A, 42–52, doi:10.1016/j.jag.2013.08.007, 2014b.

Pope, R. M. and Fry, E. S.: Absorption spectrum ~380–700 nm! of pure water. II. Integrating cavity measurements, *Appl. Opt.*, 36(33), 8710–8723, 1997.

Roy, D. P., Wulder, M. A., Loveland, T. R., C.E., W., Allen, R. G., Anderson, M. C., Helder, D., Irons, J. R., Johnson, D. M., Kennedy, R., Scambos, T. A., Schaaf, C. B., Schott, J. R., Sheng, Y., Vermote, E. F., Belward, A. S., Bindschadler, R., Cohen, W. B., Gao, F., Hipple, J. D., Hostert, P., Huntington, J., Justice, C. O., Kilic, A., Kovalsky, V., Lee, Z. P., Lymburner, L., Masek, J. G., McCorkel, J., Shuai, Y., Trezza, R., Vogelmann, J., Wynne, R. H. and Zhu, Z.: Landsat-8: Science and product vision for terrestrial global change research, *Remote Sens. Environ.*, 145, 154–172, doi:10.1016/j.rse.2014.02.001, 2014.

Selmes, N., Murray, T. and James, T. D.: Fast draining lakes on the Greenland Ice Sheet, *Geophys. Res. Lett.*, 38(15), n/a–n/a, doi:10.1029/2011GL047872, 2011.

Seo, K.-W., Waliser, D. E., Lee, C.-K., Tian, B., Scambos, T., Kim, B.-M., van Angelen, J. H. and van den Broeke, M. R.: Accelerated Mass Loss from Greenland Ice Sheet: Links to Atmospheric Circulation in the North Atlantic, *Glob. Planet. Change*, doi:10.1016/j.gloplacha.2015.02.006, 2015.

Smith, L. C., Chu, V. W., Yang, K., Gleason, C. J., Pitcher, L. H., Rennermalm, A. K., Legleiter, C. J., Behar, A. E., Overstreet, B. T., Moustafa, S. E., Tedesco, M., Forster, R. R., LeWinter, A. L., Finnegan, D. C., Sheng, Y. and Balog, J.: Efficient meltwater

drainage through supraglacial streams and rivers on the southwest Greenland ice sheet, *Proc. Natl. Acad. Sci.*, 201413024, doi:10.1073/pnas.1413024112, 2015.

Smith, R. C. and Baker, K. S.: Optical properties of the clearest natural waters (200-800 nm), *Appl. Opt.*, 20(2), 177–184, 1981.

Sneed, W. A. and Hamilton, G.: Evolution of melt pond volume on the surface of the Greenland Ice Sheet, *Geophys. Res. Lett.*, 34, L03501, 2007.

Sneed, W. A. and Hamilton, G. S.: Validation of a method for determining the depth of glacial melt ponds using satellite imagery, *Ann. Glaciol.*, 52(59), 15–22, 2011.

Stevens, L., Behn, M. D., McGuire, J., Das, S. B., Joughin, I., Herring, T., Shean, D. and King, M. A.: Hydrologically Induced Basal Slip Triggers Greenland Supraglacial Lake Drainages, San Francisco., 2014.

Stumpf, R. P., Holderied, K. and Sinclair, M.: Determination of water depth with high-resolution satellite imagery over variable bottom types, *Limnol. Oceanogr.*, 48(1part2), 547–556, doi:10.4319/lo.2003.48.1\_part\_2.0547, 2003.

Sundal, A. V., Shepherd, A., Nienow, P., Hanna, E., Palmer, S. and Huybrechts, P.: Melt-induced speed-up of Greenland ice sheet offset by efficient subglacial drainage, *Nature*, 469(7331), 521–524, doi:10.1038/nature09740, 2011.

Tedesco, M. and Steiner, N.: In-situ multispectral and bathymetric measurements over a supraglacial lake in western Greenland using a remotely controlled watercraft, *The Cryosphere*, 5(2), 445–452, 2011.

Tedesco, M., Lüthje, M., Steffen, K., Steiner, N., Fettweis, X., Willis, I., Bayou, N. and Banwell, A.: Measurement and modeling of ablation of the bottom of supraglacial lakes in western Greenland, *Geophys. Res. Lett.*, 39, L02502, doi:10.1029/2011GL049882, 2012.

Tedesco, M., Willis, I. C., Hoffman, M. J., Banwell, A. F., Alexander, P. and Arnold, N. S.: Ice dynamic response to two modes of surface lake drainage on the Greenland ice sheet, *Environ. Res. Lett.*, 8(3), 034007, doi:10.1088/1748-9326/8/3/034007, 2013.

USGS: Using the USGS Landsat 8 Product, [online] Available from: [http://landsat.usgs.gov/Landsat8\\_Using\\_Product.php](http://landsat.usgs.gov/Landsat8_Using_Product.php) (Accessed 4 April 2015), 2013.

Wientjes, I. G. M., Van de Wal, R. S. W., Reichert, G. J., Sluijs, A. and Oerlemans, J.: Dust from the dark region in the western ablation zone of the Greenland ice sheet, *The Cryosphere*, 5(3), 589–601, doi:10.5194/tc-5-589-2011, 2011.

Willis, M. J., Herried, B. G., Bevis, M. G. and Bell, R. E.: Recharge of a subglacial lake by surface meltwater in northeast Greenland, *Nature*, 518(7538), 223–227, doi:10.1038/nature14116, 2015.

