

This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

Glaciers change over the last century, Caucasus Mountains, Georgia, observed by the old topographical maps, Landsat and ASTER satellite imagery

L. G. Tielidze

Department of Geomorphology, Vakhushti Bagrationi Institute of Geography, Ivane Javakhishvili Tbilisi State University, 6 Tamarashvili st. Tbilisi 0177, Georgia

Received: 12 May 2015 – Accepted: 17 June 2015 – Published: 17 July 2015

Correspondence to: L. G. Tielidze (levan.tielidze@tsu.ge)

Published by Copernicus Publications on behalf of the European Geosciences Union.

TC D

9, 3777–3806, 2015

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The study of glaciers in the Caucasus began in the first quarter of the 18th century. The first data on glaciers can be found in the works of great Georgian scientist Vakhushti Bagrationi. After almost hundred years the foreign scientists began to describe the glaciers of Georgia. Information about the glaciers of Georgia can be found in the works of W. Abich (1865), D. Freshfield (1869), G. Radde (1873), N. Dinik (1884), I. Rashevskiy (1904), A. Reinhardt (1916, 1917) etc. The first statistical information about the glaciers of Georgia are found in the catalog of the Caucasus glaciers compiled by K. Podozerskiy in 1911 (Podozerkiy, 1911). Then, in 1960s the large-scale (1 : 25 000, 1 : 50 000) topographic maps were published, which were compiled in 1955–1960 on the basis of the space images. On the basis of the mentioned maps R. Gobejishvili gave quite detailed statistical information about the glaciers of Georgia (Gobejishvili, 1989). Then in 1975 the glaciological catalog of the former USSR was published (The Catalog of Glaciers of the USSR, Vol. 8–9, 1975), where the statistical information about the glaciers of Georgia was obtained on the basis of the space images of 1970–1975. Thus, complete statistical information on the glaciers of Georgia has not been published for about last 40 years. Data obtained by us by processing of the space images of Landsat and ASTER is the latest material, which is the best tool for identification of the change in the number and area of the glaciers of Georgia during the last one century. The article presents the percentage and quantitative changes in the number and area of the glaciers of Georgia in the years of 1911–1960–1975–2014, according to the individual river basins. The air temperature course of the Georgia's high mountain weather stations has been studied. The river basins have been revealed, where there are the highest indices of the reduction in area and number of the glaciers and the reasons have been explained.

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

The current global warming has already lasted for longer than 150 years. In the middle of the 19th century, the Little Ice Age had come to its end and everywhere mountain glaciers began to decrease (Solomina, 2000). Since the end of the 1950s until the middle of the 1970s, the glaciers were in a quasi-stationary state in most of the mountain areas in Eurasia (Dyurgerov, 2005). Now, the glaciers degrade in all mountain areas of Eurasia. This is reflected in the fact that small glaciers disappear, the termini retreat, the area and volume of glaciers decrease, their surfaces are covered with moraines and large spaces of dead ice are being formed. Compound glaciers are broken into simpler components. Estimations of these changes were published in a number of papers; however, the general picture of the modern Eurasian glaciers to the present moment is not complete. Permanent, regular and detailed observations of the glacier behaviors are necessary to be performed in different regions (Barry, 2006; Khromova et al., 2014). Among them Caucasus, where the glaciers are an important source of water for agricultural production in Georgia and Azerbaijan, and runoff in large glacially-fed rivers (Kodori, Enguri, Rioni, Tskhenistskali, Nenskra) supplies several hydroelectric power stations. Caucasian glaciers also play a role in water levels in the Caspian Sea, the largest endoreic body of water on Earth. Since the mid-1970s, lake levels have risen ~ 2 m, with major socio-economic impacts (Arpe et al., 2000) on bordering coastal nations. Future trends in glacier change are thus a topic of considerable interest to the region.

2 Study area

One of the main centres of mountain glaciation in Europe is the Greater Caucasus Mountains located between the Black and Caspian Seas in the densely populated southwest of Russia and Georgia (Shahgedanova et al., 2014). The current number of glaciation is ~ 2000, with a total area of ~ 1100 km² and volume ~ 68 km³ (Radicì et al.,

TCD

9, 3777–3806, 2015

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2014). According to the morphological and morphometric characteristics the Greater Caucasus can be divided into three parts within Georgia – Western Caucasus, Central Caucasus and Eastern Caucasus (Maruashvili, 1971; Gobejishvili, 1995; Tielidze, 2014) (Fig. 1).

5 *Western Caucasus* region includes the part, which is located to the west of the Dalari Pass. It has a sublatitudinal direction in Georgia. The relief of its southern slope is characterized by complex orographic structure. The main watershed range is the highest morphological unit here. The Greater Caucasus branch-ranges: Gagra, Bzipi, Chkhalti (Abkhazeti) and Kodori, located in echelon, are also sharply distinguished morphologically and morphometrically (Geomorphology of Georgia, 1973).

10 *Central Caucasus* sector is the highest hypsometrically; it is characterized by a complex geological structure and is very interesting by glacial-geomorphological point of view because in the Pleistocene (Gobejishvili et al., 2011) and even today the main center of glaciation is located in the Central Caucasus. Its western boundary coincides with the Dalari pass and runs along the Enguri and Kodori Rivers' watershed (Kharikhra range), while its east boundary coincides with the Jvari Pass and then runs along the bottom of the river gorges of Tergi-Bidara-Mtiuleti's Aragvi (Maruashvili, 1971). In terms of the glaciers distribution, the several orographic units can be distinguished in the Central Caucasus: svaneti, Samegrelo, Letchkhumi, Shoda-Kedela and Java ranges.

20 To *Eastern Caucasus* belongs the part of the Greater Caucasus range, which is located to the east of the Georgian Military Road (Jvari Pass). Both the southern and northern slopes of the Caucasus range get within the Georgia's boundaries. Eastern Caucasus is quite high hypsometrically: heights of its peaks – Kuro, Komito, Shani, Amgha, Tebulosmta and others exceed 4000 m. Though, because of the relatively dry climate and morphological features of the relief, the contemporary glaciers are more weakly represented in the Eastern Caucasus than in the hypsometrically lower Western Caucasus.

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



according to the number of the contemporary glaciers, and according to the area – it is only behind the Enguri River basin. According to the data of 1911 there were 85 glaciers in the Rioni River basin with the area of 78.1 km². According to the data of 1960 the number of the glaciers was 112 with the total area of 75.1 km². By the data of 1975 there were 124 glaciers in this basin with the total area of 62.9 km². And according to the data of 2014 there are 97 glaciers with the total area of 46.7 km² (Table 1). The largest glacier in the Rioni River basin is Kirtisho with the area of 4.4 km².

By relatively low hypsometrical location is distinguished the Liakhvi River basin, which is located to the east of the Rioni River basin. According to the data of 1911 there were 12 glaciers in the basin with the area of 5.1 km². There were 16 glaciers in the Liakhvi River basin with the total area of 4.0 km² according to the data of 1960. By the data of 1975 there were 22 glaciers in this basin with the total area of 6.6 km². And according to the data of 2014 there are 10 glaciers in the Liakhvi River basin with the total area of 1.9 km² (Table 1).

The easternmost basin of the Central Caucasus, where the contemporary glaciers are presented, is the Aragvi River basin. According to the data of 1911 there were 3 glaciers with the total area of 2.2 km². According to the data of 1960 the area of all glaciers was 0.9 km². By the data of 1975 there were 6 glaciers in this basin with the total area of 1.6 km². And according to the data of 2014 the only one glacier (Abudelauri) is remained in the basin with the area of 0.3 km² (Table 1).

In total, the glaciers area decreased by 145.7 km² (34.8%) in the Central Caucasus during the last one century, while their number increased by 112 (40.9%) in the same period (Figs. 2 and 3).

4.1.3 Eastern Caucasus

In Georgia the Eastern Caucasus is represented both by southern and northern slopes. The basins of the rivers such as Tergi, Asa, Arghuni and Pirikita Alazani are located there. All of the river basins are distributed on the northern slopes of the Caucasus.

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The Tergi River basin is a main glaciation center of the Eastern Caucasus. Some of the peaks' heights exceed 5000 m here (Mkinvartsveri/Kazbegi 5033 m). According to the number of glaciers the Tergi River basin is in the fourth place after Enguri, Kodori and Rioni and its share is 9.1 % in the total number of the glaciers of Georgia. It is also in the fourth place by the area after Enguri, Rioni and Kodori, and its share in the total area of the glaciers of Georgia is 10.0 %. By the data of 1911 there were 63 glaciers in the Tergi River basin with the total area of 89.1 km². By the data of 1960 there were 99 glaciers with the total area of 67.2 km². According to the data of 1975 there were 129 glaciers with the area of 72.1 km². And by the data of 2014 there are 58 glaciers with the total area of 35.6 km² (Table 1).

The Asa River basin is located on the northern slope of the Greater Caucasus. Its name is Arkhotistskali in the territory of Georgia. Heights of some of the peaks in this region exceed 3700 m. By the data of 1911 there were 17 glaciers in the Asa River basin with the total area of 4.1 km². By the data of 1960 there were 9 glaciers in this basin with the total area of 2.6 km². According to the data of 1975 there were 3 glaciers with the area of 1.1 km². And by the data of 2014 there are still 3 glaciers with a total area of 0.6 km² (Table 1).

The Arghuni River basin is located on the northern slope of the Greater Caucasus and it has the meridional direction. Although the hypsometric benchmarks of the relief are quite high, the contemporary glaciation is presented in small scales, and the glaciers are characterized by the small sizes. By the data of 1911 there were 10 glaciers in the Arghuni River basin with a total area of 5.4 km². By the data of 1960 there were 17 glaciers with the total area of 2.7 km². According to the data of 1975 there were 14 glaciers with the area of 1.7 km². And by the data of 2014 there are only 6 glaciers with the total area of 0.5 km² (Table 1).

Pirikita Alazani River basin is located on the northern slope of the Greater Caucasus and is of latitudinal direction. Here the individual peaks' height is over 3800–4000 m. According to the data of 1911 there were 23 glaciers there with the total area of 19.1 km². By the data of 1960 the glaciers were reduced in size and though the num-

the relief as well. The relief of some of the river basins is built by Jurassic sedimentary rocks, which suffer heavy denudation. That is why the Pleistocene glaciation forms, where the snow is well-kept and collected, and therefore, is one of the important conditions for the existence of glaciers, are poorly preserved there (Gobejishvili et al., 2011; Tielidze, 2014).

As it was mentioned above, the main glaciation center on the *Central Caucasus* is the Enguri and Rioni River basins. According to the materials available to us, the area of the glaciers in the Rioni River basin was reduced by only 3.8% in the years of 1911–1960, while the area of the glaciers in the Enguri River basin was reduced only by 3.7%. In our opinion, the mentioned data is not true, because, as it was mentioned above, certain glaciers in the Rioni and Enguri basins are difficult to access for the plane table surveying; therefore, the first topographical survey of the Caucasus was conducted, the firm contours of the mentioned glaciers were incorrectly depicted, and some small glaciers were completely omitted. The catalog of 1911 by K. Podozerskiy, which is compiled based on the mentioned maps, is distinguished by the certain defects. As in the same period of 1911–1960 in the Rioni and Enguri basins the number of the glaciers considerably increased, namely: in the Rioni basin more than 27 glaciers, in the Enguri basin more than 125 glaciers, it is natural that the number of the glaciers would not have been increased so sharply due to such a low rate of the reduction in the area of the glaciers. As for the period of 1960–2014, the areas of the glaciers in the Rioni and Enguri basins were decreased quite greatly, respectively by 37.8 and 32.8%.

As for the *Western Caucasus* it should be noted that the Bzipi and Kelasuri River basin are the only two in Georgia, where the number of the glaciers has not been changed since 1960 (Table 1), one of the conditioning factors of which is a fact that in winter period falls more solid precipitation in the Western Caucasus (Abkhazeti sector) than in the Central and Eastern Caucasus (Kordzakhia, 1967; Gobejishvili, 1995), which is one of the necessary conditions for feeding and maintaining the glaciers.

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



6 Conclusions

As a result of our research we concluded that the area of the glaciers of Georgia has been reduced from 613.3 to 555.9 km² in the years of 1911–1960, while their number has been increased from 515 to 786 (Fig. 9). In the mentioned years the number of the glaciers has been increased in almost all of the river basins (with the exception of the Asa River basin), which was caused by the division of the large size of glaciers during their degradation.

In the years of 1960–1975 the decrease in the number of the glaciers from 785 to 755 and reduction in the area from 555.9 to 514.4 km² are observed as well (Fig. 9). Only in the Eastern Caucasus a slight increase is observed, which is caused by the fact that the temporary snow spots and snow areas are considered as glaciers in the Tergi River basin, which is not true. In 1975–2014 the area of the glaciers has been reduced from 514.1 to 355.8 km² and their number was reduced from 755 to 637 (Fig. 9). In 1975–2014 the simultaneous reduction in the number and area of the glaciers is caused due to the fact that for the years of 1960–1970 in Georgia dominated the small size of glaciers of cirque type, which have completely disappeared during the last half century. In total, the area of the glaciers of Georgia reduced by 42.0% in the years of 1911–2014, while their number increased by 23.7%.

As a result of the research it was identified that in the end of the 19th century and early 20th century, the largest glacier of Georgia was Tviberi (Fig. 10a). According to the topographical map of 1887 the glacier area was 49.0 km² and its tongue was ended at a height of 2030 m above sea level. Before 1960, the Kvitoldi glacier was separated from the Tviberi glacier's left side, which became an independent glacier (Fig. 10b2). In the topographical map of 1960 the area of the Tviberi was 24.7 km² and the glacier tongue was ended at the height of 2140 m a.s.l. (Fig. 10b1). In the Landsat aerial image of 2014 can be well seen the Tviberi degradation after 1960, when the relatively small size simple valley type of glaciers and even smaller cirque type of

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Kaab, A.: Monitoring high-mountain terrain deformation from repeated air- and spaceborne optical data: examples using digital aerial imagery and ASTER data, ISPRS J. Photogramm., 57, 39–52, 2002.

Khromova, T., Nosenko, G., Kutuzov, S., Muravievand, A., and Chernova, L.: Glacier area changes in Northern Eurasia, Environ. Res. Lett., 9, 015003, doi:10.1088/1748-9326/9/1/015003, 2014.

Kordzakhia, R.: Enguri and Tskhenistskhali river basins climate features within the Svaneti, Acts of Georgian Geographical Society, Vol. IX–X, Tbilisi, 110–125, 1967 (in Georgian).

Maruashvili, L.: Physical Geography of Georgia, Monograph. Publ. “Metsniereba”, Tbilisi, 1971 (in Georgian).

Podozerskiy, K. I.: Ledniki Kavkazskogo Khrebta (Glaciers of the Caucasus Range): Zapiski Kavkazskogo otdela Russkogo Geograficheskogo Obshchestva, Publ. Zap. KORGO., Tiflis, 29, 1, 200 pp., 1911 (in Russian).

Radicì V., Bliss, A., Beedlow, A. C., Hock, R., Miles, E., and Cogley, J. G.: Regional and global projections of twenty-first century glacier mass changes in response to climate scenarios from global climate models, Clim. Dynam., 42, 37–58, doi:10.1007/s00382-013-1719-7, 2014.

Shahgedanova, M., Nosenko, G., Kutuzov, S., Rototaeva, O., and Khromova, T.: Deglaciation of the Caucasus Mountains, Russia/Georgia, in the 21st century observed with ASTER satellite imagery and aerial photography, The Cryosphere, 8, 2367–2379, doi:10.5194/tc-8-2367-2014, 2014.

Solomina, O. N.: Retreat of mountain glaciers of northern Eurasia since the Little Ice Age maximum, Ann. Glaciol., 31, 26–30, 2000.

Tielidze, L. G.: Glaciers of Georgia, Monograph. Publ. “Color”, 254 pp., Tbilisi, 2014 (in Georgian).

Tielidze, L. G., Lomidze, N., and Asanidze, L.: Glaciers retreat and climate change effect during the last one century in the Mestiachala River Basin, Caucasus Mountains, Georgia, Earth Sci., 4, 72–79, doi:10.11648/j.earth.20150402.12, 2015a.

Tielidze, L. G., Kumladze, R., and Asanidze, L.: Glaciers reduction and climate change impact over the last one century in the Mulkhura River Basin, Caucasus Mountains, Georgia, Int. J. Geosci., 6, 465–472, doi:10.4236/ijg.2015.65036, 2015b.

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Mean annual temperatures of Georgia’s medium and high mountain meteorological stations.

Years	Mestia Mean annual °C	Years	Mamisoni Mean annual °C	Years	Jvari Pass Mean annual °C	Years	Kazbegi Mean annual °C
1906–1960	+5.9	1907–1960	–2.2	1907–1960	–0.1	1907–1960	–5.8
1961–2013	+6.0	1961–1995	–2.2	1961–2009	+0.2	1961–2009	–5.6
	Temperature increase +0.1		Temperature increase 0.0		Temperature increase +0.3		Temperature increase +0.2

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze




Figure 1. Georgian Caucasus glacier outlines (in yellow) derived from Landsat and ASTER imagery, and Georgia's mountain meteorological stations location.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze




Figure 3. The change in the area (km²) of the Western, Central and Eastern Caucasus glaciers in 1911–1960–1975–2014.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze




Figure 4. The course of the mean annual air temperatures in the Mestia, Mamisoni, Jvari Pass and Kazbegi meteorological stations over the last one century.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



TCD

9, 3777–3806, 2015

Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze




Figure 5. Mean monthly air temperature course of the Kazbegi meteorological station in the years of 1907–1960 and 1961–2009.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze




Figure 6. Mean monthly air temperature course of the Jvari Pass meteorological station in the years of 1907–1960 and 1961–2009.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Glaciers change over the last century, Caucasus Mountains, Georgia

L. G. Tielidze




Figure 10. (a) Tviberi glacier, topographical map of 1887; (b) topographical map of 1960; (b1) Tviberi glacier; (b2) Kvitolodi glacier; (c) Landsat L8 imagery; (d) photo of 1884 (M. V. Dechy); (e) photo of 2011 (L. G. Tielidze).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

