

Time forecast of a break-off event from a hanging glacier

Jérôme Faillettaz¹, Martin Funk², and Marco Vagliasindi³

¹3G, UZH, University of Zürich, Zürich, Switzerland

²VAW, ETHZ, Zürich, Switzerland

³Fondazione Montagna Sicura, Courmayeur, Aosta Valley, Italy

Correspondence to: jerome.faillettaz@geo.uzh.ch

Abstract.

A cold hanging glacier located on the south face of the Grandes Jorasses (Mont Blanc, Italy) broke off on the 23rd and 29th September 2014 with a total estimated ice volume of 105.000 m³. Thanks to accurate surface displacement measurements taken up to the final break-off, this event was
5 successfully predicted 10 days in advance, enabling local authorities to take the necessary safety measures. The break-off event also confirmed that surface displacements experience a power law acceleration along with superimposed log-periodic oscillations prior to the final rupture. This paper describes the methods used to achieve a satisfactory time forecast in real time and demonstrates, using a retrospective analysis, their potential for the development of early-warning systems in real
10 time.

1 Introduction

Rockfalls, rock instabilities due to permafrost degradation, landslides, snow avalanches or avalanching glacier instabilities are gravity-driven rupture phenomena occurring in natural heterogeneous media. Such events have a potential to cause major disasters, especially when they are at the origin of a chain of processes involving other materials such as snow (snow avalanche), water (flood)
15 and/or debris (mudflow). The reliable forecasting of such catastrophic phenomena combined with a timely evacuation of the endangered areas is often the most effective way to cope with such natural hazards. Unfortunately, accurate time prediction of such events remains a somewhat daunting task as (i) natural materials are heterogeneous, (ii) the heterogeneity is difficult to quantify and measure,
20 and (iii) the rupture is a non-linear process involving such heterogeneities. Although often located in a remote high-mountain environment, avalanching glacier instabilities offer an interesting starting point for investigating early-warning perspectives of break-off events, as a glacier consists of a single material (ice) lying on well-defined bedrock. This relative simplicity of the system allows the focus to be placed on the rupture processes leading to the initiation of the instability. Recently,
25 considerable efforts in monitoring, analyzing and modeling such phenomena have led to significant

advances in understanding the destabilization process and in improving early-warning perspectives (Faillettaz et al., 2015).

In general, it is possible to distinguish three types of avalanching glacier instabilities according to the thermal properties of their ice/bedrock interface (Faillettaz et al., 2011b, 2012, 2015). If temperate, the presence of liquid water in the glacier plays a key role in the initiation and the development of the instability as its presence influences the basal properties of the ice/bedrock interface (diminution of friction, lubrication or loss of support). In such cases, several preliminary conditions to be fulfilled can be identified, but an accurate time forecast of an impending break-off event is still far from being possible. If the ice/bed interface experiences a transition from cold to temperate, the presence of melt water may reduce the basal resistance, which promotes the instability. No clear and easily detectable precursory signs are known in this case, and the only way to infer any potential instability is to monitor the temporal evolution of the thermal regime. If the ice/bedrock is cold, glaciers are entirely frozen to their bedrock. This situation appears in the case of high altitude hanging glaciers located entirely in accumulation zone. The snow accumulation is mostly compensated by periodic break-off of ice chunks (Pralong and Funk, 2006), occurring once a critical point in glacier geometry is reached. The instability results from the progressive accumulation of internal damage due to an increasing stress regime caused by glacier thickening. In this case, the rupture occurs within the ice, typically a few meters above the bedrock. The maturation of the rupture was shown to be associated with a typical time evolution of both surface velocities (Faillettaz et al., 2008) and passive seismic activity (Faillettaz et al., 2011a). This characteristic time evolution can theoretically be used to predict the occurrence of a catastrophic event. This was done a posteriori with data obtained prior to the 2005 break-off of the Weisshorn glacier.

In this context, the Whymper glacier, a cold hanging glacier located at the Grandes Jorasses (Mont Blanc, Alps, Italy), already broke off several times in the past, leading to major ice avalanches that occasionally reached the bottom of the valley. In autumn 2008, the glacier recovered its previous critical geometry from the year 1998 and a crevasse opened in the lower part of the tongue, prompting the local authorities to initiate a monitoring program to enable a time forecast of the expected break-off event. The glacier finally broke off causing no damage in autumn 2014, after more than 5 years of monitoring. The break-off was successfully predicted 2 weeks in advance.

The aim of this paper is to confirm the validity of the time forecast procedure first developed in 2005 on the Weisshorn glacier and to apply it here in real time.

After describing the glacier and the monitoring system installed on the glacier, we analyze the time evolution of the surface displacement measurements in the context of a time forecast procedure. While comparing this break-off event with the Weisshorn event of 2005 we discuss the results obtained, with the goal of improving the understanding of this phenomenon.




Figure 1. Global view of the Val Ferret with Grandes Jorasses. White rectangle highlights Whymper glacier. Light red lines indicate possible avalanche flow path (for more details see Margreth et al. (2011)), red lines indicate avalanche path from the 1998 break-off event. Inset shows geographical situation of the studied glacier.

2 Grandes Jorasses glacier

2.1 Study site

The Whymper glacier is located on the south face of the Grandes Jorasses (Mont Blanc massif, Italy) between 3900 and 4200 m asl (Fig. 1). The front of the glacier is about 90 m wide and its surface area amounts 25,000 m². This very steep cold hanging glacier (about 40 °) lies above the village of Planpincieux and the Italian Val Ferret, a famous and highly frequented tourist destination both in winter and summer. In 1997, six boreholes were drilled down to the bed and temperature profiles were measured, indicating basal temperatures below the freezing point (below $-1.6 \pm 0.4^{\circ}C$) at all locations (Pralong and Funk, 2006). Historical data and morphological evidence indicate that the glacier experienced recurrent break-off events that can be dangerous, particularly in winter, when the initial ice avalanche can drag snow in its path. This hanging glacier periodically broke off in the past leading to large avalanches that flowed down into the valley.

2.2 Break-off event history

The glacier broke off several times during last 100 years. Some of these events have been observed and reported:




Figure 2. (a) Grandes Jorasses (Whymper) glacier before (23rd August 2014), (b) after the first break-off (23rd September 2014) and (c) after the second break-off (30th September 2014)

- On 21 December 1952, after an intensive snowfall period, a huge avalanche was released below the Grandes Jorasses which destroyed a 200-year old forest and blocked the bottom of the Val Ferret over a distance of more than 1 km. The avalanche volume was estimated at more than 1,000,000 m³. It is not clear whether the snow avalanche was triggered by an ice avalanche from the Whymper glacier.
- In August 1993 and July 1996, the glacier released ice avalanches of 80,000 and 24,000 m³, respectively. These ice avalanches did not reach the bottom of the valley.
- The last major break-off event occurred in the night of 31th May to 1st June 1998. Almost the entire Whymper glacier (around 150,000 m³) broke off at one time and the triggered ice avalanche reached the bottom of the valley, fortunately without causing damage (Fig. 1). According to Pralong and Funk (2006) the formation of the upper crevasse was observed 2.5 years before failure .

2.3 Present monitoring: 2009-2014

The survey primarily consisted of surface displacement measurements with an automatic total station and GPS as well as close-range photogrammetry (Margreth et al., 2011). Two reflectors set on the rock on both sides of the glacier were used as reference, and several reflectors mounted on stakes were directly drilled into the ice, so that their exact positions could be monitored (Fig. 2). Because of instrument problems, the seismic activity unfortunately could not be monitored as initially planned.

Starting in 2010, surface displacements were continuously recorded at several stakes at 2-hour intervals (when the prisms were visible, i.e., good weather conditions) with the aim to timely detect an impending ice avalanche (Margreth et al., 2011). Using the same correction technique as described by Faillettaz et al. (2008), the surface displacements could be determined with an accuracy better than 1 cm, allowing surface velocities to be inferred.

In parallel to the monitoring program, a safety concept for the valley floor was developed considering several scenarios of falling ice volumes. The different ice avalanche scenarios were simulated

using the 2-dimensional calculation model RAMMS. The necessary safety measures were defined according to the local avalanche danger level and the potential volume of a break-off event (Margreth et al., 2011).

2.4 The 2014 break-off event

105 From 2010 on, surface displacements were surveyed without interruption. The Whymper glacier finally broke off with an estimated ice volume similar to the 1998 event (about 105,000 m³). Contrary to the 1998 event, the glacier broke off in two events on September 23rd and on September 29th 2014, without reaching the valley (Fig. 2). At the final break-off, 4 reflectors were still active, 2 of them in place for more than 2 years. Despite poor weather conditions between the 16th and 21st 110 of September, the monitoring was operational up to the final break-off. By chance, there was one reflector on each of the two unstable parts and one on the stable part, (Fig. 2).

Striking qualitative analogies with those of the 2005 Weisshorn event (Faillettaz et al., 2008) can be highlighted.

1. This steep cold hanging glacier experiences periodic break-off events.
- 115 2. The geometrical configuration of the glacier is similar before each break-off, with an upper crevasse spanning the whole glacier and a clear thickening of the glacier towards its tongue.
3. The upper crevasse marks a clear distinction between a stable upper part (where Stake 4 is located) and a downstream unstable part (where the other reflectors were located, Fig. 2, section 4.1). A crude estimation of the volume of the unstable part is thus possible.
- 120 4. Downstream of this crevasse, surface displacements experience a typical acceleration prior break-off, whereas upstream this crevasse constant velocities are recorded (Stake 4 in Fig. 3).
5. The rupture did not occur at the ice/bedrock contact but a few meters above it, within the ice (Fig. 2), possibly because of bedrock irregularities (Pralong and Funk, 2006).
- 125 6. The whole break-off occurred in two steps; a minor section at the left side of the glacier was released first.

3 Previous findings on cold glacier break-off

Based on a retrospective analysis, the main conclusion drawn by Flotron (1977) and R othlisberger (1981) was that the forecast of a break-off event from a hanging glacier was possible using surface displacements alone. The principle is to fit the characteristic acceleration of the surface motion with 130 a power law behavior of the form:

$$s(t) = s_0 + u_s t - a(t_c - t)^\theta, \quad (1)$$

where $s(t)$ is the displacement (in meters) at time t (in days), s_0 a constant in meters, u_s the constant velocity of the upstream stable part (in md^{-1}), t_c the critical time (in days), $\theta < 0$ (without units) and a (in $\text{md}^{-\theta}$) the parameters characterizing the acceleration. In this way, the critical time t_c , i.e.,
135 time at which the theoretical displacement becomes infinite, could be evaluated using such empirical law. Although the break-off event would necessarily occur earlier, this critical time represents the upper limit of the break-off timing. Moreover, an oscillating pattern superimposed on the power law acceleration of the surface displacements was evidenced prior to the 2005 Weisshorn event (Pralong et al., 2005; Faillettaz et al., 2008). This peculiar glacier dynamics was shown to be a log-
140 periodic oscillating process superimposed on this acceleration (for appearance and interpretation see Faillettaz et al. (2015)). The time evolution of the surface displacement measurements can be described with the following equation (after Sornette and Sammis, 1995; Pralong et al., 2005):

$$s(t) = s_0 + u_s t - a(t_c - t)^\theta \left[1 + C \sin \left(2\pi \frac{\ln(t_c - t)}{\ln(\lambda)} + D \right) \right], \quad (2)$$

where C the relative amplitude (without units), λ the logarithmic frequency (in days) and D the
145 phase shift of the log-periodic oscillation (without units).

Thanks to a combined analysis of surface displacement and seismic measurement, Faillettaz et al. (2011a) were able to obtain a coherent quantitative picture of the damage evolution process developing before the 2005 Weisshorn break-off. They have suggested three regimes in the evolution of the failure process leading to the break-off event:

- 150 (i) A first stable phase related to a self-organizing regime, where diffuse damage accumulates within the glacier, with a proliferation of dislocation-like defects.
- (ii) A transitional phase where the damage process goes on, micro-cracks grow and start merging in a homogeneous way. Log-periodic oscillations appear and reveal the hierarchical structure of the fracture process under development.
- 155 (iii) A catastrophic regime where damage clusters are randomly activated. Damage clusters interact and merge with a preferential direction (i.e. preparing the final rupture pattern), in contrast to the previous regime.

4 Results

4.1 Surface displacements and associated velocities

160 Fig. 3 shows the corrected surface displacements and Fig. 4 the associated derived surface velocities of the 4 stakes (Fig 2) prior to the break-off. The associated derived surface velocities are computed taking the surface displacements (smoothed over 5 points) interpolated on regular time step of one day. Note that Stakes 13 and 14 have more than 2 years of nearly continuous measurements and the




Figure 3. Surface displacements of the 4 stakes before the break-off using 19 July 2014 as reference (when Stake 2 and Stake 4 were installed). Vertical red dashed lines indicate the occurrence of the two break-offs, on 23rd and 29th September 2014. Interrupted lines indicate a period of bad weather conditions without measurements. Note that Stake 14 was not surveyed after 16th of September 2014, i.e., one week before the first break-off.

position of Stake 13 was surveyed up to the final break-off event on September 29th. After processing
 165 the raw data with two reference points located near the glacier and additionally accounting for flow
 direction, the accuracy of the results was less than a centimeter (Faillettaz et al., 2008). Note that
 this constitutes a unique dataset not only because of the great accuracy and long measurement period
 but also due to available surface displacement data up to a few hours prior to the break-off event.
 Whereas surface velocities at Stake 4 are approximately constant (Fig. 4), the three other stakes show
 170 a clear acceleration which is typical for an unstable situation. According to this observation we can
 expect that the glacier section around Stake 2, 13 and 14 will break-off, while the section around
 Stake 2 will remain stable (section 2.4).

4.2 Application to forecasting

Previous findings (section 3) were applied in order to forecast the breaking-off event in real time.
 175 As soon as a significant increase in velocity was detected, the same procedure was followed as in
 Faillettaz et al. (2008). We periodically fitted surface displacements of all stakes to a power law
 (Eq. 1) and a log periodic oscillating behavior (Eq. 2). The nonlinear least-squares curve-fitting
 was performed using the Levenberg-Marquardt algorithm. Because the results depend on the initial
 parameter estimates, especially t_c and θ , we have systematically used different initial values with a
 180 prescribed bound and selected the results corresponding to the best root mean squared error and the
 degree-of-freedom adjusted coefficient of determination.

Fig. 5a and 6a show both power law (Eq. 1) and log-periodic (Eq. 2) fits using the last month
 of available data, i.e up to 16th September for Stake 14 and 29th September for Stake 13. As both




Figure 4. Surface velocities for 4 stakes since 2012. Inset shows a closer view during the same period as in Fig 3.

fits are barely distinguishable, we have also plotted on Fig. 5b and 6b the residuals to the power law fit and show the associated log-periodic fit (minus the power law fit) as a dashed gray line ;
 185 Table 1 contains the values of the parameters in Eq. 2, taking $\lambda = 2$ d. Note that measurements are available up to the final break-off for 3 prisms (i.e., Stake 13, Stake 2 and Stake 4) and stopped on 16 September for Stake 14, i.e., 8 days before the first break-off.

It appears that the power law behavior describes well the surface displacements with a maximum
 190 discrepancy of about 5 cm for Stake 14 (8 days before break-off), about the same order of magnitude as the one observed during the 2005 Weisshorn event (Fig. 5). However, residuals indicate an oscillating pattern. When using the log-periodic function (Eq. 2), the agreement between measured and fitted values (dashed gray line) becomes significantly better, with an accuracy of the order of magnitude of the measurement accuracy (less than a centimeter). Results show that the critical time can
 195 be expected around the 3rd October for both stakes, which is fairly close to the observed break-off. Note that such an approach can be used to investigate how far in advance a reliable time forecast is possible (see section 5.4).

However, even if Stake 14 is located on a section that broke off earlier, no significant differences
 200 could be detected. Our approach is not able to detect whether the break-off will occur all at once or as successive small events.

Now when considering the entire dataset for Stake 13 (where measurements could be recorded up to the break-off) using the same method, it appears that the amplitudes of the oscillations superimposed on the power law acceleration become even larger close to the break-off - they reach values up to 30-40 cm (Fig. 6). Such a broad oscillating pattern has never been observed before, confirming
 205 that the jerky motion of the glacier (with oscillating nature) might have a physical origin (see Section 5.2).




Figure 5. a. Surface displacements of Stake 14 for the period 16th August- 16th September (last measurement of Stake 14) and the associated power law (in blue) and log-periodic fit (in gray). b. Residuals (in meters) to the power law fit (in blue) for the same period. Log-periodic fit is also shown with dashed gray line. Values for the parameters are shown in Table 1.

5 Discussion

5.1 Influence of data accuracy on the final result

To assess how the data accuracy influences the time forecast of the break-off, we artificially added
 210 two uniformly distributed random noise (between -1 and 1 cm and between -5 and 5 cm) to our
 dataset and to analyze how the obtained critical time depends on our fitting method. To ensure good
 statistical representation, this procedure was performed 100 times on Stake 14 up to 16.09.2014 and
 on Stake 13 up to the final break-off (29.09.2014). Results (see table 2) show that the forecasting
 errors are mainly due to the fitting procedure. Errors due to the data accuracy are about one order of
 215 magnitude lower than those generated by the fitting procedure.

5.2 Appearance of log-periodic behavior

The origin of the log-periodic oscillating behavior is likely due to a Discrete Scale Invariance (DSI),
 a weaker kind of scale invariance according to which the system obeys scale invariance only at a
 specific scaling factor scale (Sornette and Sammis, 1995; Sornette, 1998; Zhou and Sornette, 2002a;

Table 1. Values of the estimated coefficients of Equation 2 with $\lambda = 2$ d and the root-mean-square error (RMSE) of the fit, first two columns corresponding to the parameters of the fit used in Fig. 5 for the period 16.08-16.09, the last two columns corresponding to the parameters of the fit used in Fig. 6 for the period 30.08- 30.09. t_c is given in days after the first day of the investigated period.

Parameter	Units	Stake 13 (16.08 - 16.09)	Stake 14 (16.08 - 16.09)	Stake 13 (29.08 - 29.09)	Stake 2 (29.08 - 29.09)
t_c	d	48.02 ± 5.13	48.11 ± 5.6	41.93 ± 0.91	41.80 ± 2.18
	date	03-Oct-2014	03-Oct-2014	10-Oct-2014	10-Oct-2014
θ	-	-0.24	-0.25	-0.99	-0.98
s_0	m	$-1.47 \cdot 10^4$	-1.47×10^4	-2.03×10^4	-1.48×10^4
u_s	md^{-1}	2.00×10^{-2}	2.00×10^{-2}	2.99×10^{-2}	2.27×10^{-2}
a	$\text{md}^{-\theta}$	27.88	27.72	141.73	164.42
C	-	2.9×10^{-3}	2.3×10^{-3}	3.0×10^{-2}	2.0×10^{-2}
D	-	2.25	1.97	6.13	0.06
RMSE	md^{-1}/m	8.7×10^{-3}	7.9×10^{-3}	3.05×10^{-2}	2.51×10^{-2}




Figure 6. a Surface displacement of Stake 13 for the period 29th August 2014 - 29th September 2014 and associated power law (in blue) and logperiodic (in gray) fits. b. Residuals (in meters) to the power law fit (black point) for the same period. Log-periodic fit is shown with gray dashed line. Values of the parameters are shown in Table 1.

Stake	Initial data	$\pm 1\text{cm}$ noise	$\pm 5\text{cm}$ noise
13	$t_c = 41.93 \pm 0.91$ d	$t_c = 41.90 \pm 0.038$ d	$t_c = 41.86 \pm 0.041$ d
14	$t_c = 48.11 \pm 5.6$ d	$t_c = 48.55 \pm 0.56$ d	$t_c = 49.33 \pm 0.55$ d

Table 2. Critical time t_c evaluated with our dataset and with a uniformly distributed random noise of ± 1 cm and ± 5 cm.

220 Sornette, 2006). Whereas the hallmark of Continuous Scale Invariance is the existence of power law, the signature of DSI is the presence of power laws with complex exponents which manifests itself in data by log-periodic corrections to scaling. Several mechanisms may lead to this partial breaking of the continuous symmetry. Thanks to a combined analysis of surface displacements and seismic measurements, Faillettaz et al. (2011a) suggest that it results from the dynamic interactions between
 225 newly developed micro-cracks, as shown by Huang et al. (1997) and Sahimi and Arbabi (1996).

To identify the log-frequency, we analyzed the data in the same way as Faillettaz et al. (2008) with a Lomb periodogram analysis (Press, 1996; Zhou and Sornette, 2002b), which is designed to analyze non-uniformly sampled time series. This method enables us to determine f_{Lomb} as a function of $\cos(2\pi f_{Lomb}t)$. The parameter λ in Equation 2 can then be evaluated easily as $\lambda = e^{1/f_{Lomb}}$.
 230 Unfortunately, the critical time t_c has to be known to perform this analysis, i.e., this analysis can only be performed a posteriori. It clearly shows (Fig. 7a) peaks in Lomb power (power spectral density) at $\lambda \sim 2$ d for the three analyzed points, confirming that the oscillating behavior is not a measurement artefact but has physical origins, such as the merge of newly developed micro-cracks. Note that another strong log frequency appears at $\lambda \sim 7.4$ d for Stakes 2 and 13 (Fig. 7b), after the
 235 first break-off. The reason for the appearance of such peak is not clear, but is probably induced by the occurrence of the first break-off that changes the geometry of the glacier: Using experimental data, Moura et al. (2005, 2006) suggested that grain size and loading rate directly influence log-periodic oscillations. A possible explanation would thus be that the first release led to a sudden change in the global loading of the remaining section of the glacier, i.e., loading rate change, introducing
 240 another subharmonic log frequency and perturbing the overall behavior of the remaining section of the glacier where Stakes 2 and 13 stand.

5.3 Accurate determination of break-off occurrence

As critical time t_c given by power law or log-periodic fit indicates when surface displacements become theoretically infinite. However, the break-off event is expected before t_c . When fitting in
 245 real time the surface displacements with both power law and log-periodic behavior, it is not only possible to assess the critical time but also the time at which the derived velocities are expected to reach a given threshold (for example 50 cm d^{-1} or 1 m d^{-1}). Fitting and estimating the time at which the velocity reaches a given threshold provides a more accurate way to predict the break-off




Figure 7a. Lomb periodogram for Stakes 13 and 2 (in inset) as well as the corresponding log frequencies (λ) of the peaks.




Figure 7b. Lomb periodogram for Stake 14 as well as the corresponding log frequencies (λ) of the peaks.

event. We developed a software based on this idea by fitting in real time the measurements with both
 250 power law and log-periodic behavior and thus provide an estimate of the break-off time.

According to our knowledge, it is not possible to know in advance the velocity at which break-off will occur. However, from previous events (Weisshorn 1973 and 2005 event, Flotron (1977); R othlisberger (1981); Faillettaz et al. (2008)), it seems that break-off occurs between 50 cm d^{-1} to 1.2 m d^{-1} , but this is based on a restricted number of events.

255 Taking threshold surface velocities of 50 cm d^{-1} and 1 m d^{-1} , our analysis (using Eq. 2) performed every days from the 12 September to 16 September suggested that break-off could occur between the 23 September ($v_{th} = 50 \text{ cm d}^{-1}$) and the 29 September ($v_{th} = 1 \text{ m d}^{-1}$). Note that the two breaking-off events occurred exactly at these two days, which were forecasted 10 days in

advance. Following this analysis, alert was immediately sent to the authorities leading them to close
260 the endangered area one week before the event. Note that the definition of the velocity threshold has
an influence on the prediction itself, as we saw nearly one week is needed for the glacier to accelerate
from 50 cm d^{-1} to 1 m d^{-1} . The precise prediction would also not only be based on a correct fit of
the surface displacement data but also on a guess of this parameter. We suggest to choose 40 cm d^{-1}
as a conservative threshold to define a safe break-off danger time interval.

265 **5.4 How far in advance are time forecasts possible?**

Surface displacements were analyzed retrospectively based on the last month of data for each stake,
and the critical time as well as the time at which the fitted velocity reached 50 cm d^{-1} (v_{50}) and
 1 m d^{-1} (v_{100}) were plotted as a function of the time (Fig. 8). Associated errors (Fig. 8 bottom)
account for the fitting procedure.

270 First, the prediction is better when using log-periodic fit than power law fit. This retrospective
analysis shows that the prediction is correct after 12 September, i.e., 11 and 17 days before the
break-off with a confidence interval becoming less than 10 days with a log-periodic fit.

This analysis points out the great prediction potential - and early warning perspective - of this
method, as the exact time of the break-off could be forecasted almost 2 weeks in advance. Note that
275 both the power law and the log-periodic fits become less accurate after the first break-off for Stake
13. This might be related to the sudden change in glacier geometry that may influence the surface
displacements at Stake 13. However, note that the time at which a velocity of 1 m d^{-1} (v_{100}) is
expected remains unaffected, still pointing at September 29th.

5.5 Overall recommendations

280 According to the knowledge gained from the different studies on Weisshorn, Mönch and Grandes
Jorasses glaciers, accurate data are required to forecast an impending break-off event. As the am-
plitudes of the log-periodic oscillations are increasing towards the break-off (from 5 cm one week
before the break-off to 40 cm at the break-off), the confidence of the time forecast strongly depends
on the precision of the surveying data. To ensure a satisfactory forecast about one week in advance, a
285 surveying accuracy better than half of the expected log-periodic amplitudes, i.e., 2.5 cm, is required.
In this study an accuracy of 1 cm was achieved with an automatic total station (Leica theodolite
TM1800 combined with the DI3000S Distometer). The sampling rate needs to be adapted to the
oscillating pattern in order to enable its detection. Moreover, in such rapid changing meteorology
where clouds can momentarily hindered measurements, several measurements need to be performed
290 each days. A sampling rate of 2 hours was chosen in this study, ensuring thus several opportunities
to obtain data every day. This technique can be performed in near real time and several measure-
ments can be performed every day with a sufficient accuracy. Note that GPS measurements would
be a valuable alternative but this technique requires a long acquisition time and additional process-




Figure 8. Top: Thick lines: evaluated critical time t_c for power law (blue) and log-periodic (red) fit for Stakes 13 (left) and 14 (right) as a function of the time of analysis. Interrupted lines indicate time at which estimated derived velocity from power law and log-periodic fit reaches 50 cm d^{-1} (dashed lines, v_{50}) and 1 m d^{-1} (dot-dash line, v_{100}). Horizontal grey lines represent the observed break-off (23rd and 29th September). Bottom: Error in days on critical time fitted with power law (blue) and log-periodic (red) estimated from the 95% confidence interval. Errors on v_{50} and v_{100} are similar to the errors on critical time, as they are directly derived from these fits.

ing to achieve to required accuracy. Although independent of weather conditions, the power supply
 295 and data transmission are problems to be solved. This procedure based on power law/log-periodic
 oscillations regression requires at least two measurements points on the potentially unstable part of
 the glacier, so that the time evolution of surface motion at different points could be compared. It also
 ensures that the results are not affected by stake/prism stability issues.

An alternative surveying technique is terrestrial Insar. The advantage of this technique is that no
 300 installation on the glacier (potentially dangerous) is required. However the data accuracy which can
 be expected with this monitoring system is not completely clear yet (Preiswerk et al., 2016)

6 Conclusions

Grandes Jorasses glacier broke off twice, on 23rd and 29th September 2014. In 2008, as it was
 suspected that a large part of this glacier is becoming unstable, a monitoring program was initiated.
 305 At the time of the break-off, 4 stakes covering a large part of the glacier enabled surface displacement
 measurements up to the time of the break-off. By regularly analyzing the dataset, it was possible to
 forecast the event ten days in advance. In the following the local authorities closes the endangered
 area up to the final rupture.

It was possible to confirm for an impending ice fall that a time series of surface displacements exhibits strong log-periodic oscillations superimposed on a global power law acceleration, as first discovered for the Weisshorn event (Faillettaz et al., 2015). In the immediate vicinity of the break-off, such oscillations reached an amplitude of more than 40 cm, almost one order of magnitude larger than revealed in previous findings. By fitting our recorded surface displacements, the critical time, i.e. time at which surface displacement become infinite, can be determined. Using this critical time value as an upper bound, a good time forecast could be achieved.

The inferred surface velocities immediately prior the two events were 0.5 m d^{-1} and 1.2 m d^{-1} , in the same range as for the Weisshorn event, suggesting that break-off of a cold hanging glacier could occur as soon as surface velocities reached 0.5 m d^{-1} . We showed that evaluating the time at which extrapolated velocities (based on the log-periodic fit) reach a prescribed threshold (0.5 m d^{-1} and 1 m d^{-1}) provides a significantly better forecast. However, in the present case, surface velocity increased from 50 to 100 cm/d in the order of one week. In practice, we suggest to use a critical velocity of $v=0.4 \text{ m d}^{-1}$ to determine the period of highly likely break-off occurrence. A retrospective analysis based on this method showed that an accurate prediction of the phenomenon can be achieved two weeks before its occurrence using the last month of surface displacement data and 0.5 m d^{-1} and 1 m d^{-1} as velocity thresholds. Although enabling a crude estimation of the total unstable ice volume, this point based surveying procedure is not appropriate to determine whether the unstable ice mass will fall down in one event or disaggregate and give rise to several smaller events, as no differences in the evolution of surface displacements were detected. This has consequences for the risk evaluation, as the resulting ice avalanche (and also the chain of processes resulting from its release) depends on the falling ice volume. To conclude, our results suggest that the presented monitoring and data processing techniques exploiting the log-periodic oscillating behavior can be applied in real time to forecast a break-off event on any cold unstable hanging glacier.

Acknowledgements. The authors thank Susan Braun-Clarke for proofreading the English. The authors thanks also J. Bassis and two anonymous reviewers for their useful comments that significantly improved the manuscript.

335 References

- Faillietaz, J., Pralong, A., Funk, M., and Deichmann, N.: Evidence of log-periodic oscillations and increasing icequake activity during the breaking-off of large ice masses, *J. Glaciol.*, 54, 725–737, doi:10.3189/002214308786570845, <http://www.ingentaconnect.com/content/igsoc/jog/2008/00000054/00000187/art00016>, 2008.
- 340 Faillietaz, J., Funk, M., and Sornette, D.: Icequakes coupled with surface displacements for predicting glacier break-off, *J. Glaciol.*, 57, 453–460, doi:10.3189/002214311796905668, <http://www.ingentaconnect.com/content/igsoc/jog/2011/00000057/00000203/art00007>, 2011a.
- Faillietaz, J., Sornette, D., and Funk, M.: Numerical modeling of a gravity-driven instability of a cold hanging glacier: reanalysis of the 1895 break-off of Altelsgletscher, Switzerland, *J. Glaciol.*, 57, 817–831, doi:10.3189/002214311798043852, <http://www.ingentaconnect.com/content/igsoc/jog/2011/00000057/00000205/art00005>, 2011b.
- 345 Faillietaz, J., Funk, M., and Sornette, D.: Instabilities on Alpine temperate glaciers: new insights arising from the numerical modelling of Allalingsletscher (Valais, Switzerland), *Nat. Hazard Earth Sys.*, 12, 2977–2991, doi:10.5194/nhess-12-2977-2012, <http://www.nat-hazards-earth-syst-sci.net/12/2977/2012/>, 2012.
- 350 Faillietaz, J., Funk, M., and Vincent, C.: Avalanching glacier instabilities: review on processes and early warning perspectives, *Rev. Geophys.*, pp. 203–224, doi:10.1002/2014RG000466, <http://dx.doi.org/10.1002/2014RG000466>, 2014RG000466, 2015.
- Flotron, A.: Movement studies on hanging glaciers in relation with an ice avalanche, *J. Glaciol.*, 19, 671–672, 1977.
- 355 Huang, Y., Ouillon, G., Saleur, H., and Sornette, D.: Spontaneous generation of discrete scale invariance in growth models, *Phys. Rev. E*, 55, 6433–6447, doi:10.1103/PhysRevE.55.6433, <http://link.aps.org/doi/10.1103/PhysRevE.55.6433>, 1997.
- Margreth, S., Faillietaz, J., Funk, M., Vagliasindi, M., Diotri, F., and Broccolato, M.: Safety concept for hazards caused by ice avalanches from Whymper hanging glacier in the Mont-Blanc massif., *Cold Reg. Sci. Technol.*, 69, 194–201, doi:10.1016/j.coldregions.2011.03.006, 2011.
- 360 Moura, A., Lei, X., and Nishisawa, O.: Prediction scheme for the catastrophic failure of highly loaded brittle materials or rocks, *Journal of the Mechanics and Physics of Solids*, 53, 2435 – 2455, doi:10.1016/j.jmps.2005.06.004, <http://www.sciencedirect.com/science/article/pii/S0022509605001110>, 2005.
- 365 Moura, A., Lei, X., and Nishisawa, O.: Self-similarity in rock cracking and related complex critical exponents, *Journal of the Mechanics and Physics of Solids*, 54, 2544 – 2553, doi:10.1016/j.jmps.2006.06.010, <http://www.sciencedirect.com/science/article/pii/S002250960600127X>, 2006.
- Pralong, A. and Funk, M.: On the instability of avalanching glaciers, *J. Glaciol.*, 52, 31–48, doi:10.3189/172756506781828980, <http://www.ingentaconnect.com/content/igsoc/jog/2006/00000052/00000176/art00004>, 2006.
- 370 Pralong, A., Birrer, C., Stahel, W. A., and Funk, M.: On the predictability of ice avalanches, *Nonlinear Proc. Geoph.*, 12, 849–861, doi:10.5194/npg-12-849-2005, <http://www.nonlin-processes-geophys.net/12/849/2005/>, 2005.

- Preiswerk, L., Wlaser, F., Anandakrishnan, S., Barfucci, G., Beutel, J., Burkett, P., Dalban, P., Funk, M.,
375 Limpach, P., Marchetti, E., Meier, L., and Neyer, F.: Monitoring unstable parts in the ice-covered Weiss-
mies northwest face, in: Proceedings of INTRAPRAEVENT 2016, p. in Press, 2016.
- Press, W.: Numerical Recipes in Fortran 90: Volume 2, Volume 2 of Fortran Numerical Recipes: The Art of
Parallel Scientific Computing, Fortran numerical recipes, Cambridge University Press, <http://books.google.ch/books?id=SPEi4mCfhacC>, 1996.
- 380 Röthlisberger, H.: Eislawinen und Ausbrüche von Gletscherseen, in: P. Kasser (Ed.), Gletscher und Klima
- glaciers et climat, Jahrbuch der Schweizerischen Naturforschenden Gesellschaft, wissenschaftlicher Teil
1978, pp. 170–212, Birkhäuser Verlag Basel, Boston, Stuttgart, 1981.
- Sahimi, M. and Arbabi, S.: Scaling Laws for Fracture of Heterogeneous Materials and Rock, *Phys. Rev.
Lett.*, 77, 3689–3692, doi:10.1103/PhysRevLett.77.3689, [http://link.aps.org/doi/10.1103/PhysRevLett.77.](http://link.aps.org/doi/10.1103/PhysRevLett.77.3689)
385 3689, 1996.
- Sornette, D.: Discrete-scale invariance and complex dimensions, *Phys. Rep.*, 297, 239 –
270, doi:10.1016/S0370-1573(97)00076-8, <http://www.sciencedirect.com/science/article/pii/S0370157397000768>, 1998.
- Sornette, D.: Critical Phenomena in Natural Sciences: Chaos, Fractals, Selforganization and Disorder:
390 Concepts and Tools, Springer Series in Synergetics, Springer, <http://books.google.com.au/books?id=uYHQHV2hCpwC>, 2006.
- Sornette, D. and Sammis, C. G.: Complex Critical Exponents from Renormalization Group Theory of Earth-
quakes: Implications for Earthquake Predictions, *J. Phys. I France*, 5, 607–619, doi:10.1051/jp1:1995154,
<http://dx.doi.org/10.1051/jp1:1995154>, 1995.
- 395 Zhou, W.-X. and Sornette, D.: Generalized q analysis of log-periodicity: Applications to critical ruptures,
Phys. Rev. E, 66, 046111, doi:10.1103/PhysRevE.66.046111, [http://link.aps.org/doi/10.1103/PhysRevE.66.](http://link.aps.org/doi/10.1103/PhysRevE.66.046111)
046111, 2002a.
- Zhou, W.-X. and Sornette, D.: Evidence of intermittent cascades from discrete hierarchical dissipation in
turbulence, *Physica D: Nonlinear Phenomena*, 165, 94 – 125, doi:10.1016/S0167-2789(02)00390-1, <http://www.sciencedirect.com/science/article/pii/S0167278902003901>, 2002b.
400