

Brief Communication: Glaciers in the Hunza Catchment (Karakoram) are in balance since the 1970s

Tobias Bolch^{1,2}, Tino Pieczonka², Kriti Mukherjee², Joseph Shea³

¹Department of Geography, University of Zurich, 8057 Zurich, Switzerland

5 ²Institute for Cartography, Technische Universität Dresden, 01069 Dresden, Germany

³International Centre for Mountain Development (ICIMOD), Kathmandu, Nepal

Correspondence to: Tobias Bolch (tobias.bolch@geo.uzh.ch)

Abstract. Previous geodetic estimates of mass changes in the Karakoram revealed balanced budgets or a possible slight
10 mass gain since the year ~2000. Indications for longer-term stability exist **but no mass budget analyses are available before**
2000. Here, we show that glaciers in the Hunza River basin (Central Karakoram) were on average in balance or showed
slight insignificant mass loss within the period ~1973 – 2009. Heterogeneous behaviour and frequent surge activities were
also characteristic for the period before 2000. Surge-type and non-surge-type glaciers showed on average no significantly
different mass change values. However, some individual glacier mass change rates differed significantly for the periods
15 before and after ~2000. **These analyses are based on low cost stereo Hexagon KH9 images from the 1970s, freely available**
and automatically generated digital terrain models (DTMs) from ~2009 ASTER data, and the SRTM DTM.

1 Introduction

Glacier melt water is of high importance for the run-off of the Indus River (Immerzeel et al., 2010) but the exact glacier
share is not known. This is partly due to the lack of knowledge about precipitation, snow cover, and snow water equivalent,
20 but also about glacier mass balance, their characteristics and their responses to climate change. Karakoram glaciers, which
occupy a large portion of the glacierized area of the Indus basin, have recently shown unusual behaviour: on average no
significant area changes but frequent advances and surge activities have been observed during the last decades (Bhambri et
al., 2013; Bolch et al., 2012; Hewitt, 2011; **Copland et al., 2011**). Geodetic mass estimates revealed balanced glacier mass
budgets or even slight mass gain since ~2000 (Rankl and Braun, 2016; Gardelle et al., 2013; Käab et al., 2015). However,
25 **almost no mass budget analyses are available** for Karakoram glaciers prior to the year 2000. The only exception is Siachen
Glacier in eastern Karakoram, for which Zaman and Liu (2015) corrected the erroneous value of $-0.51 \text{ m w.e. a}^{-1}$ given by
Bhutiyani (1999), and estimated the mass budget to be between $+0.22 \text{ m}$ and $-0.23 \text{ m w.e. a}^{-1}$. Herreid et al. (2015) found no
significant change in debris-coverage of the glaciers in the Hispar and Shimshal sub-regions of the Hunza River basin for the
period 1977 until 2014 and concluded that this might be due to balanced glacier budgets during this period. Temperature
30 measurements ~~that are~~ available since 1961 in the Karakoram show, in contrast to many other regions of the globe, a

consistent decline in summer and an increase during winter (Fowler and Archer, 2006). Hence, these measurements would support the assumption that glaciers would be in balanced or slightly positive conditions over the last several decades of the 20th century.

Declassified stereo satellite images from the 1960s and 1970s such as Corona KH-4 and Hexagon KH-9 have been proven to be suitable to generate digital terrain models (DTMs) and assess glacier mass changes since the 1960s (Bolch et al., 2008; Pieczonka et al., 2013). Hence, the aim of this study is to revisit existing information and extend the time series back to some of the earliest available satellite imagery. We focus on the Hunza catchment in the Central Karakoram (Figure 1) where high heterogeneity of glacier behaviour was found in previous studies (e.g. Bolch et al., 2012; Quincey and Luckman, 2014). Moreover, suitable Hexagon KH-9 data from the 1970s and recent stereo data from ~2010 such as ASTER and Cartosat-1 data were available. The Hunza River is a tributary to the Gilgit River, which flows into the upper Indus. The area of the basin is about 13,715 km² and approximately 25% of the basin is covered by glaciers. These glaciers constitute more than 15% of the glacierized area of the entire Karakoram. Frequent surges reported for several glaciers in this basin (Quincey and Luckman, 2014; Copland et al., 2011; Rankl et al., 2014) complicate the analysis of mass budgets as often only a certain part of a surge is captured.

15




Figure 1: Overview map of the study region

2. Data and Methods

2.1 Data

The SRTM digital terrain model (DTM) version 4, with a spatial resolution of 1 arc second (~30m, SRTM1) was utilized as reference dataset. The SRTM1 DTM was acquired by the use of two C-Band radar antennas (operating in interferometric mode) during 11 - 22 February 2000 and is frequently used for glaciological investigations. It can be assumed that the represented ice surface of the ablation region is close to the surface at the end of the 1999 ablation period, assuming a full penetration of the radar beam into snow (Paul and Haeberli, 2008). However, deeper radar penetration can be expected in the accumulation region (Berthier et al., 2006). Data voids which are mainly restricted to some accumulation areas were filled with ASTER GDEM2 data (see section 3.2).

This DTM is a merge of several ASTER scenes covering the period ~2000 – 2010 (Slater et al., 2011). We used ten on demand generated ASTER DTMs (product AST14DEM) from five different acquisition dates of the year 2008-2010. A small missing stripe was filled with a DTM from 2013 scenes (Table S1, Fig. S1). ASTER scenes were visually checked and from the most promising available (no clouds, minimum snow cover) the respective DTMs were ordered and used for DTM differencing.

To estimate rates of elevation change, an acquisition year was assigned for each glacier. We used the mean time difference for glaciers covered by two scenes acquired in different years (Table S3). Two high-resolution Cartosat-1 stereo scenes captured on 11 July 2010 (Table 1) were used to compare and investigate the consistency of the results obtained with the lower resolution ASTER DTM. Cartosat-1 (IRS-P5) was launched by Indian Space Research Organisation (ISRO) in May 2005. The satellite has two high resolution (2.5 m) panchromatic sensors recording stereo images along the track (Titarov, 2008). The major advantage of this dataset, besides the high spatial resolution, is the 12 bit radiometric resolution. Unfortunately, the spatial coverage is relatively small (25 x 25 km) and our two stereo pairs cover only one large glacier (Khurdopin Glacier, glacier nr. 6 in Figure 2) in full. Declassified Hexagon KH-9 imagery which has a ground resolution of about 8 m and a coverage of about 250 x 125 km allowed us to extend the time series back to 1973 (Table S1). The KH9-Hexagon mission was part of the US keyhole reconnaissance satellite program whose images were declassified in 2002. Imagery from this program have already been applied to investigate glacier mass changes (e.g. Pieczonka et al., 2013). The ICIMOD glacier inventory (Bajracharya and Shrestha, 2011), also available through the GLIMS database (www.glims.org), was used as a baseline data set and manually adjusted based on the utilized optical imagery for DTM generation and Landsat ETM+ scenes of the years 2000 and 2001.

2.2 DTM generation, postprocessing, differencing and uncertainty

All KH-9 DTMs were generated with Erdas Imagine 2014 Photogrammetry Suite using the frame camera model with a focal length of 30.5 cm. Image pre-processing includes the elimination of internal distortions based on the regularly distributed réseau crosses (originally included to correct film distortion effects) and their removal thereafter, following Pieczonka et al.

(2013). GCPs were collected from Landsat 7 ETM+ imagery with SRTM1 as a vertical reference (Pieczonka et al., 2013). Though GCP collection in rough terrain is challenging, we were able to identify 26/28 GCPs (Table S2) located at mountain peaks, large terrain features, and bridges. GCPs were distributed throughout the scenes and at different elevations.

Fiducials were measured manually considering the principal point in the image centre. All stereo images have been processed with a RMS of $< \sim 1.5$ pixels (Table S2). The final Hexagon KH-9 DTMs cover the entire Hunza basin. A small gap of about 20 pixels exists between the two generated DTMs.

The Cartosat-1 stereo pairs have also been processed using PCI Orthoengine 2014 with 32 and 35 GCPs. To improve the quality of the DTMs, image enhancement techniques were applied prior to DTM generation in order to overcome low image contrast and temporal differences in image acquisition. The root mean squared error (RSME) varied between 0.3 and 3.9 pixels (Table S2). The spatial resolution of all DTMs was chosen as 30 m.

In order to obtain reliable results on glacier surface elevation changes, the DTMs must be properly co-registered (Nuth and Kääb, 2011). As we observed tilts when differencing the original DTMs, we first minimized elevation differences between the different DTMs with respect to the SRTM1 master DTM by applying a first order trend correction. We considered only elevation differences (Δh) between ± 150 m over non-glacierized terrain with slopes less than 15° (Bolch et al., 2008; Pieczonka et al., 2013). Subsequently, all DTMs were further co-registered following the approach by Nuth and Kääb (2011). **The final displacement between all DTMs and SRTM1 were less than or equal to one pixel (≤ 30 m) on average.**

Each individual older co-registered DTM was then subtracted from the more recent DTM. The difference grids were finally mosaicked to facilitate processing. We used a **weighted mean elevation change**, based on area of coverage, for the overlapping ASTER DTMs (Fig. S1).

Data voids and mismatches that result in incorrect elevation values can occur in areas with low image contrast such as cast shadows and bright snow. Mismatches due to snow in the accumulation regions led often to unrealistic low elevation values using KH-9 data that would subsequently lead to unrealistic surface lowering values in parts of the accumulation region (Pieczonka and Bolch, 2015). However, thickness change distributions for glaciers with negative mass budgets typically have a minimum lowering at the glacier head with increasing values towards the glacier front following a non-linear trend (Huss et al., 2010). This pattern is different for surging glaciers that often exhibit high positive Δh values at comparatively low elevations, strong surface lowering around the ELA, and then decreasing Δh values towards the upper reaches (Gardelle et al. 2013, Rankl and Braun, 2016). Elevation change patterns can also be affected by thick debris cover where the highest lowering usually does not occur close to the terminus but upstream (Bolch et al., 2008).

As both debris-covered glaciers and surge-type glaciers are common in the investigated region we could not apply a general threshold to remove Δh outliers, but used the general assumption that lower elevations show stronger Δh variability than higher elevations. This should still be true for surging glaciers and those with balanced conditions. The related calculations followed the approach by Pieczonka and Bolch (2015) and used a sigmoid function allowing a larger range of Δh values in the middle part of the ablation region to preserve the signal of surging glaciers and a narrower range ($-20 \leq \Delta h \leq 20$ m) at the

glacier head. We filled all data gaps (including the gap between the two KH9 DTMs) by means of ordinary kriging in order to get the weighted moving average based on neighbouring pixel values.

The penetration of the radar beams into firn and snow has to be considered in case of the comparison of DTMs generated from microwave data such as the SRTM1. However, the value can only be estimated as it depends on several unknown parameters (e.g. snow depth and characteristics) and is therefore one major source of uncertainty (Kääb et al., 2015; Gardelle et al., 2013). We applied the correction suggested by Kääb et al. (2012), who analysed the beam penetration of the C-band SRTM data in a similar region of the Karakoram and found a penetration of 2.4 ± 1.4 m. The conversion of volume to mass changes needs to consider the combined ice and snow density. As both are unknown, we used a density of 850 ± 60 kg/m³ as a reasonable and widely used assumption for a longer time period (Huss, 2013).

10 There is no best method to estimate the uncertainty (ϵ) of the DEM differencing when no precise and well distributed GCPs are available. A first estimate provides the standard deviation of the non-glacierized terrain which can serve as a first estimate and is 22m for the difference between the KH9 and the SRTM DEM, 24 m (SRTM-ASTER DEM), and 26 m (KH9-ASTER DEM). However, the standard deviation can be significantly higher than the real uncertainty as the spatial correlation is not considered (Rolstad et al., 2009). Therefore, we followed the widely applied approach developed by
 15 Gardelle et al., (2013):

$$E_{\Delta h} = \frac{E_{\Delta h_i}}{\sqrt{N_{eff}}} \quad (1)$$

where $E_{\Delta h_i}$ is the standard deviation of the mean elevation change of the non-glacierized terrain of each altitude band and
 20 N_{eff} is the effective number of observations. The latter is calculated using the total number of observations N_{tot} , the pixel size R (30 m), and d is the distance of spatial autocorrelation of the elevation change maps (1025 m)

$$N_{eff} = \frac{N_{tot} \cdot R}{2d} \quad (2)$$

25 The overall uncertainty of the DEM difference is the average of $E_{\Delta h}$ weighted by the glacier hypsometry. A further uncertainty to be considered is the uncertainty of the mapped area of the glaciers E_a . We assume an uncertainty of 5% which is towards the upper bound of published estimates of the uncertainty of mapped glaciers based on similar satellite data (e.g. Paul et al. 2013) taken into consideration that the delineation of debris-covered and avalanche-fed glaciers which are both common in the study region is more difficult. The final uncertainty is calculated considering also the uncertainty of the radar
 30 penetration (E_p , ± 1.4 m) and of the volume to mass conversion (E_m , $\pm 7\%$ of the elevation change).

$$E = E_{\Delta h} + E_a + E_p + E_m \quad (3)$$

We did not apply a seasonality correction as most of the images were acquired close to the end of the ablation period but assume the effect is well within the considered uncertainty.

3. Results and Discussion

3.1 Glacier volume and mass changes, surge-type glaciers

5 Results for the period 1999 to ~2009 show heterogeneous glacier behaviour, with several surging glaciers in the study region (Figure 2). A northern tributary of Hispar Glacier thickened by approximately 150 m at the confluence of the glaciers. Khurdopin and its neighbouring glaciers in south-eastern Shimshal Valley both show significant thickening and thinning within their tongues which is typical of a surge occurring during the study period (Figure 2). In contrast, the large debris-covered Batura Glacier west of Shimshal Valley showed surface lowering throughout the tongue leading to an overall
10 volume loss (Figures 2, 3). For the entire study region we found no significant mass changes (Table 1) which is in line with previous results for the period after 1999 in Central Karakoram (e.g. Gardelle et al., 2013). In addition, we add information for glaciers east of their study region (e.g. Batura Glacier) and cover the entire Hispar Glacier, one of the largest glaciers (length 50 km) in the Karakoram. For this region we cannot confirm positive mass budgets but our study indicates a slight mass loss similar to Rankl and Braun (2016) and Kääb et al. (2015). However, our uncertainties are larger due to the
15 utilization of the lower resolution, ~~but freely available~~, ASTER DTM. In addition, we emphasize that the analysed glaciers and period slightly differ which could be a reason for the (not significant) differences to the existing studies.




Figure 2: Elevation difference between the ASTER and Hexagon KH9 DTMs (above), the Hexagon KH-9 and the SRTM1 DTMs (below, left) and the SRTM1 and ASTER DTMs (below, right). Black dots indicate surge-type glaciers, the numbers selected larger glaciers: 1: Batura, 2: Pasu, 3: Barpu, 4: Hispar, 5: Yazghil, 6: Khurdopin, 7: Vajerab

5



Our extended time series show that glaciers in Hunza Valley experienced no significant mass changes but a heterogeneous behaviour also for the period 1973 to 1999 (Table 1). Hence, central Karakoram glaciers were, on average, in balance for at

least the last 40 years. Although longer-term balanced budgets could be assumed based on existing information about (insignificant) area changes since the 1970s (Bhambri et al., 2013), on average similar debris coverage since 1977 (Herreid et al., 2015), we confirm **for the first time** on average balanced-budgets since 1973 using elevation differencing.

5 Most glaciers experienced similar mass budgets for both investigation periods. However, it seems that some glaciers had more negative budgets post-2000. This is especially true for the debris-covered Batura Glacier whose tongue showed significant lowering during 1999-2009. Surge-type glaciers also showed more negative values in the recent period. Kurdophin Glacier, for example, experienced a significant thickening near the snout and a significant lowering around the ELA, both combined resulting in an about zero mass budget for 1973 and 1999 and significant lowering at the lower part of
 10 the tongue leading probably to a mass loss for 1999 – 2009 (Figures 2, 3). These elevation change characteristics could be due the fact that the active surge event transferred ice mass to lower elevations where it is now more prone to melting. The very high surface lowering of the middle part of the tongue of Hispar Glacier hints also to a past surge event. For the recent period, different tributaries of Hispar Glacier show clear signs of surging with significant elevation gains (Figure 2). Overall, we identified 28 surge-type glaciers (including 5 tributaries) based on the DTM differencing results in combination with
 15 morphological features like looped moraines or heavily crevassed glacier surfaces (Fig. 2). We cannot exclude the possibility that a few surge-type glaciers were missed. However, most of the surge events should be covered by our study period of almost 40 years as the surge periodicity in the Karakoram is rather short with averages between ~25 and 40 years (Copland et al. 2011). Over the entire study period there is no significant difference in the mass budgets of surge-type and non-surge-type glaciers, a result also found by Gardelle et al. (2013).

20

Table 1: Glacier mass balance for different periods and glacier types

Nr.	Glacier Name	Glacier Type	Area (km ²)	Glacier Mass Balance (m w.e. a ⁻¹)		
				1973 - 1999	1999 – ca. 2009	1973 – ca. 2009
1	Batura	Debris covered	236	0.00 ± 0.10	-0.39 ± 0.26	-0.12 ± 0.09
2	Pasu	Debris free	51	+0.05 ± 0.11	-0.13 ± 0.26	-0.09 ± 0.10
3	Barpu	Surge type	90	+0.03 ± 0.08	-0.10 ± 0.18	-0.15 ± 0.08
4	Hispar	Surge-type Debris covered	345	-0.10 ± 0.08	-0.11 ± 0.21	-0.14 ± 0.08
5	Yazghil	Debris covered	99	-0.02 ± 0.13	-0.04 ± 0.32	-0.01 ± 0.12
6	Khurdopin	Surge-type Debris covered	115	-0.05 ± 0.09	-0.14 ± 0.22	-0.05 ± 0.08

7	Vijerab	Surge-type Debris covered	113	+0.03 ± 0.10	-0.31 ± 0.25	-0.22 ± 0.09
	Whole region		2868	-0.01±0.09	-0.08 ± 0.21	-0.06 ± 0.08
	Whole region	Non-surge-type	2237	0.00 ± 0.08	-0.03 ± 0.22	-0.03± 0.08
	Whole region	Surge-type	631	-0.03 ± 0.10	-0.15 ± 0.30	-0.09 ± 0.09

3.2 DTM generation and sources of uncertainty

Declassified KH-9 Hexagon data have proven to be valuable for assessing geodetic glacier mass budgets (Pieczonka and Bolch, 2015; Pieczonka et al., 2013, Mauer et al. 2016). The main challenges in obtaining accurate results are
5 miscorrelations in the accumulation regions of glaciers and significant tilts and shifts of elevation trends making careful co-registration and post-processing necessary. This leads to higher uncertainties compared to more recent data with a similar spatial resolution. The glacier volume changes calculated based on the automatically derived ASTER DTMs (AST14DEM) and the SRTM DTM were similar to those using better quality higher resolution SPOT5 DTMs for a similar region (Gardelle et al., 2013). The DTM differences between the utilized DTMs of the off-glacier area shows the in general good agreement
10 but also regions with higher deviations especially for the western part where the quality of the ASTER DEMs were lower (Fig. S1). We found no significant difference between the mass budget results of Khurdopin Glacier calculated using Cartosat-1 data and the values calculated using the ASTER data (-0.14 ± 0.21 for 1999-2009 vs. -0.16 ± 0.13 for 1999 – 2010). This gives confidence that the ASTER DTMs which can be freely obtained for scientific purposes are of high value to calculate glacier volume and mass changes over a longer period of time as also shown for other regions (e.g. Berthier et al.,
15 2016).

A further source of uncertainty are the data voids in the original SRTM data and voids due to the outlier filtering. About 20% of the total glacierized area for all analysed periods were identified as outliers and filled afterward by kriging interpolation. The voids are almost entirely located in the accumulation region where surface elevation changes are relatively small (e.g. Schwitter & Raymond, 1993) and where we restricted the maximum possible deviation. To assess the influence of the filling
20 on the result we calculated the elevation change value (a) without void filling (result: mean elevation difference -3.80 m or -0.09 m w.e. a^{-1}), (b) filling with zero (-2.98 m or -0.07 m w.e. a^{-1}), or (c) our applied interpolation method (-2.39 m or -0.06 m w.e. a^{-1}). The results show that average elevation change rates do not change significantly and the deviations are well within the uncertainty. About 10% of the glacierized area in our study region is affected by data voids in the original SRTM data and previous studies showed that there can be significant deviations to reality in the data used to fill the voids (e.g. Kääb
25 et al. 2012). In addition, the time stamp of the data is often unknown. We compared therefore the results of the void filled and the non-void filled version. Using the latter we calculated the surface elevation change for the existing pixels only. The resultant value of the mean surface elevation change for both periods differs only by about 0.02 m a^{-1} , which is also well

within the estimated uncertainty. The minor differences are, besides the possibility that the ASTER GDEM data used for void filling are quite reliable for the study region, due to the fact that most of the voids are located in the accumulation regions where we restricted the maximum possible deviation. Only one larger glacier is also affected in the ablation region. This is Shishpar Glacier, a southerly facing exposed glacier, located south of Batura Glacier. The void-filled data allowed for

5 detection of the surge activity between 1973 and 1999 with an estimated mass budget of $+0.04 \pm 0.19$ m w.e. a⁻¹.




Figure 3: Longitudinal profiles of surface elevation changes for selected glaciers for the two periods.

10 One of the major sources of uncertainty is the penetration of the radar beam into snow and ice when using the SRTM DTM. Gardelle et al. (2013) estimated a mean penetration of 3.4 m with values up to more than 9 m in the accumulation region. This value is higher than the 2.4 m we applied here following Kääb et al. (2012). However, applying the higher penetration value would only lead to a slight difference in mass change of $+0.03$ m w.e. a⁻¹. The average surface elevation change without considering any radar penetration is -0.10 m a⁻¹ for the period 1973-1999 and $+0.15$ m a⁻¹ for the period 1999-2009.

15 However, penetration is not an issue for calculation of the geodetic mass budget over the entire period (~1973-2009) as this uses optical data only. As the results of the individual periods are in good agreement with the values for the entire period, we are confident in the reliability of all mass balance calculations.

4 Conclusion

Although longer-term balanced budgets in the Karakoram region could be assumed based on existing information, such as

20 insignificant changes in glacier area or debris-covered area since the 1970s, confirm based on 1973 Hexagon and ~2009 ASTER DEM data that balanced-budgets in the investigated region are not a recent phenomenon. Since at least the 1970s, glaciers in the Hunza have experienced on average no significant overall mass changes. This is important information as glacier area and length changes represent an indirect and delayed response to climate change and deriving a clear cause and

effect relation is difficult to establish. On the other hand, during a surge, ice is transported rapidly from an upper reservoir zone to the ablation region and vice versa during the quiescent phase, albeit on an often much longer time scale. With the 40-year time-period considered here it was possible to show that the overall mass change remains about the same but that significant differences exist for individual glaciers.

5

Author Contributions

T.B. designed the study, performed all analysis, generated the figures and wrote the draft of the manuscript. K.M. generated the raw Hexagon and Cartosat-1 DTMs and co-registered the data. T.B. and T.P. supported the generation of the DTMs and the co-registration. J.S. contributed to the study design, and all authors contributed to the final form of the article.

10 **Acknowledgements**

This study was performed within the Cryosphere Initiative of the International Center for Integrated Mountain Development (ICIMOD), with the targeted support of the UK Department for International Development (DFID). ICIMOD is funded in part by the governments of Afghanistan, Bangladesh, Bhutan, China, India, Myanmar, Nepal, and Pakistan, and the Cryosphere Initiative is funded by the Norwegian Ministry of Foreign affairs. The views expressed are those of the authors and do not necessarily reflect their organizations or funding institutions. T. Bolch acknowledges funding through the ESA project Glaciers_cci (4000109873/14/I-NB), Deutsche Forschungsgemeinschaft (DFG) and The University of Colorado CHARIS project (Contribution to High Asia Runoff from Ice and Snow – (nsidc.org/charis) funded by the United States Agency for International Development We thank F. Paul for his comments on the manuscript.

References

- 20 Bajracharya, S. R. and Shrestha, B. R. (Eds.): The status of glaciers in the Hindu Kush-Himalayan Region, Kathmandu, 127 pp., 2011.
- Berthier, E., Arnaud, Y., Vincent, C. and Rémy, F.: Biases of SRTM in high-mountain areas. Implications for the monitoring of glacier volume changes, *Geophys. Res. Lett.*, 33, L08502, doi:10.1029/2006GL025862, 2006.
- Berthier, E., Cabot, V., Vincent, C. and Six, D.: Decadal region-wide and glacier-wide mass balances derived from multi-
25 temporal ASTER satellite digital elevation models. Validation over the Mont-Blanc area, *Frontiers in Earth Science*, 4, 2016, doi: 10.3389/feart.2016.00063.
- Bhambri, R., Bolch, T., Kawishwar, P., Dobhal, D., Srivastava, D. and Pratap, B.: Heterogeneity in glacier response in the Shyok valley, northeast Karakoram, *Cryosphere*, 7, 1384–1398, 2013, doi:10.5194/tc-7-1385-2013.
- Bolch, T., Buchroithner, M. F., Pieczonka, T. and Kunert, A.: Planimetric and volumetric glacier changes in Khumbu
30 Himalaya since 1962 using Corona, Landsat TM and ASTER data, *J. Glaciol.*, 54, 592–600, 2008.

- Bolch, T., Kulkarni, A., Kääb, A., Huggel, C., Paul, F., Cogley, J. G., Frey, H., Kargel, J. S., Fujita, K., Scheel, M., Bajracharya, S. and Stoffel, M.: The state and fate of Himalayan glaciers, *Science*, 336, 310–314, 2012, doi: 10.1126/science.1215828.
- Copland, L., Sylvestre, T., Bishop, M. P., Shroder, J. F., Seong, Y. B., Owen, L. A., Bush, A. and Kamp, U.: Expanded and recently increased glacier surging in the Karakoram, *Arct. Antarct. Alp. Res.*, 43, 503–516, 2011, doi: 10.1657/1938-4246-43.4.503
- Fowler, H. and Archer, D.: Conflicting signals of climate change in the Upper Indus Basin, *J. Climate*, 19, 4276–4293, 2006.
- Gardelle, J., Berthier, E., Arnaud, Y. and Kääb, A.: Region-wide glacier mass balances over the Pamir-Karakoram-Himalaya during 1999–2011, *Cryosphere*, 7, 1263–1286, 2013, doi:10.5194/tc-7-1263-2013.
- Herreid, S., Pellicciotti, F., Ayala, A. and Chesnokova, A.: Satellite observations show no net change in the percentage of supraglacial debris-covered area in northern Pakistan from 1977 to 2014, *J. Glaciol.*, 61, 524–536, 2015, doi: 10.3189/2015JoG14J227.
- Hewitt, K.: Glacier change, concentration, and elevation effects in the Karakoram Himalaya, Upper Indus Basin, *Mount. Res. Dev.*, 33, 188–200, 2011, doi: 10.1659/MRD-JOURNAL-D-11-00020.1.
- Huss, M.: Density assumptions for converting geodetic glacier volume change to mass change, *Cryosphere*, 7, 877–887, 2013, 10.5194/tc-7-877-2013.
- Huss, M., Joutet, G., Farinotti, D. and Bauder, A.: Future high-mountain hydrology: a new parameterization of glacier retreat, *Hydrol. Earth Syst. Sci.*, 14, 815–829, 2010, 10.5194/hess-14-815-2010.
- Immerzeel, W. W., van Beek, L. P. and Bierkens, M. F.: Climate change will affect the Asian water towers, *Science*, 328, 1382–1385, 2010, doi: 10.1126/science.1183188.
- Kääb, A., Berthier, E., Nuth, C., Gardelle, J. and Arnaud, Y.: Contrasting patterns of early twenty-first-century glacier mass change in the Himalayas, *Nature*, 488, 495–498, 2012, doi: 10.1038/nature11324.
- Kääb, A., Treichler, D., Nuth, C. and Berthier, E.: Brief Communication: Contending estimates of 2003–2008 glacier mass balance over the Pamir–Karakoram–Himalaya, *Cryosphere*, 557–564, 2015, doi:10.5194/tc-9-557-2015.
- Maurer, J., Rupper, S. and Schaefer, J. M.: Quantifying ice loss in the eastern Himalayas since 1974 using declassified spy satellite imagery, *Cryosphere*, 10, 2203–2215, 2016.
- Nuth, C. and Kääb, A.: Co-registration and bias corrections of satellite elevation data sets for quantifying glacier thickness change, *Cryosphere*, 5, 271–290, 2011, 10.5194/tc-5-271-2011.
- Paul, F. and Haeberli, W.: Spatial variability of glacier elevation changes in the Swiss Alps obtained from two digital elevation models, *Geophys. Res. Lett.*, 35, L21502, doi:10.1029/2008GL034718, 2008.
- Paul, F., Barrant, N., Baumann, S., Berthier, E., Bolch, T., Casey, K. A., Frey, H., Joshi, S. P., Kononov, V., LeBris, R., Mölg, N., Nosenko, G., Nuth, C., Pope, A., Racoviteanu, A., Rastner, P., Raup, B., Scharrer, K., Steffen, S. and Winsvold, S.: On the accuracy of glacier outlines derived from remote sensing data, *Ann. Glaciol.*, 54, 171–182, 2013, doi: 10.3189/2013AoG63A296.

- Pieczonka, T. and Bolch, T.: Region-wide glacier mass budgets and area changes for the Central Tien Shan between ~1975 and 1999 using Hexagon KH-9 imagery, *Global Planet. Change*, 128, 1–13, 2015, doi: 10.1016/j.gloplacha.2014.11.014.
- Pieczonka, T., Bolch, T., Wei, J. and Liu, S.: Heterogeneous mass loss of glaciers in the Aksu-Tarim Catchment (Central Tien Shan) revealed by 1976 KH-9 Hexagon and 2009 SPOT-5 stereo imagery, *Remote Sens. Environ.*, 130, 233–244, 2013, 10.1016/j.rse.2012.11.020.
- 5 Quincey, D. J. and Luckman, A.: Brief Communication: On the magnitude and frequency of Khurdopin glacier surge events, *Cryosphere*, 8, 571–574, 2014, doi:10.5194/tc-8-571-2014.
- Rankl, M. and Braun, M.: Glacier elevation and mass changes over the central Karakoram region estimated from TanDEM-X and SRTM/X-SAR digital elevation models, *Ann. Glaciol.*, 57, 273–281, 2016, doi: 10.3189/2016AoG71A024.
- 10 Rankl, M., Kienholz, C. and Braun, M.: Glacier changes in the Karakoram region mapped by multitemission satellite imagery, *Cryosphere*, 8, 977–989, 2014, doi: 10.5194/tc-8-977-2014.
- Rolstad, C., Haug, T. and Denby, B.: Spatially integrated geodetic glacier mass balance and its uncertainty based on geostatistical analysis: application to the western Svartisen ice cap, Norway, *J. Glaciol.*, 55, 666–680, 2009.
- Slater, J. A., Healy, B., Kroenung, G., Curtis, W. and Haase, J.: Global assessment of the new ASTER Global Digital Elevation Model, *Photogramm. Eng. Remote Sensing*, 77, 335–349, 2011.
- 15 Titarov, P. S.: Evaluation of Cartosat-1 geometric potential, *International Archives of Photogrammetry, Remote Sensing and Spatial Information Sciences*, 37, 841–846, 2008.