

“Heterogeneous glacier thinning patterns over the last 40 years in Langtang Himal, Nepal”

by S. Ragetti, T. Bolch and F. Pellicciotti

Dear Editor,

Thank you for the careful reading of our manuscript and for the comments. Our point-by-point responses are provided below.

With kind regards,

Silvan Ragetti,

Editor’s technical comments:

Title. Add that this is in Nepal?

It is now specified in the title that Langtang Himal is in Nepal.

Abstract L20. Is "identify" the best verb? "measure"? "compute"?

We have replaced “identify” by “compute” as suggested.

Effect of the earthquake. It has also been studied using SPOT6/7 data by Lacroix, 2016.

Probably worth mentioning this study together with Kargel et al., 2015.

Lacroix, P.: Landslides triggered by the Gorkha earthquake in the Langtang valley, volumes and initiation processes, Earth, Planets and Space, 68(1), 1–10, doi:10.1186/s40623-016-0423-3, 2016.

This is a good suggestion. We now also cite Lacroix 2016.

P8 L11. Space missing

Ok. Thank you for noticing.

P11 L27. I think "until" can be deleted

We agree and have removed “until”.

Everywhere: when you have two or more "glaciers", I do not think a cap letter is needed for "glaciers"

Both capital letters or small letters after naming two or more “glaciers” are used in the literature. We prefer to consistently use capital letters.

P17 L25. suggestion: "During the same two time intervals debris-free Kimoshung Glacier" ("debris-free" added for the sake of parallelism with Yala Glacier and to remind this fact to the reader)

Ok. We have added “debris-free” here.

P18 L25, L29. Not so easy for the reader to understand what you mean by "here" (the present study? the glacier described?). Maybe you could clarify a bit.

We have clarified this and removed “here” in both cases.

P20. L17. Maybe you could add that this is because you are only considering time period of at least 4 years in the ensemble? Then the impact is at most $0.52/4=0.16$ just above 0.12 m/yr? Otherwise it is not really clear how 0.52 m is compared to 0.12 m/yr.

This is now clarified. We now state “(... if divided over *six years, which is the shortest time interval of any $\Delta h/\Delta t$ map in the ensemble involving the October 2015 DEM*).

P21. L10. (and elsewhere) "volume loss rate". You are often using both "thinning rate" and "volume change rate" as if they were the same. I think in the context of your paper/results, it would be easier for the reader if you could stick to a single terminology (thinning rate). You could legitimately answer me that this is just a matter of dividing by the area and you would be mathematically right. But unifying your terminology (i.e. text = figure/table) will, I think, help the reader to really extract the real message.

Ok. We have replaced “volume loss rate” consistently by “thinning rate” everywhere in the text.

P26 L26. I think it would be more accurate to say here "for the same 10 glaciers as Bolch et al. (2011) in the Everest region" (because the Gardelle et al., 2013 mass balance for the whole Khumbu area is less negative than for these 10 glaciers)

We have clarified the sentence as suggested.

P27 L3. I do not think you need to define DEM again in the conclusion.

We agree. We have changed this.

Figure 2 legend. "The central marks correspond to the median" is repeated twice

We have rephrased the legend to prevent repetitions.

Figure 7. The dark blue line for the 1974-2006 dh/dt rate is not continuous. Probably because the time scale is not linear. To be explained in the legend.

We have added a sentence to clarify this (also in the legend of Figure 8): “*Note that the 1974-2006 time scale is not linear (dashed dark blue line).*”

P45 L4. " uncertainty in function" --> " uncertainty as a function" (I think). Same for figure 10.

We have corrected this as suggested.

P47. L6. data from Figure 11. Do you mean Figure 13?

We have corrected this. Thank you for noticing.

Heterogeneous glacier thinning patterns over the last 40 years in Langtang Himal, Nepal

S. Ragettli^{1,2}, T. Bolch^{2,3} and F. Pellicciotti⁴

[1]{Institute of Environmental Engineering, ETH Zürich, Switzerland}

[2]{University of Zurich, Department of Geography, Zurich, Switzerland}

[3]{Institute for Cartography, Technische Universität Dresden, Dresden, Germany}

[4]{Northumbria University, Department of Geography, Newcastle upon Tyne, UK}

Correspondence to: S. Ragettli (ragettli@ifu.baug.ethz.ch)

Abstract

This study presents volume and mass changes of seven glaciers (five partially debris-covered, two debris-free) in the upper Langtang catchment in Nepal. We use a digital elevation model (DEM) from 1974 stereo Hexagon satellite data and seven DEMs derived from 2006-2015 stereo or tri-stereo satellite imagery (e.g. SPOT6/7). The availability of multiple independent DEM differences allows identifying a robust signal and narrowing down the uncertainty about recent volume changes. The volume changes calculated over several multi-year periods between 2006 and 2015 consistently indicate that glacier thinning has accelerated with respect to the period 1974-2006. We calculate an ensemble-mean elevation change rate of $-0.45 \pm 0.18 \text{ m a}^{-1}$ for 2006-2015, while for the period 1974-2006 we identifycompute a rate of $-0.24 \pm 0.08 \text{ m a}^{-1}$. However, the behavior of glaciers in the study area is heterogeneous, and the presence or absence of debris does not seem to be a good predictor for mass balance trends. Debris-covered tongues have non-linear thinning profiles, and we show that recent accelerations in thinning correlate with the presence of supraglacial cliffs and lakes. At stagnating glacier areas near the glacier front, on the other hand, thinning rates decreased with time or remained constant. The April 2015 Nepal earthquake triggered large avalanches in the study catchment. Analysis of two post-earthquake DEMs revealed that the avalanche deposit volumes remaining six months after the earthquake are negligible in comparison to 2006-2015

1 elevation changes. However, the deposits compensate about 40% the mass loss of debris-
2 covered tongues of one average year.

3 **1 Introduction**

4 Atmospheric warming has caused widespread recent glacier thinning and retreat in the
5 Himalayan region (Bolch et al., 2012). The impact of current and future glacier changes on
6 Himalayan hydrology and downstream water supply strongly depends on the rate of such
7 changes. However, planimetric and volumetric glacier changes are difficult to characterize
8 due to limited data availability, and many recent studies have highlighted the spatially
9 heterogeneous distribution of glacier wastage in the Himalayas (Fujita and Nuimura, 2011;
10 Bolch et al., 2012; Kääb et al., 2012). Prominent examples of current-day regional differences
11 in glacier evolution across the Hindu Kush–Karakoram–Himalaya (HKH) are the reported
12 positive glacier mass balances in the Pamir and Karakoram. Glaciers in the rest of the HKH
13 are thinning and receding (Bolch et al., 2012; Kääb et al., 2012; Gardelle et al., 2013). Across
14 regions, differences in recent glacier evolution can often be associated to differences in
15 climatic regimes (Fujita, 2008), particularly to the varying influence of the south Asian
16 monsoon and westerly disturbances (Yao et al., 2012). However, also within the same
17 climatic region the rate of glacier changes can be heterogeneous (Scherler et al., 2011b). A
18 main focus of current research is on the effect of supraglacial debris-cover on glacier response
19 to climate. Thick debris cover is a common feature in the HKH (Scherler et al., 2011b;
20 Racoviteanu et al., 2015) and a homogenous layer of thick debris effectively reduces melt
21 rates of underlying ice (e.g. Östrem, 1959; Mattson et al., 1993). However, the
22 characterization of debris-covered glacier response to climate is complicated by the frequent
23 occurrence of ice cliffs and supraglacial lakes. At exposed cliffs, melt rates are much higher
24 compared to the ice covered by a thick debris mantle (Sakai et al., 1998, 2002; Immerzeel et
25 al., 2014a; Steiner et al., 2015; Buri et al., 2016), and also at supraglacial ponds energy
26 absorption is several times larger than that at the surrounding debris-covered surface (Sakai et
27 al., 2000; Miles et al., 2016a). Recent large-scale geodetic studies based on remote sensing
28 have provided evidence that the present-day surface lowering rates of some debris-covered
29 areas in the HKH might be similar to those of debris-free areas even within the same
30 altitudinal range (Kääb et al., 2012; Nuimura et al., 2012; Gardelle et al., 2013), and surmise
31 this could be due to enhanced melt from exposed ice cliffs and supraglacial lakes. Several
32 detailed modelling studies on the other hand have provided evidence for a melt reducing

1 effect of debris at the glacier scale (e.g. Juen et al., 2014; Ragettli et al., 2015), and have
2 shown how supraglacial debris prolongs the response of the glacier to warming (Banerjee and
3 Shankar, 2013; Rowan et al., 2015). Discrepancies between the different conclusions may be
4 associated to glacier samples that are not comparable or to model uncertainties (particularly
5 regarding the representation of the effect of supraglacial cliffs and lakes on total melt).
6 Models can also provide actual melt rates while geodetic studies only provide glacier thinning
7 rates, which are affected by glacier emergence velocity.

8 Programs to monitor debris-covered glaciers have been initiated in the Karakorum (e.g.
9 Mayer et al., 2006; Mihalcea et al., 2006, 2008) and in the Central Himalaya (e.g. Pratap et
10 al., 2015; Ragettli et al., 2015). Yet, due to the logistical and financial constraints, long-term
11 mass balance measurements are basically inexistent in the HKH. To document changes in
12 debris-covered glacier thinning over time, declassified high-resolution reconnaissance
13 satellite data available from the 1960s and 1970s are an important source of information
14 (Bolch et al., 2008, 2011; Maurer and Rupper, 2015). However, a common problem of
15 previous multi-temporal geodetic studies is the relatively low statistical significance of
16 detected changes in glacier thinning over time. The uncertainties in digital elevation models
17 (DEMs) derived from optical data and mass change measurements by DEM differentiation in
18 the HKH arise from the difficult conditions for photogrammetric elevation analysis (due to
19 extreme topography, surfaces with low contrast like bright snow cover or cast shadows).
20 Radar derived DEMs provide more accurate results in snow covered areas but have even
21 higher problems in steep terrain due to the side looking geometry. Unknown penetration of
22 the radar beam into snow and ice is another shortcoming of DEMs derived from radar data.
23 Uncertainties in volume loss estimates are therefore usually higher than identified acceleration
24 in glacier thinning (Nuimura et al., 2012). The uncertainties are especially high over short
25 periods. For long periods with much larger absolute elevation changes, the effect of errors in
26 the DEMs weighs less and uncertainties in glacier volume changes are lower.

27 This study presents volume and mass changes of seven glaciers (five partially debris-covered,
28 two debris-free) in the upper Langtang catchment in Nepal. The aim of this study is to
29 determine changes in thinning rates with high confidence by considering multiple
30 independent DEM differences for short periods. For this we use seven DEMs derived from
31 2006-2015 stereo or tri-stereo satellite imagery and one DEM obtained from 1974 stereo
32 Hexagon satellite data. We obtain an ensemble of multi-annual elevation changes that

1 provides a range of plausible values for the period between October 2006 and October 2015.
2 We then assess if the elevation changes between different overlapping periods between 2006
3 and 2015 show similar characteristics. If this is the case, the ensemble of results can be used
4 to identify statistically significant changes in ~~volume loss~~thinning rates with respect to the
5 longer period 1974-2006. Three main research questions are then addressed. First, we assess
6 if overall thinning of glaciers in the region has accelerated. Second, we determine if spatial
7 thinning patterns have changed over time. To explain changes in thinning rates we derive a
8 number of glacier surface properties and glacier surface velocities. Third, we evaluate if there
9 are major differences between the response of debris-covered and debris-free glaciers in the
10 sample. Finally, we also look at the cryospheric impact of the April 2015 Nepal earthquake
11 (7.8 magnitude, epicenter approximately 80 km west of the Langtang Valley). The earthquake
12 devastated large parts of the Langtang catchment by triggering large avalanches (~~Kargel et al.,~~
13 2016)-and landslides (Kargel et al., 2016; Lacroix, 2016). Two post-earthquake DEMs from
14 May and October 2015 are used to quantify the impact of the avalanche events on the mass
15 balance of the debris-covered glacier tongues and assess its significance in comparison to
16 multi-annual volume changes.

17

18 **2 Study Site**

19 We analyze the seven largest glaciers in the Langtang Valley (Langtang, Langshisha,
20 Shalbachum, Lirung, Ghanna, Yala, Kimoshung), located in the monsoon-dominated Central
21 Himalaya in Nepal, approximately 50 km north of Kathmandu and 100 km west of the
22 Everest region. While Yala and Kimoshung Glaciers are debris-free glaciers, all other studied
23 glaciers have tongues that are almost entirely covered by supraglacial debris (Figure 1).
24 Langtang Glacier is the largest glacier in the valley with an area of 46.5 km² in 2006 (Table 1)
25 and a total length of approximately 18 km. The smallest glacier is Ghanna Glacier with an
26 area of 1.4 km².

27 Critical debris thicknesses leading to a reduction of melt rates are exceeded over most parts of
28 the debris-covered glacier area (Ragettli et al., 2015). Relatively thin debris appears only at
29 the transition zone between accumulation and ablation area. At Lirung, Shalbachum, Ghanna
30 and Langshisha Glaciers the upper margins of debris-covered sections are located at the foot
31 of steep cirques and icefalls, and transition zones are therefore very short. Ice cliffs and

1 supraglacial ponds increase the heterogeneity of glacier surface characteristics in the
2 Langtang Valley (Pellicciotti et al., 2015).

3 The ablation season of glaciers in the Langtang Valley lasts from April to September. The
4 monsoon season (mid June – September) is at the same time the warmest and the wettest
5 period of the year. Snow cover at the lower elevation of debris-covered glaciers is common
6 only in winter (December – March). However, outside the monsoon period precipitation is
7 limited and winters are rather dry (Collier and Immerzeel, 2015).

8

9 **3 Data and methods**

10 **3.1 Satellite imagery**

11 Multitemporal high-resolution data from different sensors are applied to assess glacier change
12 in the upper Langtang catchment. Each type of remote sensing data employed to calculate
13 glacier elevation changes is listed below. Spatial and radiometric resolutions and base to
14 height (b/h) ratios are provided in Table 2.

- 15 • The oldest data originate from Hexagon KH-9 stereo satellite images from November
16 1974 (Surazakov and Aizen, 2010; Pieczonka et al., 2013; Maurer and Rupper, 2015).
17 These are declassified images from a US reconnaissance satellite program (Burnett,
18 2012).
- 19 • Cartosat-1 is a remote sensing satellite built by the Indian Space Research
20 Organisation (Tiwari et al., 2008). We purchased radiometrically corrected along-track
21 stereo imagery (processed at level ‘ortho-kit’) of the upper Langtang catchment from
22 October 2006 and November 2009. Cartosat-1 data have been previously used for
23 DEM generation e.g. in the Khumbu region in the Nepal Himalaya by Bolch et al.
24 (2011) and Pieczonka et al. (2011).
- 25 • ALOS-PRISM (Advanced Land Observing Satellite - Panchromatic Remote-Sensing
26 Instrument for Stereo Mapping) was an optical sensor mounted on a Japanese satellite
27 system which operated from January 2006 to April 2011 (Bignone and Umakawa,
28 2008; Tadono and Shimada, 2009; Lamsal et al., 2011; Holzer et al., 2015). We

1 purchased a radiometrically calibrated along-track triplet mode scene from December
2 2010.

- 3 • SPOT6/7 (Système pour l’Observation de la Terre) along-track tri-stereo images
4 were acquired upon request in April 2014, May 2015 and October 2015. SPOT6 and 7
5 are the newest satellites of the SPOT series which have been frequently used for
6 geodetic glacier mass balance studies (e.g. Berthier et al., 2007, 2014; Pieczonka et al.,
7 2013). We acquired stereoscopic images in panchromatic mode corrected for
8 radiometric and sensor distortions. Two of the three SPOT6/7 scenes used in this study
9 were acquired in April/May which means that limited amounts of winter snow is still
10 present on the images. However, the imagery has a high spation resolution (1.5 m) and
11 high radiometric depth of 12bit which leads to good correlation results also over
12 snowy parts.
- 13 • Overlapping pairs of high-resolution images acquired by the WorldView-2 and 3
14 satellites in February 2014 provide the basis of 8m DEMs downloaded from
15 <http://www.pgc.umn.edu/elevation> (Noh and Howat, 2015).

16 **3.2 DEMs and elevation changes**

17 **3.2.1 DEM generation**

18 The Hexagon DEM used here was generated for the study by Pellicciotti et al. (2015). We
19 therefore refer to this study for further technical details regarding the Hexagon DEM. The
20 SPOT6/7, Cartosat-1 and ALOS PRISM DEMs were generated for this study using the
21 OrthoEngine module of PCI Geomatica 2015. We used the same parameters for DEM
22 generation as proposed by Berthier et al. (2014) except setting the parameter ‘DEM detail’ to
23 ‘very high’ instead of ‘low’, which provided better results for the rugged debris-covered
24 glacier surfaces. The basis for the georectification were six differential global positioning
25 system (dGPS) points collected on Lirung Glacier on 23 October 2014 (Brun et al., 2016).
26 Because glacier motion and ablation have to be accounted for when using on-glacier dGPS
27 points, we first generated a DEM from an across-track Pléiades stereo image pair from 1 and
28 9 November 2014 using the available dGPS points as ground-control points (GCPs). Glacier
29 melt between 23 October and the acquisition dates of the Pléiades scenes is negligible due to
30 the low temperatures during this period. The horizontal shift due to glacier motion during this

1 period is less than the grid size of the Pléiades image (0.5 m) and is therefore also negligible.
2 Subsequently, we determined 17 off-glacier GCPs on the basis of the Pléiades scene which
3 were then used to derive a DEM from the SPOT6 April 2014 tri-stereo scene. The Pléiades
4 DEM itself is not used in the following to calculate glacier elevation changes since it covers
5 only a small part of the catchment and since only low stereo matching scores were achieved at
6 elevations higher than 4300 m a.s.l. due to snowfall onset between 1 and 9 November 2014.
7 To guarantee high quality GCPs, only pixels with correlation scores higher than 0.7 were
8 considered for GCPs. Since the Pléiades scene covers only about one fourth of the upper
9 Langtang catchment, an additional 60 GCPs were determined on the basis of the April 2014
10 SPOT6 scene for the DEM extraction from the Cartosat-1, ALOS Prism and SPOT7 scenes.
11 In addition to the GCPs, approximately 100 tie points for each scene were used to match
12 stereo pairs before DEM extraction.

13 The WorldView DEMs are 8 m posting DEMs produced using the Surface Extraction with
14 TIN-based Search-space Minimization (SETSM) by Noh and Howat (2015). The WorldView
15 DEMs rely on the satellite positioning model to locate the surface in space. The scenes from
16 February 2015 which provide the basis of the two WorldView DEMs used in this study were
17 acquired only 20 days apart (Table 2) and are adjacent to each other. The Worldview-2 DEM
18 covers the western part of the study catchment and the WorldView-3 DEM the eastern part.
19 Those DEMs were merged for this study and in the following are referred to as one single
20 DEM representative of February 2015.

21 In addition to the DEMs discussed above, the 2000 SRTM (Shuttle Radar Topography
22 Mission) 1 Arc-Second Global DEM (30 m spatial resolution) was used to calculate slopes
23 and accumulation area ratios (AARs) of glaciers (Table 1) and to define 50 m altitude bands.
24 However, the SRTM DEM was not used for DEM differencing because of the uncertainty
25 regarding the penetration depth of the radar signal into snow and ice (Gardelle et al., 2013;
26 Kääb et al., 2015; Pellicciotti et al., 2015). Only DEMs extracted from optical stereo imagery
27 are therefore employed to calculate elevation changes in this study.

28 **3.2.2 Co-registration and DEM differencing**

29 Co-registration of DEM-pairs is applied in order to minimize the errors associated with shifts.
30 Systematic errors in the elevation change maps due to tectonic uplift which could be relevant
31 after the April 2015 Nepal earthquake are also corrected with the co-registration. For this

1 purpose we exclude from each DEM the non-stable terrain such as glaciers and in general all
2 off-glacier area at elevations higher than 5400 m a.s.l. (which is the estimated equilibrium line
3 altitude (ELA) in the Upper Langtang catchment (Ragettli et al., 2015)). The correlation score
4 maps, indicating which pixels have been matched successfully during the DEM extraction
5 process, are used to exclude all DEM grid cells with a correlation score below 0.5. Then,
6 horizontal shifts are determined by minimizing the aspect-dependent bias of elevation
7 differences (Nuth and Kääb, 2011) between each DEM pair. Because of the slope dependency
8 of the method all terrain below a slope of 10° is excluded. The ‘older’ DEM is then resampled
9 (bilinear interpolation) according to the determined horizontal shift. In a second step the
10 vertical DEM shifts and possible tilts are corrected using second order trend surfaces fitted to
11 all gently inclined ($\leq 15^\circ$) stable terrain (Bolch et al., 2008; Pieczonka et al., 2011; Pieczonka
12 and Bolch, 2015).

13 We resample all DEMs bilinearly to the grid size of the coarsest DEM (30 m) to reduce the
14 effect of different resolutions. Elevation differences are calculated by subtracting the older
15 from the younger DEM (such that glacier thickening values are positive) and are converted to
16 elevation change rates by dividing by the number of ablation seasons between the acquisition
17 dates. Seasonal effects on elevation change rates are neglected when discussing time intervals
18 between DEMs of 4 years or longer, since elevation changes during the winter half-year are
19 usually minor. On average, less than 20% of annual precipitation occur during post-monsoon
20 and winter (Immerzeel et al., 2014b), and less than 3% of annual glacier ice-melt (Ragettli et
21 al., 2015). Area-average glacier elevation change rates are calculated using always the
22 maximum glacier extent between two acquisition dates.

23 **3.2.3 Processing of elevation change maps**

24 Processing of the elevation change ($\Delta h/\Delta t$) maps involves two main steps: i) removal of pixel
25 values identified as outliers and ii) filling of gaps.

26 Outlier removal

27 The stereo matching score maps provided by PCI Geomatica are used to identify elevation
28 data that can be considered for elevation change calculations. If the correlation score of a
29 given DEM pixel is below 0.5, this indicates a poor matching score (Pieczonka et al. 2011)
30 and therefore the corresponding $\Delta h/\Delta t$ values are treated as ‘no data’. Very unrealistic
31 elevation change data (exceeding ± 150 m) are also excluded from the analysis.

1 We use the standard deviation (σ) of observed elevation changes to identify $\Delta h/\Delta t$ outliers.
2 Outliers are defined separately for debris-covered glacier areas and debris-free glacier areas.
3 For the latter we additionally distinguish between glacier area below and above the ELA
4 (estimated at 5400 m a.s.l., see above). σ -levels are thus calculated for each of the three area
5 types in every $\Delta h/\Delta t$ map. Below the ELA (both debris-free and debris-covered area), pixels
6 are defined as outliers if $\Delta h/\Delta t$ values differ from the average by $>3\sigma$ (e.g. Gardelle et al.,
7 2013). This means that only very few data are classified as outliers, since three standard
8 deviations account for 99.7% of the sample (assuming the distribution is normal). The
9 conservative outlier definitions are justified by the shallow slopes and high contrast, which
10 also explains why stereo matching scores are generally higher below the ELA (Figure 2c).
11 Above the ELA, steep terrain or featureless snow surfaces lead to low DEM accuracy and
12 therefore the outlier criteria should be more restrictive (e.g. Pieczonka et al., 2013; Pieczonka
13 and Bolch, 2015). On debris-free glacier area above the ELA, pixels are therefore defined as
14 outliers if $\Delta h/\Delta t$ values differ from the average by $>1\sigma$ (which applies to approximately 32%
15 of the values if the distribution is normal). A stricter criterion for the accumulation area is also
16 justified by the fact that it can be assumed that elevation changes in the accumulation areas
17 over periods of several years are on average small (Schwitter and Raymond, 1993; Huss et al.,
18 2010). Because we use different σ thresholds above and below the ELA we test the sensitivity
19 of calculated glacier volume changes to a ± 100 m ELA uncertainty. Furthermore, we test the
20 sensitivity to different outlier definitions by comparing our results to the results obtained with
21 a 2σ -level applied to all area types.

22 Gap filling

23 On the glacier areas below the ELA, with only very few data gaps, missing data are replaced
24 using inverse distance weighting (IDW). In the accumulation areas, on the other hand, data
25 gaps can extend over a wide elevation range if the terrain is steep or if the gaps are very large.
26 Because of the elevation dependency of $\Delta h/\Delta t$ values (e.g. Huss et al., 2010) only values from
27 the same altitudinal range should be used to fill data gaps. We thus replace missing data in the
28 accumulation areas by median $\Delta h/\Delta t$ values per 50-m elevation band considering all available
29 data for a given glacier (also from $\Delta h/\Delta t$ maps representative of different periods). For this,
30 we first calculate the mean elevation change rates per 50-m elevation band of each glacier and
31 every $\Delta h/\Delta t$ map and then determine the median of the ensemble. $\Delta h/\Delta t$ maps that are

1 rejected from the ensemble (see Section 3.2.5 below) and in general all values representative
2 of short periods ($\Delta t < 4$ years) are not considered to calculate the ensemble-median values.

3 **3.2.4 Uncertainty**

4 Elevation change uncertainty estimates are based on the standard error $E_{\Delta h}$ calculated per
5 elevation band (Gardelle et al., 2013). The standard error quantifies the effect of random
6 errors on uncertainty according to the standard principles of error propagation:

$$7 \quad E_{\Delta h} = \frac{\sigma_{\Delta h, noglac}}{\sqrt{N_{eff}}} \quad (1)$$

$$8 \quad N_{eff} = \frac{N_{tot} \times PS}{2d} \quad (2)$$

9 $\sigma_{\Delta h, noglac}$ is the standard deviation of the mean elevation change of non-glacierized terrain per
10 elevation band, N_{eff} is the effective and N_{tot} the total number of observations. PS is the pixel
11 size (30 m) and d is the distance of spatial autocorrelation. d is equal to the range of the
12 spherical semivariogram obtained by least squares fit to the experimental, isotropic variogram
13 of all off-glacier elevation differences (Wang and Kääb, 2015; Magnússon et al., 2016). The
14 distance of spatial autocorrelation of the elevation change maps varies between 260 m and
15 730 m with an average of 495 m.

16 To quantify the elevation change uncertainty of glacier area spanning several elevation bands,
17 weighted averages of $E_{\Delta h}$ are calculated. $E_{\Delta h}$ of each individual elevation band is weighted by
18 the glacier hypsometry. Elevation change uncertainties therefore vary for each individual
19 glacier because of the different glacier area-elevation distributions. $E_{\Delta h}$ tends to increase with
20 altitude (Figure 3, Figure 4) due to steeper slopes, snow and deep shadows, which are factors
21 that decrease the accuracy of DEMs derived from stereo data (e.g. Nuimura et al., 2011).
22 Uncertainty estimates for each individual glacier therefore account for the spatially non-
23 uniform distribution of uncertainty. Elevation change uncertainties of glaciers with a high
24 accumulation area such as Kimoshung and Lirung Glaciers (Table 1) are 50%-100% higher
25 than those of other glaciers, in accordance with lower DEM matching scores (Figure 2). The
26 low uncertainty associated to debris-covered areas agrees with the 30%-100% lower off-
27 glacier errors on shallow slopes ($s < 18^\circ$, 95th percentile of debris-covered glacier slopes) than
28 on steeper slopes ($s < 45^\circ$, 95th percentile of glacier slopes; Figure S1).

1 The standard error can be interpreted as the 68% confidence interval of the sample mean if the
2 distribution is normal. Since we are conservatively assuming no error compensation across
3 elevation bands the approximate confidence level in our uncertainty estimates per glacier is
4 higher than 68%.

5 This study aims at obtaining an ensemble of results about elevation change rates from the set
6 of seven DEMs available for the period 2006-2015 and we thus calculate an ensemble
7 uncertainty. The uncertainty in a sample mean is different from the uncertainty in individual
8 observations about recent volume change rates. To identify the range of ensemble values
9 (hereafter ‘ensemble uncertainty’) we use the standard deviation of the ensemble values
10 multiplied by 1.96. By multiplication with 1.96 we obtain 95% confidence levels, assuming
11 normal distribution.

12 For overall mass budget uncertainties we assume an ice density of 850 kg/m^3 to convert the
13 volume change into mass balance (Sapiano et al., 1998; Huss, 2013) and consider the
14 elevation change rate uncertainties and an ice density uncertainty of 60 kg/m^3 .

15 **3.2.5 Ensemble selection**

16 The possible two-fold combinations of all available DEMs are classified in two groups: maps
17 that involve the Hexagon 1974 DEM and maps that represent only 21st century elevation
18 changes (2006-2015). From the first group we only use the 1974-2006 $\Delta h/\Delta t$ map, to strictly
19 separate our two main study periods 1974-2006 and 2006-2015. The redundancy of
20 information between 2006 and 2015 allows extracting 12 maps of $\Delta h/\Delta t$ with a time interval
21 larger than 4 years. Since $\Delta h/\Delta t$ uncertainties increase with shorter time intervals between
22 DEMs (Figure 5, Table 3) $\Delta h/\Delta t$ maps with $\Delta t < 4$ years are not considered for the 2006-2015
23 ensemble. After careful evaluation of the DEMs in terms of $\Delta h/\Delta t$ uncertainties and the 2015
24 Nepal earthquake impact we discard from the 2006-2015 ensemble also all $\Delta h/\Delta t$ maps
25 involving the ALOS PRISM 2010 and the SPOT7 May 2015 scenes. The 2006-2015
26 ensemble consists therefore of six maps of $\Delta h/\Delta t$ (Figure S2).

27 The ALOS PRISM DEM is discarded because uncertainties associated to $\Delta h/\Delta t$ maps
28 involving this DEM are 30-100% higher than if other DEMs are involved (Table 3). The
29 ALOS-PRISM sensor has a radiometric resolution of 8-bit, which means that in comparison
30 to a 12-bit image (SPOT6/7, Table 2), $2^4=16$ times less information is provided per

1 panchromatic image pixel. The image contrast is therefore lower, which decreases the
2 accuracy of this DEM.

3 The May 2015 DEM is not considered for the 2006-2015 ensemble because of massive
4 deposits of avalanched snow and ice as a consequence of the April 2015 Nepal earthquake,
5 which strongly limits the representativeness of this DEM for the 2006-2015 period. The
6 October 2015 SPOT7 DEM is still considered for the 2006-2015 ensemble because ~~until~~ six
7 months after the earthquake most avalanche material disappeared from the glacier (section
8 4.6).

9 Due to the incomplete representation of Langtang Glacier on the SPOT6 Apr 2014 scene (the
10 scene does not cover the area north of 28°19'N, Figure S2a and d), $\Delta h/\Delta t$ maps involving this
11 DEM are excluded when discussing ensemble results for Langtang Glacier.

12 **3.3 Delineation of glaciers, debris-covered areas, and supraglacial cliffs/lakes**

13 The glacier outlines were manually delineated. We used the orthorectified satellite images
14 with the least snow cover (the Cartosat-1 2006 and 2009 scenes) to delineate the accumulation
15 areas, and assumed no changes in the accumulation area over time. The tongues of the seven
16 studied glaciers and debris extents were re-delineated for every year for which satellite images
17 are available (1974, 2006, 2009, 2010, 2014 and 2015), using the corresponding orthorectified
18 satellite images. A first operator delineated the outlines and a second operator provided
19 feedback in order to improve delineation accuracy. To quantify the uncertainty in derived
20 glacier area changes we consider a 0.5 pixel size delineation uncertainty (Paul et al., 2013).

21 The four largest glaciers in the valley were already delineated manually by Pellicciotti et al.
22 (2015) for the years 1974 and 2000. However, we decided not to use those outlines because of
23 the considerably higher resolution of the images that are available for this study and for
24 consistency in the procedure applied for different outlines. We also re-delineated the
25 catchment boundaries using the SRTM 30 m DEM and flow accumulation to accurately
26 identify the ice divides between neighboring catchments. As a result, the calculated glacier
27 areas (Table 1) changed considerably with respect to Pellicciotti et al. (2015). The 1974
28 glacier area of Langshisha Glacier changed by -40.4% (Figure S3), mostly due to clipping
29 with the catchment mask which reduced the extent of the accumulation areas. The 1974 areas
30 of Langtang, Shalbachum and Lirung changed by -8.7%, -9.5% and +8.0%, respectively.

1 To identify glacier area associated to small glaciers in the catchment that are not discussed in
2 this study we used the glacier outlines provided by the GAMDAM glacier inventory
3 (Nuimura et al., 2015). Those areas were masked out from off-glacier terrain for the co-
4 registration of the DEMs and stable terrain accuracy assessments.

5 Six quality checked maps of supraglacial cliffs and lakes are used to characterize debris-
6 covered glacier surfaces (Steiner et al., 2016). The cliff and lake inventories were generated
7 based on the available satellite imagery for the period 2006-2015 (Oct 2006, Nov 2009, Dec
8 2010, Apr 2014, May 2015 and Oct 2015). As for the glacier outlines, cliff and lake outlines
9 have been delineated by two independent operators. To further improve the accuracy of the
10 inventories, a third operator used slope and elevation change maps to identify potential cliff
11 and lake locations. The first two operators then used these indications to review the
12 inventories. All outlines have been obtained by manual delineation on the basis of the
13 orthorectified satellite images.

14 We calculated the fraction of pixels including lakes and cliffs per 50 m elevation band of each
15 debris-covered tongue (excluding tributary branches, Figure 6). In the following, we only
16 discuss median 2006-2015 cliff and lake area fractions to minimize seasonal effects. Large
17 avalanche cones, such as those present on Lirung and Langtang Glacier after the April 2015
18 earthquake, are masked out from the inventories before calculating median values.

19 **3.4 Surface velocities**

20 To assist with the interpretation of volumetric changes, we use glacier velocities determined
21 with the COSI-Corr cross-correlation feature-tracking algorithm (Leprince et al., 2007) and
22 the available satellite imagery. The orthorectified Cartosat-1 Nov 2009 and ALOS-PRISM
23 Dec 2010 images were used for this purpose. Other image pairs were not considered due to
24 longer periods between acquisitions (leading to image decorrelation) or the presence of snow
25 patches at lower elevations (SPOT6 April 2014, SPOT7 May 2015). The selected
26 orthorectified images (5 m resolution) were adjusted according to the shifts determined by co-
27 registration (Section 3.2.2). Since the window size must be large enough to avoid correlating
28 only noise but small enough not to degrade the output resolution (Dehecq et al., 2015), we
29 tested several configurations. The best results for the COSI-Corr correlation analysis were
30 achieved using multiscale window sizes of 128 down to 32 pixels, as also proposed by
31 Scherler et al. (2008). To post-process the velocity data we removed pixels with x- or y-

1 velocity values greater than 40 m/a, since these were identified as errors by manually
2 measuring the surface displacement on the basis of the orthorectified images and prominent
3 features. We then ran a median filter on the data to remove areas which show a local reversal
4 in x or y directions. Missing values were then filled with the mean of the adjacent 8 values.
5 Finally, the velocity map was resampled to 30 m resolution with a bicubic algorithm.

6 **3.5 Assessment of the April 2015 earthquake impact**

7 We quantify the impact of the avalanche events after the April 2015 earthquake on volume
8 changes of debris-covered tongues. For this purpose we use the April 2014 - May 2015 Δh
9 map to quantify the accumulated volumes less than two weeks after the earthquake, and the
10 April 2014 - Oct 2015 Δh map to quantify the remaining volumes after one ablation season.
11 To identify glacier area where avalanche material accumulated we consider all glacier grid
12 cells with significant positive elevation changes ($\Delta h > 5$ m). Approximately 7.9% (1.9 km²) of
13 all debris-covered areas were affected by avalanches according to this definition. To account
14 for pre-earthquake volume losses we first subtract from the DEM differences the elevation
15 changes between April 2014 and April 2015 determined on the basis of the Oct 2006 - Feb
16 2015 mean thinning rates. Note that we do not use the Feb 2015 - May 2015 and the Feb 2015
17 - Oct 2015 Δh maps to quantify avalanche debris volumes because the calculated uncertainties
18 associated to these maps are up to 300% higher than the uncertainties associated to
19 differential DEMs involving the Apr 2014 scene (Table S1).

20

21 **4 Results**

22 **4.1 Mean glacier surface elevation changes**

23 The 2006-2015 ensemble consistently indicates an increase in mean glacier thinning rates in
24 comparison to the period 1974-2006 (Figure 7h). For 2006-2015 we calculate an ensemble-
25 mean thinning rate of -0.45 ± 0.18 m a⁻¹, while for the period 1974-2006 we identify a
26 thinning rate of -0.24 ± 0.08 m a⁻¹ (Table 4). This corresponds to an increase in determined
27 mean thinning rates by 0.21 m a⁻¹ or 87.5%. The error bounds associated to the two periods
28 are overlapping at the extremes. However, error bounds are not overlapping at 80%
29 confidence levels (assuming normal distribution). Given the probability of less than 10% for

1 1974-2006 and 2006-2015 thinning rates for being above or below this confidence interval,
2 the estimated confidence level of accelerated thinning rates is higher than 99%.

3 From the seven studied glaciers in the valley, the thinning rates of Langtang, Langshisha and
4 Yala Glaciers have accelerated at 99% confidence levels (Figure 7, Table 4). At Shalbachum
5 Glacier the error bounds are overlapping but the estimated probability of thinning acceleration
6 is more than 90%. At Lirung and Kimoshung Glaciers the mean elevation change rates have
7 likely remained approximately constant (Table 4). Mean thinning rates of these glaciers
8 increased by less than 0.10 m a^{-1} between 1974-2006 and 2006-2015. Also at Ghanna Glacier
9 the 1974-2006 value and the 2006-2015 ensemble mean differ by only 0.05 m a^{-1} (Table 4).
10 However, the scatter in the 2006-2015 values is such that no trend can be identified. Ghanna
11 Glacier is the only glacier where the ensemble of values available for the period 2006-2015
12 did not narrow down the uncertainty associated to individual periods (Figure 7).

13 The most negative elevation change for 1974-2006 was observed at Shalbachum (-0.43 ± 0.08
14 m a^{-1} , Table 4) and Ghanna Glacier ($-0.51 \pm 0.05 \text{ m a}^{-1}$). The least negative values were
15 calculated for Langshisha ($-0.12 \pm 0.09 \text{ m a}^{-1}$) and Kimoshung Glaciers ($0.06 \pm 0.13 \text{ m a}^{-1}$).
16 Comparing the period 1974-2006 and the 2006-2015 ensemble mean values, the strongest
17 thinning acceleration took place at Yala Glacier (from $-0.33 \pm 0.06 \text{ m a}^{-1}$ to $-0.89 \pm 0.23 \text{ m a}^{-1}$,
18 Table 4). Yala Glacier was also the glacier with the highest 2006-2015 ensemble mean
19 thinning rate.

20 Elevation change rates are also calculated separately for the five debris-covered tongues
21 (Figure 8, Table 4). An increase in ~~volume loss~~ thinning rates is evident on the Langtang,
22 Langshisha, Shalbachum and Lirung tongues. Thinning rates increased between 15%
23 (Langtang tongue) and 68% (Langshisha and Shalbachum tongues). Changes in thinning rates
24 are not significant for Ghanna tongue, but five out of six members of the 2006-2015 ensemble
25 decreased as compared to the previous period.

26 Of all debris-covered areas, the downwasting rates on Lirung tongue are the highest. This
27 applies to both the period 1974-2006 ($-1.03 \pm 0.05 \text{ m a}^{-1}$, Table 4) and to the 2006-2015
28 ensemble mean ($-1.67 \pm 0.59 \text{ m a}^{-1}$, Table 4). The 2006-2015 ensemble uncertainty is very
29 large on Lirung tongue ($\pm 0.59 \text{ m a}^{-1}$), which we believe is due to systematic errors in the
30 2009-2014 differential DEM that represents an outlier in the ensemble (Figure 8). However,
31 neither on Lirung nor on Langtang tongue (the two glaciers most affected by post-earthquake

1 avalanches, see section 4.6) post-earthquake elevation changes (2006-Oct 2015 or 2009-Oct
2 2015) represent outliers with respect to other 2006-2015 multi-annual periods. The lowest
3 ~~volume loss~~ thinning rates are identified for Ghanna tongue (Figure 8, Table 4). Here, the
4 2006-2015 ensemble mean value ($-0.50 \pm 0.20 \text{ m a}^{-1}$) is about 30% of the thinning rate at
5 Lirung tongue.

6 **4.2 Sensitivity to outlier correction and ELA definitions**

7 Mean elevation change values are most sensitive to outlier definitions for Langshisha Glacier
8 1974-2006 (Table 5). If a 2σ -level is used to define outliers for all area types (instead of a 3σ -
9 level above and a 1σ -level below the ELA, Section 3.2.3), $\Delta h/\Delta t_{1974-2006}$ for Langshisha
10 Glacier changes by -0.09 m a^{-1} from $-0.12 \pm 0.09 \text{ m a}^{-1}$ to $-0.21 \pm 0.09 \text{ m a}^{-1}$. If we compare
11 the results obtained with an estimated ELA at 5300 m a.s.l. to the results obtained with an
12 ELA at 5500 m a.s.l., mean elevation changes of individual glaciers differ by up to -0.23 m a^{-1}
13 (Shalbachum Glacier 1974-2006). However, only for two glaciers the sensitivity values
14 exceed the uncertainty values estimated from off-glacier elevation change errors (at
15 Shalbachum and Yala Glacier 1974-2006, Table 5). In both cases the differences can be
16 explained by unrealistic patterns (strongly negative elevation changes above 5400 m a.s.l.),
17 that are not identified as outliers with a 3σ threshold applied to areas below 5500 m a.s.l. Our
18 analysis thus shows that elevation change estimates are in most cases not significantly
19 influenced by outlier definitions and ELA estimates.

20 **4.3 Altitudinal distribution of elevation changes**

21 The altitudinal distribution of mean elevation changes clearly show that the thinning patterns
22 of all debris-covered tongues have changed over time (Figure 9, Figure 10). Areas with clear
23 increases in thinning rates can be identified for Langtang Glacier 5000-5150 m a.s.l.
24 (25%-100% thinning rate increase), for Langshisha Glacier 4650-5100 m a.s.l. (25%-260%),
25 for Shalbachum Glacier 4500-4800 m a.s.l. (25%-180%) and for Lirung Glacier 4300-4350
26 m a.s.l. (80%-170%). Thinning rates have remained mostly constant in the lower third of the
27 elevation ranges of the tongues (Langtang, Shalbachum and Lirung Glaciers). At Ghanna
28 Glacier, thinning rates have recently declined near the glacier terminus at 4800-4850 m a.s.l.
29 (60-90% thinning rate decrease, Figure 9e). This pattern contrasts with all other temporal
30 patterns for debris-covered glacier areas.

1 On Langshisha Glacier thinning patterns in 1974-2006 and 2006-2015 are different near the
2 terminus (Figure 9b). Here, the glacier tongue became very narrow in the last decade and
3 ultimately a small part below 4500 m a.s.l. disconnected from the main tongue (Figure 1)
4 between 2010 and 2014. The fragmentation of the tongue lead to mean thinning rates close to
5 zero at elevation bands where a substantial part of the glacier area disappeared.

6 Overall, the thinning profiles of 2006-2015 ensemble members show very similar
7 characteristics (Figure 9, Figure 10). The profiles diverge for the uppermost elevation bands
8 of the tongues and in the accumulation areas. This agrees with the larger error that is
9 attributed to higher elevations (Figure 3). Above 5500 m a.s.l. it is impossible to separate
10 uncertainty from actual differences in thinning rates.

11 On Yala Glacier there has been a three-fold increase in thinning rates below 5400 m a.s.l.,
12 comparing 1974-2006 to the 2006-2015 ensemble results (Figure 10d). Maximal thinning
13 takes place at the terminus and then decreases nearly linearly with altitude until it reaches
14 values close to zero. This is in clear contrast to the much less uniform patterns on debris-
15 covered glaciers (Figure 10a-c). On debris-covered glaciers, the elevation corresponding to
16 the maximum thinning rates is different from glacier to glacier. On Shalbachum and Lirung
17 Glaciers the maximum is reached somewhere close to the upper end of the tongue (4650-4750
18 m a.s.l. and 4300-4400 m a.s.l., respectively, Figure 9c and d), on Langtang and Ghanna
19 Glaciers more in the middle part (4950 – 5150 m a.s.l. and 4900-5000 m a.s.l., respectively,
20 Figure 9a and e) and on Langshisha Glacier closer to the terminus (4450-4700 m a.s.l., Figure
21 9b). On the large debris-covered glaciers, areas of maximum thinning seem to have shifted
22 and extended to higher elevations only at Langtang Glacier, where during the period 1974-
23 2006 maximum thinning occurred between 4850 and 4950 m a.s.l. (Figure 9a). On
24 Shalbachum Glacier maximum thinning during the period 1974-2006 occurred slightly higher
25 up at 4750 – 4800 m a.s.l. (Figure 9c).

26 Note that the altitudinal $\Delta h/\Delta t$ profiles (Figure 9, Figure 10) always refer to the same position
27 in space, since 50 m elevation bands were delimited only once on the basis of the SRTM
28 1 Arc-Second Global DEM. To account for the up-valley movement of on-glacier elevation
29 bands over time due to surface lowering, profiles would have to be slightly shifted relative to
30 each other. However, given the maximum thinning rates of 1-1.5 ma^{-1} in 1974-2006, the
31 maximum relative adjustment of values in Figure 9 and Figure 10 would never exceed one

1 50 m elevation band. Accounting for the shifting of elevation bands over time would therefore
2 not lead to different conclusions regarding changes in spatial $\Delta h/\Delta t$ patterns.

3 **4.4 Glacier area changes**

4 Debris-free Yala Glacier experienced the strongest increase in relative annual area loss of all
5 studied glaciers (1974-2006: $-0.43 \pm 0.05\% \text{ a}^{-1}$, 2006-2015: $-1.77 \pm 0.16\% \text{ a}^{-1}$, Table 6).
6 | During the same two time intervals debris-free Kimoshung Glacier shrank only at rates of
7 $0.08 \pm 0.01\% \text{ a}^{-1}$ and $0.05 \pm 0.02\% \text{ a}^{-1}$, respectively. This represents significantly lower retreat
8 rates for the second period than at Yala Glacier. The differences in area change rates are
9 consistent with the identified differences in mean glacier surface elevation changes, where the
10 two glaciers also represent opposite extremes (Section 4.1).

11 In comparison to the current retreat rates of Yala Glacier, all debris-covered glaciers are
12 shrinking at a much slower pace, with retreat rates between $-0.04 \pm 0.04\% \text{ a}^{-1}$ and $-0.40 \pm$
13 $0.12\% \text{ a}^{-1}$ (Table 6). Also debris-covered glaciers for which we observe high annual volume
14 losses have nearly stationary fronts (e.g. Shalbachum Glacier: 2006-2015 thinning rate $-0.53 \pm$
15 0.19 m a^{-1} , 2006-2015 area loss $-0.04 \pm 0.04\% \text{ a}^{-1}$). Ghanna Glacier in contrast shows a
16 slightly more significant retreat ($-0.40 \pm 0.12\% \text{ a}^{-1}$, Table 6), although the mean thinning rates
17 are the least negative of all debris-covered areas (Figure 8).

18 **4.5 Surface velocities and supraglacial cliff/lake areas**

19 Approximately 10% of all grid cells for the three largest debris-covered tongues (Langtang,
20 Langshisha, Shalbachum) contain supraglacial cliff features ('Cliff Area' in Table 7). At
21 Lirung and Ghanna tongues this value decreases to 8% and 3%, respectively. For Ghanna
22 tongue practically no supraglacial lakes could be identified, while at the other debris-covered
23 tongues 'Lake Area' is between 2.3% and 3.3%.

24 The mean surface velocities of the tongues range between 1.6 m a^{-1} (Ghanna tongue) and
25 7 m a^{-1} (Langhsisha tongue). The mean and the standard deviation of off-glacier surface
26 velocities are 1.3 m a^{-1} and 1.9 m a^{-1} , respectively. At Ghanna and Lirung tongue, which both
27 have a mean surface velocity below 3 m a^{-1} , it is therefore practically impossible to
28 discriminate moving ice from quasi-stagnant ice. Following Scherler et al. (2011b), all glacier
29 grid cells with a surface velocity of less than 2.5 m a^{-1} are therefore termed 'stagnant' for

1 simplicity. According to this definition, the tongue area classified as ‘stagnant’ (Table 7)
2 ranges from 20% (Langshisha tongue) to 85% (Ghanna tongue).

3 In our sample of five debris-covered glaciers, cliffs and lakes seem to appear more frequently
4 on glaciers which are dynamically active. We identify a highly significant negative correlation
5 (Pearson's linear correlation coefficient $r=-0.99$) between cliff area fraction per tongue and the
6 percentage of stagnant tongue area. ‘Lake Area’ and ‘% stagnant area’ are also negatively
7 correlated ($r=-0.87$). At the scale of individual tongues, a correlation between surface
8 | velocities and cliff appearance is evident at Shalbachum Glacier (Figure 11c). Here, were we
9 identify a correlation of 0.85 (respectively 0.68) between the altitudinal velocity profile and
10 cliff (respectively lake) areas per 50 m elevation band. Also on the two other large debris-
11 covered tongues in the valley, on Langtang and Langshisha tongues, cliff appearance clearly
12 | decreases towards the termini where the glaciers are quasi-stagnant. However, hereon these
13 two glaciers the surface velocities and cliff appearance are not linearly correlated since the
14 highest cliff area densities are identified 200-300 m below the altitude ranges corresponding
15 to maximum surface velocity.

16 To investigate a possible link between accelerated thinning and the presence of supraglacial
17 lakes and cliffs we compare ‘Cliff Area’ and ‘Lake Area’ (as provided in Table 7) to changes
18 in mean thinning rates per tongue ($\Delta \Delta h/\Delta t$, difference between ‘1974-2006’ and ‘ensemble
19 mean 2006-2015’ as provided in Table 4). Overall, the correlation coefficient between
20 fractional cliff area per tongue and $\Delta \Delta h/\Delta t$ is -0.62 (and -0.50 between lake area and
21 $\Delta \Delta h/\Delta t$). The likely reduced thinning rates on Ghanna tongue (Figure 8e) indeed correspond
22 to low cliff and lake area fractions (3.2% and 0.4%, respectively). On Lirung, Shalbachum
23 and Langshisha tongues thinning accelerated by 0.47-0.64 m a^{-1} , whereas fractional cliff and
24 lake areas are similar (cliff area: 8.0-10.5%, lake area: 2.3-2.6%). Also Langtang tongue is
25 characterized by relatively high cliff and lake area fractions (10% and 3.3%, respectively,
26 Table 7) but the identified changes in thinning rates are only minor. The acceleration of mean
27 thinning rates at Langtang tongue is significant at the 95% confidence level (Figure 8a), but
28 the difference in mean thinning rates 1974-2006 and 2006-2015 is only -0.12 m a^{-1} (Table 4).

29 At locations where thinning rates did not increase significantly we mostly identify low cliff
30 area fractions below 10% (e.g. on Langtang tongue below 4750 m a.s.l. and above 5150 m
31 a.s.l., at Shalbachum below 5500 m a.s.l. and at Ghanna tongue). Conversely, cliff area

1 fractions are generally higher than 10% where the 2006-2015 ensemble consistently indicates
2 thinning acceleration (Figure 11). Exception to this observation are the high cliff area
3 fractions at Langtang Glacier 4750-4900 m a.s.l., where thinning rates did not change
4 significantly (Figure 11a), and low cliff area fractions at Shalbachum Glacier 4750-4800 m
5 a.s.l., where thinning rates increased (Figure 11c). Lirung tongue also shows a different
6 behavior, except at the lowest elevation band. However, maximum thinning acceleration at
7 4300 m a.s.l. corresponds to a relatively high lake area fraction of 6% (Figure 11d).

8 Altitude bands with no significant increases in thinning rates on Langtang Glacier consistently
9 coincides with relatively low surface velocities below 5 m a^{-1} . At Langhisha and Shalbachum
10 tongues this is also the case (Figure 11). Across all debris-covered glacier tongues, 77% of all
11 elevation bands where thinning accelerated ($\Delta \Delta h/\Delta t < -0.2 \text{ m a}^{-1}$) are not stagnating
12 (velocities above 2.5 m a^{-1}), and in 72% of all elevation bands where thinning rates remained
13 constant or declined ($\Delta \Delta h/\Delta t \geq -0.2 \text{ m a}^{-1}$) we observe velocities below 2.5 m a^{-1} .

14 **4.6 Impacts of the April 2015 earthquake**

15 We calculate a total volume of post-earthquake avalanche debris in May 2015 of $2.5 \cdot 10^7 \text{ m}^3$,
16 which is equivalent to a cube length of 292 m. 40% of the avalanche material remained until 6
17 Oct 2015 (Table 8). The two glaciers which were most affected by avalanches were Langtang
18 Glacier (receiving 58% of the total volume) and Lirung Glacier (29%). The avalanche cone at
19 Lirung Glacier piled up to a height of nearly 60 m, while the avalanche material at Langtang
20 Glacier was more spread (Figure 12). Consequently, more material remained until 6 Oct 2015
21 at Lirung Glacier (57%), while at Langtang Glacier 31% remained (Table 8). Field visits at
22 the end of October 2015 to Lirung and Langtang Glaciers revealed that a smooth debris layer
23 melted out of the avalanche material and covered the surface uniformly with a thickness of a
24 few centimeters (P. Buri and P. Egli, personal communication).

25 Volume loss from glacier area where avalanche material accumulated between May and
26 October 2015 was 30 times higher than during an average ablation season (May 2015 – Oct
27 2015: $-1.5 \cdot 10^7 \text{ m}^3$, Oct 2006 – Apr 2014: $-5 \cdot 10^5 \text{ m}^3 \text{ a}^{-1}$). After this rapid initial downwasting
28 the avalanche deposits diminished to a volume of 10^7 m^3 , equivalent to an average positive
29 surface elevation change over all debris-covered glacier area of $0.52 \pm 0.19 \text{ m}$ (Table 8, or
30 0.06—0.09 m a⁻¹ if divided over six to nine years, which is the shortest time interval of any
31 $\Delta h/\Delta t$ map in the ensemble involving the October 2015 DEM). The avalanche impact on the

1 Oct 2015 DEM is thus within the uncertainty range associated to multi-annual $\Delta h/\Delta t$ values
2 ($\pm 0.12 \text{ m a}^{-1}$, Table 3) and justifies why the October 2015 DEM is considered for the 2006-
3 2015 ensemble.

4 The avalanche traces are still visible six months after the earthquake at Lirung Glacier (4350-
5 4400 m a.s.l.), at Langtang Glacier (4500-4900 m a.s.l.), at Langshisha Glacier (4800 m a.s.l.)
6 and at Shalbachum Glacier (4750 m a.s.l.) (Figure 9). Except for Lirung Glacier at 4350
7 m a.s.l. the 2006-Oct 2015 and 2009-Oct 2015 thinning profiles are within the error bounds
8 associated to other multi-annual periods.

9

10 **5 Discussion**

11 **5.1 Elevation changes of debris-covered glaciers**

12 Elevation changes in the debris-covered area are primarily independent of elevation (Figure
13 9), as previously identified in the Langtang catchment (Pellicciotti et al., 2015) and elsewhere
14 in high-mountain Asia (e.g. Bolch et al., 2011; Dobhal et al., 2013; Pieczonka et al., 2013;
15 Pieczonka and Bolch, 2015; Ye et al., 2015). Such patterns have usually been explained by
16 downglacier increase of debris thickness and by ablation associated with supraglacial lakes
17 and exposed ice cliffs. Our analysis shows that, with few exceptions, the highest thinning
18 rates and the strongest increase in thinning rates can be associated to areas with a high
19 concentration of ice cliffs and supraglacial ponds (Figure 11, Figure S4). While previous
20 studies have pointed out that debris-covered areas with a large presence of supraglacial cliffs
21 and lakes make a disproportionately large contribution to ablation (Reid and Brock, 2014;
22 Buri et al., 2016; Miles et al., 2016a; Thompson et al., 2016), this is the first study which
23 documents the relation between accelerations in ~~volume loss~~ thinning rates and the large
24 presence of supraglacial cliffs and lakes.

25 Supraglacial cliffs seem to appear more frequently on slowly moving ice ($5\text{-}10 \text{ m a}^{-1}$, Figure
26 11) and not where the glacier is stagnant (Sakai et al., 2002; Bolch et al., 2008; Thompson et
27 al., 2016). This can be explained by compressive stresses associated with flow deceleration
28 that may initiate fracturing (Benn et al., 2009). Such stresses are usually not large enough to
29 initiate open surface crevasses, but in combination with elevated water pressure due to local
30 water inputs lead to hydrologically driven fracture propagation (hydrofracturing) and

1 englacial conduit formation (Benn et al., 2009). The collapse of large englacial voids
2 destabilizes the debris layers and leads to the formation of new ice cliffs. Accordingly,
3 accelerated thinning of debris-covered area in the Upper Langtang catchment does not take
4 place on stagnating parts of the tongues, but where the transition between the active and the
5 stagnant ice can be expected (Figure 11).

6 The appearance of supraglacial lakes, on the other hand, is strongly related to the surface
7 gradient (Sakai and Fujita, 2010; Miles et al., 2016b). Large supraglacial lakes can only form
8 where the slope is less than 2° (Reynolds, 2000) and where local water input is high. These
9 conditions are not met on debris-covered glacier sections in the Upper Langtang catchment,
10 since local surface slope is consistently above 5° (Pellicciotti et al., 2015). It is interesting to
11 note that the highest lake area fractions (Lake Area $> 6\%$) are found on avalanche deposition
12 zones at Langtang Glacier (4750-4800 m a.s.l., Figure 7a and Figure 11a) and at Lirung
13 Glacier (4300 m a.s.l., Figure 7d and Figure 11d). This is likely related to high local surface
14 water inputs from melting of avalanche snow and ice. On Langtang Glacier frequent
15 avalanche inputs may explain why thinning did not accelerate at the altitude range between
16 4750 m a.s.l. and 4900 m a.s.l., in spite of the presence of exposed ice (Cliff Area $> 13\%$,
17 Figure 11a).

18 Several studies show that lakes and cliffs are important but cannot explain the mass loss alone
19 (e.g. Sakai et al., 2002; Juen et al., 2014). The high thinning magnitudes on the upper sections
20 of Shalbachum tongue (4750-4800 m a.s.l.) likely cannot be attributed to lakes and cliffs
21 (cliff/lake area fractions are below 5%, Figure S4c), and thin layers of deposited debris in the
22 upper sections of the glacier tongue could explain such patterns.

23 Reduced ice fluxes also contribute to thinning accelerations. To assess how much this factor
24 contributes to the observed accelerations in thinning it would be necessary to quantify
25 changes in ice flux over time (e.g. Nuimura et al., 2011; Berthier and Vincent, 2012; Nuth et
26 al., 2012). However, information about the evolution of surface velocities over long time
27 periods would be required, which our dataset cannot provide. Over the timescales considered
28 in this study, on the other hand, high warming rates have been identified in this part of the
29 Himalaya (Shrestha et al., 1999; Lau et al., 2010). The rise in air temperatures directly
30 impacts glacier melt rates, and can explain rapid acceleration of thinning where ice is not
31 insulated from warming by thick debris.

1 Banerjee and Shankar (2013) numerically investigated the response of extensively debris-
2 covered glaciers to rising air-temperatures and describe the dynamical response as follows:
3 during an initial period the fronts remain almost stationary and in the ablation region a slow-
4 flowing quasi-stagnant tongue develops. During this period, which may last more than 100
5 years, glaciers lose volume by thinning. After this initial period glaciers start to retreat with a
6 higher rate, while annual volume loss decreases because of thickening debris layers. Since
7 thinning rates near the fronts of the large debris-covered glaciers in the valley (Langtang,
8 Langshisha and Shalbachum Glaciers) have not yet started to significantly decrease (Figure
9 11a-c) and the glacier tongues are still dynamically active (Figure 13) it can be assumed that
10 the quasi-stationary length period will persist for these glaciers in the near future. The model
11 of Banerjee and Shankar (2013) does not account for supraglacial cliffs and lakes, which
12 likely contribute to thinning acceleration (Figure 11). However, we have shown that they
13 primarily appear on parts of the glacier tongues which are still dynamically active (Table 7).
14 Our results suggest that they become less abundant with decreasing flow. The presence of
15 cliffs and lakes therefore does not interfere with the dynamical response of debris-covered
16 glaciers as described by Banerjee and Shankar (2013).

17 Near the snout of Ghanna Glacier a deceleration in thinning rates by -80% can be clearly
18 identified (Figure 9e, Figure 11e, 4800-4850 m a.s.l.). Previous studies have provided
19 evidence that ablation rates of debris-covered ice may decrease over time as a consequence of
20 thickening debris cover, in spite of rising air-temperatures (Banerjee and Shankar, 2013;
21 Rowan et al., 2015).

22 **5.2 Elevation changes of debris-free glaciers**

23 2006-2015 downwasting rates on Yala Glacier are 0.5-1.2 m a⁻¹ higher than on Kimoshung
24 Glacier (Table 4). However, the two glaciers have a very different hypsometry (Figure S5).
25 Kimoshung Glacier has a very steep tongue that reaches to similarly low elevations as the
26 debris-covered glacier tongues (Table 1). The glacier is nearly in equilibrium with the climate
27 (Table 4), which explains the low thinning rates at low elevations (Figure S5). Currently the
28 estimated AAR of Yala Glacier is 40% (Table 1), which is a common value in the HKH
29 region (Kääb et al., 2012). The estimated AAR of 86% at Kimoshung Glacier, on the other
30 hand, corresponds to an exceptionally high value for the HKH (Khan et al., 2015). The
31 differences in ~~volume loss~~thinning rates point to the role of glacier hypsometry for the

1 response of debris-free glaciers to climatic changes (e.g. Jiskoot et al., 2009). Almost
2 balanced mass budgets in recent years (Table 4) and only minor area changes (Table 6) are
3 associated to Kimoshung Glacier. Thinning did not increase significantly with respect to the
4 period 1974-2006 (Figure 7g). Due to the steep tongue of this glacier the AAR is not sensitive
5 to changes in the ELA (Table 5). Only a small fraction of area is therefore additionally
6 exposed to temperatures above freezing level in case of atmospheric warming, which causes
7 the glacier to be less sensitive to observed warming trends in the region (Shrestha et al., 1999;
8 Lau et al., 2010). One possible explanation for the balanced conditions of Kimoshung Glacier
9 could therefore be that precipitation in recent decades remained approximately stable, which
10 agrees with the findings of studies on precipitation trends in this part of the Himalaya
11 (Shrestha et al., 2000; Immerzeel, 2008; Singh et al., 2008). However, further analysis is
12 required for justification. The mass balance of Yala Glacier, on the other hand, is sensitive to
13 fluctuations in temperature. A hypothetical rise of the ELA by 100 m at this glacier causes
14 30% of its area to turn from accumulation into ablation area (Table 5). Accordingly, thinning
15 of Yala Glacier is accelerating rapidly (Figure 9f). Due to the common AAR of Yala Glacier
16 and the extreme topography of Kimoshung Glacier it can be assumed that other debris-free
17 glaciers in the region are also thinning and that balanced conditions such as observed on
18 Kimoshung Glacier are exceptional.

19 **5.3 Differences between debris-free and debris-covered glaciers**

20 The response of debris-covered and debris-free glaciers to warming is substantially different,
21 as described in the two sections above and exemplified by the altitudinal elevation change
22 profiles in Figure 10. Our observations do not support the findings of previous studies about
23 similar present-day lowering rates of debris-covered and debris-free glacier areas at the same
24 elevation (Kääb et al., 2012; Nuimura et al., 2012; Gardelle et al., 2013). Also for debris-
25 covered elevation bands where up to 18% of the area is covered by supraglacial cliffs and
26 lakes (e.g. at Langtang tongue 5050 m a.s.l. or at Langshisha tongue 4750 m a.s.l.) thinning
27 rates do not exceed 1.8 m a^{-1} , while for Yala Glacier the lowering rates are already above this
28 value at 5250 m a.s.l. and further increase downglacier (Figure 10). Within the same
29 altitudinal range (5200-5300 m a.s.l.) thinning rates of debris-covered glaciers do not exceed
30 35%-75% of the thinning rates of Yala Glacier.

1 Regarding the mean surface elevation changes (Table 4), our observations reveal a
2 heterogeneous response to climate of both the debris-free and the debris-covered glaciers. As
3 discussed in the two sections above, there are examples for both types of glaciers where
4 thinning has increased significantly or where thinning remained approximately constant. A
5 significant difference in thinning trends between debris-free and debris-covered glaciers in
6 our sample cannot be identified. In our sample, the best predictor for thinning accelerations
7 seems to be the altitude distributions of glaciers. Glaciers with a high AAR (Kimoshung) or
8 which reach the highest elevations (Lirung) have the most balanced mass budgets and show
9 no significant changes in volume loss over time (Figure 7, Table 4). Glaciers which are most
10 sensitive to ELA changes (more than $\pm 10\%$ AAR change in response to ± 100 m ELA
11 uncertainty, Table 5) such as Yala, Langtang and Langshisha Glaciers reveal the most
12 significant thinning accelerations (Figure 7, Table 4). However, debris-free Yala Glacier is
13 currently downwasting at 60%-100% higher rates than the large debris-covered glaciers in the
14 valley. Considering the common characteristics of Yala Glacier and given that this glacier has
15 been denominated as a benchmark glacier for the Nepal Himalayas (Fujita and Nuimura,
16 2011) it seems important that future geodetic or field based studies extend our analysis to
17 larger glacier samples.

18 **5.4 Post-earthquake avalanche impacts**

19 Accumulation by debris-laden avalanches is one of the most important processes for debris-
20 covered glacier formation (Scherler et al., 2011a). The tongue of Lirung Glacier would likely
21 not exist without accumulation through avalanches (Ragettli et al., 2015). It is detached from
22 the accumulation area (Figure 1) and reaches 200-700 m lower elevations than all other
23 debris-covered glaciers (Table 1). Our volume calculations of the post-earthquake avalanche
24 impact allow quantifying the avalanche impact on mass balance and comparing it to mass loss
25 during an average year. Given the avalanche deposits remaining on Lirung tongue by 6 Oct
26 2015 (divided by the area of the tongue: 3.87 ± 0.23 m, Table 8) and the average $\Delta h/\Delta t$ rates
27 between Oct 2006 and Feb 2015 of -1.64 ± 0.10 m a⁻¹ (Figure 8d), the avalanche after the
28 earthquake compensated by 240% the volume loss of one average year. At the scale of all
29 debris-covered area in the valley this value amounts to 50% (0.52 ± 0.19 m avalanche
30 deposits and -1.02 ± 0.08 m a⁻¹ average thinning). According to Scally and Gardner (1989)
31 avalanche deposit density increases until the end of the ablation season to about 80% of ice

1 density. The mass deposits therefore compensate mass loss during a normal year by about
2 180% at Lirung tongue (40% at the catchment scale). Still, our analysis has revealed that the
3 impacts are not significant in comparison to the 2006-2015 ensemble uncertainty (Section 4.6,
4 Figure 8d and f).

5 **5.5 Comparison to other studies**

6 The four largest debris-covered glaciers in the valley (Langtang, Langshisha, Shalbachum,
7 Lirung) have been the focus of a recent geodetic mass balance study by Pellicciotti et al.
8 (2015), who reconstructed elevation and mass changes using the 1974 Hexagon DEM which
9 is also used in this study (spatial resolution 30 m) and the 2000 SRTM3 DEM (90 m). They
10 found that all four glaciers lost mass over the study period but with different rates (on average
11 -0.32 ± 0.18 m w.e. a^{-1}). We find an overall glacier mass balance for the period 1974-2006 of
12 the four glaciers which is slightly less negative (-0.22 ± 0.08 m w.e. a^{-1}). However, the results
13 match within the uncertainties. The lower uncertainty estimates by our study are justified by
14 the high resolution and quality of the 2006 Cartosat-1 DEM (Table 3). Differences in the mass
15 balance of Langtang, Lirung and Shalbachum Glacier are within uncertainty bounds and can
16 be attributed to differences in used glacier masks, study period, outlier correction approaches,
17 density assumptions and uncertainties regarding the penetration depth of the SRTM radar
18 signal (Kääb et al., 2015). However, for Langshisha Glacier we calculate a mass balance
19 which is substantially less negative than in Pellicciotti et al. (2015). While we identify almost
20 balanced conditions for the period 1974-2006 (-0.10 ± 0.08 m w.e. a^{-1} , Table 4), the mass
21 balance indicated by Pellicciotti et al. (2015) is very negative (-0.79 ± 0.18 m w.e. a^{-1}). The
22 discrepancy can be explained by the overestimated extent of the accumulation areas by
23 Pellicciotti et al. (2015) (Figure S3) in combination with unrealistic lowering rates of up
24 to -2 m a^{-1} at about 6000 m a.s.l. (Figure 4d in Pellicciotti et al., 2015). The more realistic
25 elevation change values obtained by the present study for the accumulation areas (ranging
26 between -0.4 and $+0.4$ m a^{-1} , Figure 10b) point to the need of restrictive outlier definitions and
27 the advantage of having information from multiple datasets available for gap filling.

28 Yala Glacier has been frequently visited for field measurements in the last 25 years.
29 Sugiyama et al. (2013) calculated mean thinning rates of Yala Glacier for the periods 1982-
30 1996 (-0.69 ± 0.25 m a^{-1}) and 1996-2009 (-0.75 ± 0.24 m a^{-1}) on the basis of ground
31 photogrammetry and GPS surveys. The values suggest a more moderate acceleration of

1 | ~~volume loss~~thinning rates than in our study ($-0.33 \pm 0.06 \text{ m a}^{-1}$ 1974-2006 to -0.89 ± 0.23
2 | m a^{-1} 2006-2015, Table 4). However, similarly to our study Sugiyama et al. (2013) identified
3 | a rapid acceleration of thinning rates at the lowest elevations. At higher elevations the
4 | uncertainty of photogrammetric surveys increases because of low contrast due to
5 | homogeneous snow layers.

6 | The acceleration in mass loss in recent periods identified by this study agrees with other
7 | studies from the Nepalese Himalaya which assess multi-temporal elevation changes (Bolch et
8 | al., 2011; Nuimura et al., 2012). Bolch et al. (2011) identify an increase in mass loss rates by
9 | $0.47 \text{ m w.e. a}^{-1}$ comparing the two periods 1970-2007 ($-0.32 \pm 0.08 \text{ m w.e. a}^{-1}$) and 2002-2007
10 | ($-0.79 \pm 0.52 \text{ m w.e. a}^{-1}$). Nuimura et al. (2012) calculate increasing mass losses in the same
11 | study region between 1992-2008 ($-0.26 \pm 0.24 \text{ m w.e. a}^{-1}$) and 2000-2008 ($-0.45 \pm 0.60 \text{ m}$
12 | w.e. a^{-1}). However, the identified acceleration in glacier thinning is not significant given the
13 | largely overlapping error bounds. Moreover, the mass loss estimates of Gardelle et al. (2013)
14 | for the ~~Khumbu region and same 10 glaciers as Bolch et al. (2011) in the Everest region and~~
15 | ~~for~~ the period 2001-2011 (average of $-0.41 \pm 0.21 \text{ m w.e. a}^{-1}$) are in the same order as
16 | calculated by Bolch et al. (2011) for 1970-2007. The ensemble approach of this study can
17 | therefore substantially strengthen previous conclusions that mass loss of glaciers in the
18 | Central Himalaya is accelerating. The volume changes calculated over several multi-year
19 | periods between 2006 and 2015 consistently indicate that glacier thinning has indeed
20 | accelerated (Figure 7h).

22 | **6 Conclusions**

23 | This study presents glacier volume changes of seven glaciers (five partially debris-covered,
24 | two debris-free) in the upper Langtang catchment in Nepal, using a ~~digital elevation model~~
25 | ~~(DEM)~~ from 1974 stereo Hexagon satellite data and seven DEMs derived from 2006-2015
26 | stereo or tri-stereo satellite imagery. We carefully selected elevation change maps which are
27 | least affected by uncertainty to obtain multiple independent DEM differences for the period
28 | 2006-2015.

29 | Our results point to increasing thinning rates, from $-0.24 \pm 0.08 \text{ m a}^{-1}$ in 1974-2006 to $-0.45 \pm$
30 | 0.18 m a^{-1} in 2006-2015, where the estimated confidence level of accelerated thinning rates is
31 | higher than 99%. This study therefore supports the findings of previous studies (Bolch et al.,

1 2011; Nuimura et al., 2012) that glacier wastage in the Central Himalaya is accelerating.
2 However, whereas a majority of glaciers in the study region are thinning rapidly, glaciers with
3 a high accumulation area have almost balanced mass budgets and experience no or only
4 insignificant accelerations in thinning.

5 Our observations also reveal that thinning has mostly accelerated in the upper reaches of the
6 tongues (up to +150%, comparing the periods 1974-2006 and 2006-2015), while the nearly
7 stagnant areas near the terminus show constant or decreasing thinning rates (up to -80%). The
8 highest thinning rates and the strongest increase in thinning rates can be associated to areas
9 with a high concentration of ice cliffs and supraglacial ponds. Constant or decelerating
10 thinning rates can be associated to areas with relatively homogeneous debris layers near the
11 termini of glaciers. We conclude that the response of extensively debris-covered glaciers to
12 global warming is largely determined by feedback processes associated to different surface
13 characteristics.

14 The behavior of glaciers in the study area is highly heterogeneous, and the presence or
15 absence of debris is not a good predictor for mass balance trends. However, the spatial
16 thinning patterns on debris-covered glaciers are fundamentally different than those on debris-
17 free glaciers. Debris-free glaciers in our sample present thinning rates that are linearly
18 dependent on elevation, while debris-covered glaciers have highly non-linear altitudinal
19 elevation change profiles. Our observations do not provide evidence for the existence of a so-
20 called debris-cover anomaly, where the insulating effect of thick supraglacial debris is
21 compensated by enhanced melt from exposed ice cliffs or due to high energy absorption at
22 supraglacial ponds. Within the same altitudinal range, lowering rates on debris-free Yala
23 Glacier are 35%-300% higher than on debris-covered glacier area. On debris-free Kimoshung
24 Glacier the thinning rates are similar to those of debris-covered area, but this result must be
25 explained by compressive flows that compensate for surface lowering by ablation due to its
26 exceptionally balanced conditions.

27 Geodetic mass balance studies such as this have been increasingly revealing heterogeneous
28 patterns of changes and a complex response of debris-covered glaciers that call for an
29 enhanced understanding of processes over debris-covered glaciers. Their ablation, mass
30 balance and response to climate is modulated by debris supply, transport, glacier flow, lakes
31 and cliffs developments and a complex subglacial hydrology and hydraulics that all need to be

1 understood in the future to be able to predict future changes of these glaciers over multiple
2 time scales.

3 **Acknowledgements**

4 This study is funded by the Swiss National Science Foundation (SNF) project UNCOMUN
5 (Understanding Contrasts in High Mountain Hydrology in Asia). T. Bolch acknowledges
6 funding through German Research Foundation (DFG, code BO 3199/2-1) and European
7 Space Agency (Glaciers_cci project, code 400010177810IAM). We thank Evan Miles for
8 helping with the glacier delineations and the post-processing of surface velocity data. Jakob
9 Steiner and Pascal Buri manually delineated cliffs and lakes for the inventories used in this
10 study, and they are gratefully acknowledged. We thank Etienne Berthier for the Pléiades
11 image data and Fanny Brun for her help with the identification of GCPs. DigitalGlobe
12 imagery was used to produce the WorldView-1 and 2 digital elevation models. We thank
13 Etienne Berthier (Scientific Editor), Eyjolfur Magnússon and one anonymous reviewer for
14 extensive and thorough comments that considerably helped to improve the paper.

15

1 **References**

- 2 Banerjee, A. and Shankar, R.: On the response of Himalayan glaciers to climate change, *J.*
3 *Glaciol.*, 59(215), 480–490, doi:10.3189/2013JoG12J130, 2013.
- 4 Benn, D., Gulley, J., Luckman, A., Adamek, A. and Glowacki, P. S.: Englacial drainage
5 systems formed by hydrologically driven crevasse propagation, *J. Glaciol.*, 55(191), 513–523,
6 doi:10.3189/002214309788816669, 2009.
- 7 Berthier, E. and Vincent, C.: Relative contribution of surface mass-balance and ice-flux
8 changes to the accelerated thinning of Mer de Glace, French Alps, over 1979–2008, *J.*
9 *Glaciol.*, 58(209), 501–512, doi:10.3189/2012JoG11J083, 2012.
- 10 Berthier, E., Arnaud, Y., Kumar, R., Ahmad, S., Wagnon, P. and Chevallier, P.: Remote
11 sensing estimates of glacier mass balances in the Himachal Pradesh (Western Himalaya,
12 India), *Remote Sens. Environ.*, 108(3), 327–338, doi:10.1016/j.rse.2006.11.017, 2007.
- 13 Berthier, E., Vincent, C., Magnússon, E., Gunnlaugsson, Á. Þ., Pitte, P., Le Meur, E.,
14 Masiokas, M., Ruiz, L., Pálsson, F., Belart, J. M. C. and Wagnon, P.: Glacier topography and
15 elevation changes derived from Pléiades sub-meter stereo images, *Cryosph.*, 8(6), 2275–2291,
16 doi:10.5194/tc-8-2275-2014, 2014.
- 17 Bignone, F. and Umakawa, H.: Assessment of ALOS PRISM digital elevation model
18 extraction over Japan, *Int. Arch. Photogramm. Remote Sens. Spat. Inf. Sci.*, 37, 1135–1138,
19 2008.
- 20 Bolch, T., Buchroithner, M. and Pieczonka, T.: Planimetric and volumetric glacier changes in
21 the Khumbu Himal, Nepal, since 1962 using Corona, Landsat TM and ASTER data, *J.*
22 *Glaciol.*, 54(187), 592–600, doi:10.3189/002214308786570782, 2008.
- 23 Bolch, T., Pieczonka, T. and Benn, D. I.: Multi-decadal mass loss of glaciers in the Everest
24 area (Nepal Himalaya) derived from stereo imagery, *Cryosph.*, 5(2), 349–358, doi:10.5194/tc-
25 5-349-2011, 2011.
- 26 Bolch, T., Kulkarni, A., Käab, A., Huggel, C., Paul, F., Cogley, J. G., Frey, H., Kargel, J. S.,
27 Fujita, K., Scheel, M., Bajracharya, S. and Stoffel, M.: The state and fate of Himalayan
28 glaciers., *Science*, 336(6079), 310–314, doi:10.1126/science.1215828, 2012.
- 29 Brun, F., Buri, P., Miles, E. S., Wagnon, P., Steiner, J., Berthier, E., Ragetli, S., Immerzeel,
30 W. W. and Pellicciotti, F.: Quantifying volume loss from ice cliffs on debris-covered glaciers
31 using high resolution terrestrial and aerial photogrammetry, *J. Glaciol.*, in press, 1–12,
32 doi:10.1017/jog.2016.54, 2016.
- 33 Buri, P., Pellicciotti, F., Steiner, J. F., Miles, E. S. and Immerzeel, W. W.: A grid-based model
34 of backwasting of supraglacial ice cliffs on debris-covered glaciers, *Ann. Glaciol.*, 57(71),
35 199–211, doi:10.3189/2016AoG71A059, 2016.

- 1 Burnett, M.: Hexagon (KH-9) Mapping Program and Evolution, National Reconnaissance
2 Office, Chantilly, Virginia., 2012.
- 3 Collier, E. and Immerzeel, W.: High-resolution modeling of atmospheric dynamics in the
4 Nepalese Himalaya, *J. Geophys. Res. Atmos.*, 120(19), 9882–9896,
5 doi:10.1002/2015JD023266.Received, 2015.
- 6 Dehecq, A., Gourmelen, N. and Trouve, E.: Deriving large-scale glacier velocities from a
7 complete satellite archive: Application to the Pamir–Karakoram–Himalaya, *Remote Sens.*
8 *Environ.*, 162, 55–66, doi:10.1016/j.rse.2015.01.031, 2015.
- 9 Dobhal, D. P., Mehta, M. and Srivastava, D.: Influence of debris cover on terminus retreat
10 and mass changes of Chorabari Glacier, Garhwal region, central Himalaya, India, *J. Glaciol.*,
11 59(217), 961–971, doi:10.3189/2013JoG12J180, 2013.
- 12 Fujita, K.: Effect of precipitation seasonality on climatic sensitivity of glacier mass balance,
13 *Earth Planet. Sci. Lett.*, 276(1-2), 14–19, doi:10.1016/j.epsl.2008.08.028, 2008.
- 14 Fujita, K. and Nuimura, T.: Spatially heterogeneous wastage of Himalayan glaciers., *Proc.*
15 *Natl. Acad. Sci. U. S. A.*, 108(34), 14011–14014, doi:10.1073/pnas.1106242108, 2011.
- 16 Gardelle, J., Berthier, E., Arnaud, Y. and Kääb, a.: Region-wide glacier mass balances over
17 the Pamir-Karakoram-Himalaya during 1999–2011, *Cryosph.*, 7(4), 1263–1286,
18 doi:10.5194/tc-7-1263-2013, 2013.
- 19 Holzer, N., Vijay, S., Yao, T., Xu, B., Buchroithner, M. and Bolch, T.: Four decades of
20 glacier variations at Muztagh Ata (eastern Pamir): a multi-sensor study including Hexagon
21 KH-9 and Pléiades data, *Cryosph.*, 9(6), 2071–2088, doi:10.5194/tc-9-2071-2015, 2015.
- 22 Huss, M.: Density assumptions for converting geodetic glacier volume change to mass
23 change, *Cryosph.*, 7(3), 877–887, doi:10.5194/tc-7-877-2013, 2013.
- 24 Huss, M., Juvet, G., Farinotti, D. and Bauder, A.: Future high-mountain hydrology: a new
25 parameterization of glacier retreat, *Hydrol. Earth Syst. Sci.*, 14(5), 815–829,
26 doi:10.5194/hess-14-815-2010, 2010.
- 27 Immerzeel, W.: Historical trends and future predictions of climate variability in the
28 Brahmaputra basin, *Int. J. Climatol.*, 28(2), 243–254, doi:10.1002/joc.1528, 2008.
- 29 Immerzeel, W. W., Kraaijenbrink, P. D. a., Shea, J. M., Shrestha, A. B., Pellicciotti, F.,
30 Bierkens, M. F. P. and de Jong, S. M.: High-resolution monitoring of Himalayan glacier
31 dynamics using unmanned aerial vehicles, *Remote Sens. Environ.*, 150, 93–103,
32 doi:10.1016/j.rse.2014.04.025, 2014a.
- 33 Immerzeel, W. W., Petersen, L., Ragettli, S. and Pellicciotti, F.: The importance of observed
34 gradients of air temperature and precipitation for modeling runoff from a glacierized
35 watershed in the Nepalese Himalayas, *Water Resour. Res.*, 50(3), 2212–2226,
36 doi:10.1002/2013WR014506, 2014b.

- 1 Jiskoot, H., Curran, C. J., Tessler, D. L. and Shenton, L. R.: Changes in Clemenceau Icefield
2 and Chaba Group glaciers, Canada, related to hypsometry, tributary detachment, length–slope
3 and area–aspect relations, *Ann. Glaciol.*, 50(53), 133–143,
4 doi:10.3189/172756410790595796, 2009.
- 5 Juen, M., Mayer, C., Lambrecht, ~~a~~A., Han, H. and Liu, S.: Impact of varying debris cover
6 thickness on ablation: a case study for Koxkar Glacier in the Tien Shan, *Cryosph.*, 8(2), 377–
7 386, doi:10.5194/tc-8-377-2014, 2014.
- 8 Käab, A., Berthier, E., Nuth, C., Gardelle, J. and Arnaud, Y.: Contrasting patterns of early
9 twenty-first-century glacier mass change in the Himalayas, *Nature*, 488(7412), 495–498,
10 doi:10.1038/nature11324, 2012.
- 11 Käab, A., Treichler, D., Nuth, C. and Berthier, E.: Brief Communication: Contending
12 estimates of 2003–2008 glacier mass balance over the Pamir–Karakoram–Himalaya,
13 *Cryosph.*, 9(2), 557–564, doi:10.5194/tc-9-557-2015, 2015.
- 14 Kargel, J., Leonard, G., Shugar, D. et al.: Geomorphic and geologic controls of geohazards
15 induced by Nepal’s 2015 Gorkha earthquake, *Science*, 351(6269), 1–18,
16 doi:10.1126/science.aac8353, 2016.
- 17 Khan, A., Naz, B. S. and Bowling, L. C.: Separating snow, clean and debris covered ice in the
18 Upper Indus Basin, Hindukush-Karakoram-Himalayas, using Landsat images between 1998
19 and 2002, *J. Hydrol.*, 521, 46–64, doi:10.1016/j.jhydrol.2014.11.048, 2015.
- 20 [Lacroix, P.: Landslides triggered by the Gorkha earthquake in the Langtang valley, volumes](#)
21 [and initiation processes, *Earth, Planets Sp.*, 68\(1\), 46, doi:10.1186/s40623-016-0423-3, 2016.](#)
- 22 Lamsal, D., Sawagaki, T. and Watanabe, T.: Digital terrain modelling using Corona and
23 ALOS PRISM data to investigate the distal part of Imja Glacier, Khumbu Himal, Nepal, *J.*
24 *Mt. Sci.*, 8(3), 390–402, doi:10.1007/s11629-011-2064-0, 2011.
- 25 Lau, W. K. M., Kim, M.-K., Kim, K.-M. and Lee, W.-S.: Enhanced surface warming and
26 accelerated snow melt in the Himalayas and Tibetan Plateau induced by absorbing aerosols,
27 *Environ. Res. Lett.*, 5(2), 025204, doi:10.1088/1748-9326/5/2/025204, 2010.
- 28 Leprince, S., Barbot, S., Ayoub, F. and Avouac, J.-P.: Automatic and precise
29 orthorectification, coregistration, and subpixel correlation of satellite images, application to
30 ground deformation measurements, *IEEE Trans. Geosci. Remote Sens.*, 45(6), 1529–1558,
31 2007.
- 32 Magnússon, E., Muñoz-Cobo Belart, J., Pálsson, F., Ágústsson, H. and Crochet, P.: Geodetic
33 mass balance record with rigorous uncertainty estimates deduced from aerial photographs and
34 lidar data – Case study from Drangajökull ice cap, NW Iceland, *Cryosph.*, 10(1), 159–177,
35 doi:10.5194/tc-10-159-2016, 2016.
- 36 Mattson, L. E., Gardner, J. S. and Young, G. J.: Ablation on Debris Covered Glaciers: an
37 Example from the Rakhiot Glacier, Punjab, Himalaya, *IAHS Publ.*, 218, 289–296, 1993.

- 1 Maurer, J. and Rupper, S.: Tapping into the Hexagon spy imagery database: A new automated
2 pipeline for geomorphic change detection, *ISPRS J. Photogramm. Remote Sens.*, 108, 113–
3 127, doi:10.1016/j.isprsjprs.2015.06.008, 2015.
- 4 Mayer, C., Lambrecht, A., Belo, M., Smiraglia, C. and Diolaiuti, G.: Glaciological
5 characteristics of the ablation zone of Baltoro glacier, Karakoram, Pakistan, *Ann. Glaciol.*,
6 43(1), 123–131, doi:10.3189/172756406781812087, 2006.
- 7 Mihalcea, C., Mayer, C., Diolaiuti, G., Lambrecht, A., Smiraglia, C. and Tartari, G.: Ice
8 ablation and meteorological conditions on the debris-covered area of Baltoro glacier,
9 Karakoram, Pakistan, *Ann. Glaciol.*, 43(1), 292–300, doi:10.3189/172756406781812104,
10 2006.
- 11 Mihalcea, C., Mayer, C., Diolaiuti, G., Agata, C. D., Smiraglia, C., Lambrecht, A.,
12 Vuillermoz, E. and Tartari, G.: Spatial distribution of debris thickness and melting from
13 remote-sensing and meteorological data, at debris-covered Baltoro glacier, Karakoram,
14 Pakistan, *Ann. Glaciol.*, 48(1), 49–57, doi:10.3189/172756408784700680, 2008.
- 15 Miles, E. S., Pellicciotti, F., Willis, I. C., Steiner, J. F., Buri, P. and Arnold, N. S.: Refined
16 energy-balance modelling of a supraglacial pond, Langtang Khola, Nepal, *Ann. Glaciol.*,
17 57(71), 29–40, doi:10.3189/2016AoG71A421, 2016a.
- 18 | Miles, E. S., Willis, I. C., Arnold, N. S., [Steiner, J.](#) and Pellicciotti, F.: Spatial, seasonal, and
19 interannual variability of supraglacial ponds in the Langtang Valley of Nepal, 1999 to 2013, *J.*
20 *Glaciol.*, under revision, 2016b.
- 21 Noh, M.-J. and Howat, I. M.: Automated stereo-photogrammetric DEM generation at high
22 latitudes: Surface Extraction with TIN-based Search-space Minimization (SETSM) validation
23 and demonstration over glaciated regions, *GIScience Remote Sens.*, 52(2), 198–217,
24 doi:10.1080/15481603.2015.1008621, 2015.
- 25 Nuimura, T., Fujita, K., Fukui, K., Asahi, K., Aryal, R. and Ageta, Y.: Temporal Changes in
26 Elevation of the Debris-Covered Ablation Area of Khumbu Glacier in the Nepal Himalaya
27 since 1978, *Arctic, Antarct. Alp. Res.*, 43(2), 246–255, doi:10.1657/1938-4246-43.2.246,
28 2011.
- 29 Nuimura, T., Fujita, K., Yamaguchi, S. and Sharma, R. R.: Elevation changes of glaciers
30 revealed by multitemporal digital elevation models calibrated by GPS survey in the Khumbu
31 region, Nepal Himalaya, 1992–2008, *J. Glaciol.*, 58(210), 648–656,
32 doi:10.3189/2012JoG11J061, 2012.
- 33 Nuimura, T., Sakai, A., Taniguchi, K., Nagai, H., Lamsal, D., Tsutaki, S., Kozawa, A.,
34 Hoshina, Y., Takenaka, S., Omiya, S., Tsunematsu, K., Tshering, P. and Fujita, K.: The
35 GAMDAM glacier inventory: a quality-controlled inventory of Asian glaciers, *Cryosph.*, 9(3),
36 849–864, doi:10.5194/tc-9-849-2015, 2015.

- 1 Nuth, C. and Kääb, A.: Co-registration and bias corrections of satellite elevation data sets for
2 quantifying glacier thickness change, *Cryosph.*, 5(1), 271–290, doi:10.5194/tc-5-271-2011,
3 2011.
- 4 Nuth, C., Schuler, T. V., Kohler, J., Altena, B. and Hagen, J. O.: Estimating the long-term
5 calving flux of Kronebreen, Svalbard, from geodetic elevation changes and mass-balance
6 modelling, *J. Glaciol.*, 58(207), 119–133, doi:10.3189/2012JoG11J036, 2012.
- 7 Östrem, G.: Ice melting under a thin layer of moraine, and the existence of ice cores in
8 moraine ridges, *Geogr. Ann.*, 41(4), 228–230, 1959.
- 9 Paul, F., Barrant, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S.
10 P., Konovalov, V., Bris, R. Le, Mölg, N., Nosenko, G., Nuth, C., Pope, a., Racoviteanu, a.,
11 Rastner, P., Raup, B., Scharrer, K., Steffen, S. and Winsvold, S.: On the accuracy of glacier
12 outlines derived from remote-sensing data, *Ann. Glaciol.*, 54(63), 171–182,
13 doi:10.3189/2013AoG63A296, 2013.
- 14 Pellicciotti, F., Stephan, C., Miles, E., Immerzeel, W. W. and Bolch, T.: Mass-balance
15 changes of the debris-covered glaciers in the Langtang Himal, Nepal, 1974–99, *J. Glaciol.*,
16 61(225), doi:10.3189/2015JoG13J237, 2015.
- 17 Pieczonka, T. and Bolch, T.: Region-wide glacier mass budgets and area changes for the
18 Central Tien Shan between ~1975 and 1999 using Hexagon KH-9 imagery, *Glob. Planet.*
19 *Change*, 128, 1–13, doi:10.1016/j.gloplacha.2014.11.014, 2015.
- 20 Pieczonka, T., Bolch, T. and Buchroithner, M.: Generation and evaluation of multitemporal
21 digital terrain models of the Mt. Everest area from different optical sensors, *ISPRS J.*
22 *Photogramm. Remote Sens.*, 66(6), 927–940, doi:10.1016/j.isprsjprs.2011.07.003, 2011.
- 23 Pieczonka, T., Bolch, T., Junfeng, W. and Shiyin, L.: Heterogeneous mass loss of glaciers in
24 the Aksu-Tarim Catchment (Central Tien Shan) revealed by 1976 KH-9 Hexagon and 2009
25 SPOT-5 stereo imagery, *Remote Sens. Environ.*, 130, 233–244,
26 doi:10.1016/j.rse.2012.11.020, 2013.
- 27 Pratap, B., Dobhal, D. P., Mehta, M. and Bhambri, R.: Influence of debris cover and altitude
28 on glacier surface melting: a case study on Dokriani Glacier, central Himalaya, India, *Ann.*
29 *Glaciol.*, 56(70), 9–16, doi:10.3189/2015AoG70A971, 2015.
- 30 Racoviteanu, A. E., Arnaud, Y., Williams, M. W. and Manley, W. F.: Spatial patterns in
31 glacier characteristics and area changes from 1962 to 2006 in the Kanchenjunga–Sikkim area,
32 eastern Himalaya, *Cryosph.*, 9(2), 505–523, doi:10.5194/tc-9-505-2015, 2015.
- 33 Ragetti, S., Pellicciotti, F., Immerzeel, W. W., Miles, E. S., Petersen, L., Heynen, M., Shea,
34 J. M., Stumm, D., Joshi, S. and Shrestha, A.: Unraveling the hydrology of a Himalayan
35 catchment through integration of high resolution in situ data and remote sensing with an
36 advanced simulation model, *Adv. Water Resour.*, 78, 94–111,
37 doi:10.1016/j.advwatres.2015.01.013, 2015.

- 1 Reid, T. D. and Brock, B. W.: Assessing ice-cliff backwasting and its contribution to total
2 ablation of debris-covered Miage glacier, Mont Blanc massif, Italy, *J. Glaciol.*, 60(219), 3–13,
3 doi:10.3189/2014JoG13J045, 2014.
- 4 Reynolds, J.: On the formation of supraglacial lakes on debris-covered glaciers, *IAHS Publ.*,
5 (264), 153–161, 2000.
- 6 Rowan, A. V., Egholm, D. L., Quincey, D. J. and Glasser, N. F.: Modelling the feedbacks
7 between mass balance, ice flow and debris transport to predict the response to climate change
8 of debris-covered glaciers in the Himalaya, *Earth Planet. Sci. Lett.*, 430, 427–438,
9 doi:10.1016/j.epsl.2015.09.004, 2015.
- 10 Sakai, A. and Fujita, K.: Formation conditions of supraglacial lakes on debris-covered
11 glaciers in the Himalaya, *J. Glaciol.*, 56(195), 177–181, doi:10.3189/002214310791190785,
12 2010.
- 13 Sakai, A., Nakawo, M. and Fujita, K.: Melt rate of ice cliffs on the Lirung Glacier, Nepal
14 Himalayas, 1996, *Bull. Glacier Res.*, 16, 57–66, 1998.
- 15 Sakai, A., Takeuchi, N., Fujita, K. and Nakawo, M.: Role of supraglacial ponds in the
16 ablation process of a debris-covered glacier in the Nepal Himalayas, *Debris-Covered Glaciers*,
17 *IAHS Publ.*, 265, 119–130, 2000.
- 18 Sakai, A., Nakawo, M. and Fujita, K.: Distribution characteristics and energy balance of ice
19 cliffs on debris-covered glaciers, Nepal Himalaya, *Arctic, Antarct. Alp. Res.*, 34(1), 12–19,
20 doi:10.2307/1552503, 2002.
- 21 Sapiano, J., Harrison, W. and Echelmeyer, K.: Elevation, volume and terminus changes of
22 nine glaciers in North America, *J. Glaciol.*, 44(146), 119–135, doi:10.3198/1998JoG44-146-
23 119-135, 1998.
- 24 Scally, F. De and Gardner, J.: Evaluation of avalanche mass determination approaches: an
25 example from the Himalaya, Pakistan, *J. Glaciol.*, 35(120), 248–252, 1989.
- 26 Scherler, D., Leprince, S. and Strecker, M.: Glacier-surface velocities in alpine terrain from
27 optical satellite imagery—Accuracy improvement and quality assessment, *Remote Sens.*
28 *Environ.*, 112(10), 3806–3819, doi:10.1016/j.rse.2008.05.018, 2008.
- 29 Scherler, D., Bookhagen, B. and Strecker, M. R.: Hillslope-glacier coupling: The interplay of
30 topography and glacial dynamics in High Asia, *J. Geophys. Res.*, 116(F02019), 1–21,
31 doi:10.1029/2010JF001751, 2011a.
- 32 Scherler, D., Bookhagen, B. and Strecker, M. R.: Spatially variable response of Himalayan
33 glaciers to climate change affected by debris cover, *Nat. Geosci.*, 4(3), 156–159,
34 doi:10.1038/ngeo1068, 2011b.
- 35 Schwitter, M. and Raymond, C.: Changes in the longitudinal profiles of glaciers during
36 advance and retreat, *J. Glaciol.*, 39(133), 582–590, 1993.

- 1 Shrestha, A., Wake, C., Mayewski, P. and Dibb, J.: Maximum temperature trends in the
2 Himalaya and its vicinity: An analysis based on temperature records from Nepal for the
3 period 1971-94, *J. Clim.*, 12, 2775–2786, doi:10.1175/1520-
4 0442(1999)012<2775:MTTITH>2.0.CO;2, 1999.
- 5 Shrestha, A., Wake, C., Dibb, J. and Mayewski, P.: Precipitation fluctuations in the Nepal
6 Himalaya and its vicinity and relationship with some large scale climatological parameters,
7 *Int. J. Climatol.*, 20(3), 317–327, doi:10.1002/(SICI)1097-0088(20000315)20:3<317::AID-
8 JOC476>3.0.CO;2-G, 2000.
- 9 Singh, P., Kumar, V., Thomas, T. and Arora, M.: Changes in rainfall and relative humidity in
10 river basins in northwest and central India, *Hydrol. Process.*, 22(16), 2982–2992,
11 doi:10.1002/hyp.6871, 2008.
- 12 Steiner, J. F., Pellicciotti, F., Buri, P., Miles, E. S., Immerzeel, W. W. and Reid, T. D.:
13 Modelling ice-cliff backwasting on a debris-covered glacier in the Nepalese Himalaya, *J.*
14 *Glaciol.*, 61(229), 889–907, doi:10.3189/2015JoG14J194, 2015.
- 15 Steiner, J. F., Buri, P., Miles, E. S., Ragetti, S. and Pellicciotti, F.: Life and death of ice cliffs
16 and lakes on debris covered glaciers - insights from a new dataset, *Geophys. Res. Abstr.*,
17 18(EGU2016-13922), 2016.
- 18 Sugiyama, S., Fukui, K., Fujita, K., Tone, K. and Yamaguchi, S.: Changes in ice thickness
19 and flow velocity of Yala Glacier, Langtang Himal, Nepal, from 1982 to 2009, *Ann. Glaciol.*,
20 54(64), 157–162, doi:10.3189/2013AoG64A111, 2013.
- 21 Surazakov, A. and Aizen, V.: Positional Accuracy Evaluation of Declassified Hexagon KH-9
22 Mapping Camera Imagery, *Photogramm. Eng. Remote Sens.*, 76(5), 603–608,
23 doi:10.14358/PERS.76.5.603, 2010.
- 24 Tadono, T. and Shimada, M.: Calibration of PRISM and AVNIR-2 onboard ALOS “Daichi,”
25 *IEEE Trans. Geosci. Remote Sens. Sens.*, 47(12), 4042–4050,
26 doi:10.1109/TGRS.2009.2025270, 2009.
- 27 Thompson, S., Benn, D. I., Mertes, J. and Luckman, A.: Stagnation and mass loss on a
28 Himalayan debris-covered glacier: processes, patterns and rates, *J. Glaciol.*, 1–19,
29 doi:10.1017/jog.2016.37, 2016.
- 30 Tiwari, P., Pande, H., Punia, M. and Dadhwal, V. K.: Cartosat-I: Evaluating mapping
31 capabilities, *Int. J. Geoinformatics*, 4(1), 51–56 [online] Available from:
32 <http://creativitycity.gssc.osaka-cu.ac.jp/IJG/article/view/609> (Accessed 26 January 2016),
33 2008.
- 34 Wang, D. and Käab, A.: Modeling Glacier Elevation Change from DEM Time Series, *Remote*
35 *Sens.*, 7(8), 10117–10142, doi:10.3390/rs70810117, 2015.
- 36 Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H.,
37 Xu, B., Pu, J., Lu, A., Xiang, Y., Kattel, D. B. and Joswiak, D.: Different glacier status with

1 atmospheric circulations in Tibetan Plateau and surroundings, *Nat. Clim. Chang.*, 2(9), 663–
2 667, doi:10.1038/nclimate1580, 2012.

3 Ye, Q., Bolch, T., Naruse, R., Wang, Y., Zong, J., Wang, Z., Zhao, R., Yang, D. and Kang,
4 S.: Glacier mass changes in Rongbuk catchment on Mt. Qomolangma from 1974 to 2006
5 based on topographic maps and ALOS PRISM data, *J. Hydrol.*, 530, 273–280,
6 doi:10.1016/j.jhydrol.2015.09.014, 2015.

7

1 Figures and Tables

2 Table 1. Characteristics of the studied glaciers in the upper Langtang catchment. The
 3 measures are based on the SRTM 1 Arc-Second Global DEM and glacier outlines of 2006.

Name	Area	Debris	Mean slope	Mean slope	AAR**	Elevation range
	km ²	cover	%	glacier tongue*		
1 Langtang	46.5	15.5	17.1	7.2	52%	4479-6615
2 Langshisha	16.3	4.5	17.7	7.5	55%	4415-6771
3 Shalbachum	10.2	2.6	16.9	9.1	52%	4231-6458
4 Lirung	6.5	1.1	34.0	9.9	49%	4044-7120
5 Ghanna	1.4	0.7	20.9	15.5	15%	4721-5881
6 Kimoshung	4.4	-	24.4	32.1	86%	4385-6648
7 Yala	1.9	-	22.7	20.3	40%	5122-5676

*Here we consider the debris-covered area for glaciers with debris-covered tongues and all glacier area below 5400 m a.s.l. for debris-free glaciers.

**Assuming an equilibrium line altitude of 5400 m a.s.l. (Sugiyama et al., 2013; Ragetti et al., 2015)

4 Table 2. Remote-sensing data used

Sensor	Date of acquisition	Stereo mode (b/h-ratio)	Spatial/radiometric Resolution	Role
Hexagon KH-9	23 Nov 1974	Stereo (0.4)	6-9m/8-bits	DEM differencing, glacier outlines
Cartosat-1	15 Oct 2006	Stereo (0.62)	2.5m/10-bits	DEM differencing, glacier outlines
Cartosat-1	9 Nov 2009	Stereo (0.62)	2.5m/10-bits	DEM differencing, velocities, glacier outlines
ALOS PRISM	3 Dec 2010	Tri-stereo (0.5)	2.5m/8-bits	DEM differencing, velocities, glacier outlines
SPOT6	21 Apr 2014	Tri-stereo (0.5)	1.5m/12-bits	DEM differencing, glacier outlines
WorldView-2	2 Feb 2015	Stereo (0.5)	0.46m/11-bits	DEM differencing
WorldView-3	22 Feb 2015	Stereo (0.5)	0.31m/11-bits	DEM differencing
SPOT7	7 Mai 2015	Tri-stereo (0.64)	1.5m/12-bits	DEM differencing, glacier outlines
SPOT7	6 Oct 2015	Tri-stereo (0.68)	1.5m/12-bits	DEM differencing
Pléiades	1 and 9 Nov 2014	Across track stereo (0.4)	0.5m/12-bits	Basis for georectification

5

1 Table 3. Mean* uncertainties associated to different sets of elevation change ($\Delta h/\Delta t$) maps.

	No. of maps in category	All glacier area	Debris-covered glacier area
All $\Delta h/\Delta t$ maps, $\Delta t < 4$ a	9	1.18 m a ⁻¹	0.47 m a ⁻¹
All $\Delta h/\Delta t$ maps, $4 \text{ a} \leq \Delta t < 10$ a	12	0.29 m a ⁻¹	0.12 m a ⁻¹
All $\Delta h/\Delta t$ maps involving Hexagon 1974 DEM	7	0.07 m a ⁻¹	0.03 m a ⁻¹
DEM involved, $4 \text{ a} \leq \Delta t < 10$ a			
Cartosat-1 Oct 2006	5	0.22 m a ⁻¹	0.09 m a ⁻¹
Cartosat-1 Nov 2009	4	0.29 m a ⁻¹	0.11 m a ⁻¹
ALOS-PRISM Dec 2010	4	0.43 m a ⁻¹	0.19 m a ⁻¹
SPOT6 April 2014	2	0.24 m a ⁻¹	0.09 m a ⁻¹
WorldView Feb 2015	3	0.32 m a ⁻¹	0.14 m a ⁻¹
SPOT7 May 2015	3	0.31 m a ⁻¹	0.12 m a ⁻¹
SPOT7 October 2015	3	0.24 m a ⁻¹	0.10 m a ⁻¹

* Uncertainties associated to individual maps are shown in Table S1

2 Table 4. Glacier volume and mass changes 1974-2006, ensemble mean 2006-2015*.

	Average elevation differences (m a ⁻¹)		Average mass balance (m w.e. a ⁻¹)	
	Nov1974-Oct2006	Ensemble mean* 2006-2015	Nov1974-Oct2006	Ensemble mean* 2006-2015
Glaciers				
Langtang	-0.28 ± 0.08	-0.55 ± 0.13	-0.24 ± 0.08	-0.47 ± 0.13
Langshisha	-0.12 ± 0.09	-0.45 ± 0.19	-0.10 ± 0.08	-0.38 ± 0.18
Shalbachum	-0.43 ± 0.08	-0.53 ± 0.19	-0.36 ± 0.09	-0.45 ± 0.18
Lirung	-0.17 ± 0.13	-0.22 ± 0.16	-0.14 ± 0.11	-0.19 ± 0.14
Ghanna	-0.51 ± 0.05	-0.46 ± 0.43	-0.43 ± 0.07	-0.39 ± 0.36
Kimoshung	0.07 ± 0.13	-0.02 ± 0.17	0.05 ± 0.10	-0.02 ± 0.13
Yala	-0.33 ± 0.06	-0.89 ± 0.23	-0.28 ± 0.07	-0.76 ± 0.24
Average	-0.24 ± 0.08	-0.45 ± 0.18	-0.21 ± 0.08	-0.38 ± 0.17
Debris-covered areas				
Langtang	-0.79 ± 0.03	-0.91 ± 0.05	-0.67 ± 0.07	-0.78 ± 0.10
Langshisha	-0.69 ± 0.03	-1.16 ± 0.23	-0.58 ± 0.07	-0.98 ± 0.25
Shalbachum	-0.78 ± 0.04	-1.30 ± 0.20	-0.66 ± 0.08	-1.10 ± 0.23
Lirung	-1.03 ± 0.05	-1.67 ± 0.59	-0.87 ± 0.10	-1.42 ± 0.56
Ghanna	-0.58 ± 0.03	-0.50 ± 0.20	-0.49 ± 0.06	-0.43 ± 0.19
Average	-0.78 ± 0.03	-1.02 ± 0.18	-0.66 ± 0.07	-0.87 ± 0.20

*Average of 6 overlapping periods between Oct 2006 and Oct 2015 (Figure S4)

3

4

1 Table 5. Sensitivity to outlier correction and Equilibrium Line Altitude (ELA) definitions. $\Delta_{2\sigma}$
2 is the difference in results if a 2σ -level is used to define outliers at all area types, instead of a
3 3σ -level above and a 1σ -level below the ELA. The estimated ELA is 5400 m a.s.l. (Sugiyama
4 et al., 2013; Ragetti et al., 2015). $\Delta_{\text{ELA}\pm 100\text{m}}$ represents the differences in results obtained with
5 an ELA at 5500 m a.s.l. in comparison to results obtained with an ELA at 5300 m a.s.l.

Name	AAR		Elevation differences (m a^{-1})			
	ELA -100 m	ELA +100 m	Nov1974-Oct2006		Ensemble mean 2006-2015	
			$\Delta_{\text{ELA}\pm 100\text{m}}$	$\Delta_{2\sigma}$	$\Delta_{\text{ELA}\pm 100\text{m}}$	$\Delta_{2\sigma}$
1 Langtang	61%	43%	0.02	-0.01	-0.01	0.01
2 Langshisha	60%	45%	0.06	-0.09	0.02	-0.02
3 Shalbachum	60%	37%	-0.23	0.08	0.04	-0.02
4 Lirung	52%	46%	-0.01	0.01	0.05	-0.01
5 Ghanna	20%	12%	-0.01	0.02	0.00	0.04
6 Kimoshung	88%	80%	0.07	0.06	0.05	-0.16
7 Yala	70%	13%	-0.13	-0.04	-0.11	0.02
All Glacier Area	61%	44%	0.00	-0.01	0.02	-0.01

Sensitivity values that exceed uncertainty ranges as indicated in Table 4 are printed in bold letters.

6

7 Table 6. Glacier area changes over the periods 1974-2006 and 2006-2015.

ID	Glacier name	1974-2006		2006-2015	
		km^2	$\% \text{ a}^{-1}$	km^2	$\% \text{ a}^{-1}$
1	Langtang	-2.65 ± 0.03	-0.17 ± 0.01	-0.45 ± 0.07	-0.11 ± 0.02
2	Langshisha	-0.48 ± 0.09	-0.09 ± 0.02	-0.13 ± 0.05	-0.09 ± 0.04
3	Shalbachum	-0.28 ± 0.06	-0.08 ± 0.02	-0.03 ± 0.04	-0.04 ± 0.04
4	Lirung	-0.45 ± 0.08	-0.20 ± 0.03	-0.05 ± 0.05	-0.08 ± 0.08
5	Ghanna	-0.16 ± 0.03	-0.33 ± 0.05	-0.05 ± 0.01	-0.40 ± 0.12
6	Kimoshung	-0.11 ± 0.01	-0.08 ± 0.01	-0.02 ± 0.01	-0.05 ± 0.02
7	Yala	-0.31 ± 0.03	-0.43 ± 0.05	-0.31 ± 0.03	-1.77 ± 0.16

8

9

1 Table 7. Characteristics of the debris-covered tongues (debris-covered glacier area excluding
2 tributary branches).

	Cliff Area	Lake Area	Mean velocity	% stagnant
Langtang	10.0%	3.3%	5.9 m a ⁻¹	31%
Langshisha	10.5%	2.3%	7.0 m a ⁻¹	20%
Shalbachum	10.3%	2.6%	5.4 m a ⁻¹	29%
Lirung	8.0%	2.3%	2.8 m a ⁻¹	48%
Ghanna	3.2%	0.4%	1.6 m a ⁻¹	85%

Cliff and lake area corresponds to the percentage of 30-m pixels containing cliffs/lakes (median of 6 available cliff and lake maps from the period 2006-2015). Mean velocity is calculated on the basis of 2009-2010 surface velocities (Figure 13). To discriminate moving ice from quasi-stagnant ice we use a threshold of 2.5 m a⁻¹ (cf. Scherler et al., 2011b), which also corresponds to the approximate uncertainty of remote-sensing derived surface velocity.

3 Table 8. Elevation changes of debris-covered glacier tongues due to avalanches triggered by
4 the Nepal earthquake on 25 April 2015. The first three data columns provide the volume
5 changes of avalanche affected area ($\Delta h > 5$ m in May 2015) divided by the total debris-cover
6 area (Table 1).

	21 Apr 2014- 25 Apr 2015* (m)	25 Apr 2015- 7 May 2015 (m)	25 Apr 2015- 6 Oct 2015 (m)	6 Oct 2015, volume remaining (%)
Langtang**	-0.10 ±0.05	1.33 ±0.42	0.42 ±0.20	31.3%
Langshisha	-0.04 ±0.05	0.32 ±0.37	0.10 ±0.19	31.6%
Shalbachum	-0.11 ±0.05	0.74 ±0.35	0.31 ±0.20	42.5%
Lirung	-0.87 ±0.06	6.79 ±0.38	3.87 ±0.23	57.0%
Average	-0.13 ±0.05	1.31 ±0.35	0.52 ±0.19	39.5%

*Estimation based on average annual melt Oct 2006 – Apr 2014

**Only lower part (south of 28°19'N), upper part not on April 2014 scene

7

8 Figure 1. Map of the upper Langtang catchment. The numbers on the map correspond to the
9 glaciers listed in Table 1. Monsoon snow-cover frequency is based on Landsat 1999 to 2013
10 land cover classifications (Miles et al., 2016b). 1974 glacier area (dotted lines) is shown for
11 the seven studied glaciers only.

12

13 Figure 2. a) Uncertainty estimates of average elevation change rates ($\Delta h/\Delta t$) per individual
14 glacier and per debris-covered tongue. ~~The central mark is the median of the ensemble ($\Delta h/\Delta t$
15 maps that are rejected according to Section 3.2.5 are excluded).~~ b) Ensemble of stereo
16 matching scores per individual glacier and debris-covered tongue and c) per glacier area

1 above and below the estimated ELA. The central marks correspond to the median of all
2 ~~DEMs (except Hexagon 1974 and WorldView 2015 DEMs for which matching scores are not~~
3 ~~available)- $\Delta h/\Delta t$ maps or DEMs in the ensemble.~~ The edges of each box are the 25th and 75th
4 percentiles. The whiskers extend to the most extreme data points.

5
6 Figure 3. Off-glacier elevation change error ($\Delta h/\Delta t$) per 50-m elevation band. The black line
7 represents the median error in the ensemble of six $\Delta h/\Delta t$ maps ~~(excluding $\Delta h/\Delta t$ maps that are~~
8 ~~rejected according to Section 3.2.5).~~ Error bars represent 95% confidence intervals. The color
9 bars represent hypsometries of glacier area, off-glacier area and debris-covered glacier areas,
10 respectively.

11
12 Figure 4. Elevation change rates ($\Delta h/\Delta t$) derived from a) Hexagon Nov 1974 and Cartosat-1
13 Oct 2006 DEMs and (b) Cartosat-1 Oct 2006 and WorldView Feb 2015 DEMs.

14
15 Figure 5. Uncertainties in elevation change rates ($\Delta h/\Delta t$) ~~in~~ as a function of the time interval
16 between DEMs (Δt). Median results of all available 28 $\Delta h/\Delta t$ maps. Error bars extend to the
17 most extreme data points.

18
19 Figure 6. Supraglacial cliffs and lakes as identified from the Oct 2006 Cartosat-1 satellite
20 image: a) Langtang and Ghanna Glaciers, b) Shalbachum Glacier, c) Lirung Glacier, d)
21 Langshisha Glacier. Cliff area shows the median fraction (%) of 30-m pixels per 50m
22 elevation band that contain cliffs, considering all 6 available cliff maps from 2006-2015.

23
24 Figure 7. Mean elevation change rates ($\Delta h/\Delta t$) per period and glacier. For better readability,
25 only the maximum width of error bounds corresponding to individual periods 2006-2015 are
26 shown. Note that the 1974-2006 time scale is not linear (dashed dark blue line).

27
28 Figure 8. Mean elevation change rates ($\Delta h/\Delta t$) per period and debris-covered glacier area. For
29 better readability, only the maximum width of error bounds corresponding to individual

1 | periods 2006-2015 are shown. Note that the 1974-2006 time scale is not linear (dashed dark
2 | blue line).

3
4 | Figure 9. Altitudinal distribution of mean annual elevation change ($\Delta h/\Delta t$) over 50 m
5 | elevation bands of debris-covered tongues (debris-covered area of each glacier excluding
6 | tributary branches). Uncertainty bounds correspond to uncertainty was a function of elevation
7 | derived for each $\Delta h/\Delta t$ map individually (Figure 3).

8
9 | Figure 10. Altitudinal distribution of mean annual elevation change ($\Delta h/\Delta t$) and altitudinal
10 | distribution of glacier area (%) over 50 m elevation bands of selected glaciers. Uncertainty
11 | bounds correspond to uncertainty was a function of elevation derived for each $\Delta h/\Delta t$ map
12 | individually (Figure 3). Ensemble median values shown here are used to replace missing data
13 | in the accumulation areas of glaciers after outlier exclusion (Section 3.2.3). Note that the
14 | x-axis ranges are different for each sub-figure.

15
16 | Figure 11: Altitudinal distribution of cliff and lake area fractions, glacier velocity and changes
17 | in thinning rates. Cliff and lake area is shown as % of 30-m pixels containing cliffs/lakes per
18 | 50 m elevation band, whereas the values represent the median of 6 available cliff and lake
19 | maps from the period 2006-2015. Glacier velocities (m/a) represent the median per 50 m
20 | elevation band of data shown in Figure 4+13 and error bars represent the standard deviation in
21 | pixel values per elevation band. Changes in thinning rates ($\Delta(\Delta h/\Delta t)$ [m/a]) are calculated
22 | comparing 1974-2006 and the 2006-2015 ensemble-mean. Negative $\Delta(\Delta h/\Delta t)$ values
23 | represent thinning accelerations. Error bars represent the maximum variations in $\Delta(\Delta h/\Delta t)$
24 | considering all individual periods within the 2006-2015 ensemble.

25
26 | Figure 12. Avalanche affected sections of Lirung and Langtang glacier, pre- and after the
27 | earthquake on 25 April 2015, and corresponding surface elevation changes (Δh). Imagery
28 | ©Airbus DS 2014/2015.

1 Figure 13: Surface velocities 2009-2010 cropped to catchment boundaries. Values have units
2 of meters per year and are derived by cross-correlation feature tracking. Off-glacier velocities
3 are shown in transparent color.
4

1 **Supplementary Figures and Tables**

2

3 Figure S1. Off-glacier mean elevation differences and b) standard deviations (σ). On each
4 box, the central mark is the median and the edges of the box are the 25th and 75th percentiles
5 of the ensemble (~~six maps of $\Delta h/\Delta t$ maps that are rejected according to Section 3.2.5 are~~
6 ~~excluded~~). The whiskers extend to the most extreme data points. ME_{noglac} and σ_{noglac} are
7 calculated excluding the steepest slopes (slope $<45^\circ$). For $ME2_{\text{noglac}}$ and $\sigma2_{\text{noglac}}$ areas with a
8 monsoon snow-cover frequency higher than 20% (Figure 1) are masked out (slope $<45^\circ$,
9 $sc \leq 20\%$). For $ME3_{\text{noglac}}$ and $\sigma3_{\text{noglac}}$ the threshold slope is defined as the 95th percentile of
10 debris-covered glacier slope (slope $<18^\circ$, $sc \leq 20\%$).

11

12 Figure S2. 2006-2015 ensemble of elevation change rates ($\Delta h/\Delta t$).

13

14 Figure S3. Comparison of glacier outlines for Langshisha Glacier used by the present study
15 and by Pellicciotti et al. (2015).

16

17 Figure S4. Altitudinal distribution of mean annual elevation changes ($\Delta h/\Delta t$) of debris-
18 covered tongues (as in Figure 9) and altitudinal distribution of cliff and lake area fractions (as
19 in Figure 11).

20

21 Figure S5. Altitudinal distribution of mean annual elevation changes ($\Delta h/\Delta t$) and altitudinal
22 distribution of glacier area (%) of debris-free Yala and Kimoshung Glaciers. For better
23 readability only the elevation changes corresponding to the two periods Oct 2006 – Oct 2015
24 and Oct 2009 – Oct 2015 are shown.

1 Table S1. Uncertainties associated to each elevation change map.

Δh map	All glacier area	Debris-covered glacier area
Nov 1974 - Oct 2006	0.08 m a ⁻¹	0.03 m a ⁻¹
Nov 1974 - Nov 2009	0.07 m a ⁻¹	0.03 m a ⁻¹
Nov 1974 - Dec 2010	0.09 m a ⁻¹	0.04 m a ⁻¹
Nov 1974 - Apr 2014	0.05 m a ⁻¹	0.02 m a ⁻¹
Nov 1974 - Feb 2015	0.07 m a ⁻¹	0.03 m a ⁻¹
Nov 1974 - May 2015	0.07 m a ⁻¹	0.03 m a ⁻¹
Nov 1974 - Oct 2015	0.06 m a ⁻¹	0.02 m a ⁻¹
Oct 2006 - Nov 2009	0.41 m a ⁻¹	0.17 m a ⁻¹
Oct 2006 - Dec 2010	0.46 m a ⁻¹	0.20 m a ⁻¹
Oct 2006 - Apr 2014	0.12 m a ⁻¹	0.05 m a ⁻¹
Oct 2006 - Feb 2015	0.21 m a ⁻¹	0.08 m a ⁻¹
Oct 2006 - May 2015	0.16 m a ⁻¹	0.06 m a ⁻¹
Oct 2006 - Oct 2015	0.15 m a ⁻¹	0.06 m a ⁻¹
Nov 2009 - Dec 2010	2.15 m	0.88 m
Nov 2009 - Apr 2014	0.36 m a ⁻¹	0.14 m a ⁻¹
Nov 2009 - Feb 2015	0.29 m a ⁻¹	0.12 m a ⁻¹
Nov 2009 - May 2015	0.26 m a ⁻¹	0.10 m a ⁻¹
Nov 2009 - Oct 2015	0.24 m a ⁻¹	0.10 m a ⁻¹
Dec 2010 - Apr 2014	0.62 m a ⁻¹	0.27 m a ⁻¹
Dec 2010 - Feb 2015	0.45 m a ⁻¹	0.20 m a ⁻¹
Dec 2010 - May 2015	0.51 m a ⁻¹	0.21 m a ⁻¹
Dec 2010 - Oct 2015	0.33 m a ⁻¹	0.14 m a ⁻¹
Apr 2014 - Feb 2015	1.27 m	0.53 m
Apr 2014 - May 2015	1.08 m	0.35 m
Apr 2014 - Oct 2015	0.52 m a ⁻¹	0.19 m a ⁻¹
Feb 2015 - May 2015	1.61 m	0.62 m
Feb 2015 - Oct 2015	1.69 m	0.71 m
May 2015 - Oct 2015	1.23 m	0.47 m

2