

Reply to comments by G. Liu on “Mapping and inventorying active rock glaciers in the Northern Tien Shan (China) using satellite SAR interferometry”

General comments

After more than two decades silence, this manuscript offers us new knowledge, evidences and research methods about rock glaciers and permafrost distribution in the Northern Tien Shan of China. The paper based on 261 active rock glaciers which were recognized by combines SAR interferometry and optical images from Google Earth, give a detail discussing of their locations, geomorphic parameters, and down-slope velocities, and marking permafrost lower limit. This research is reference significance for alpine Periglacial landform research and permafrost mapping in the remote regions.

We thank Prof. G. Liu for his positive comments on our approach and results. We also appreciate his careful consideration and detailed comments. Our replies are highlighted in blue.

Specific comments

(1) Recognition of rock glaciers:

- According to the manuscript and the supplement, most of the 261 active rock glaciers were correctly identified, yet some misreading appeared, for example, ARG 70, ARG 131 should be MARG.

Authors: Thanks for your correction. We have accordingly corrected the types of ARG70 and ARG131 as MARGs. We also revised the relevant statistics related to the MARGs and TARGs through the paper.

- By the limitation of satellite data and research technique, a large number of small rock glaciers are not identified and compiled, especially the talus derived. For example, rock glaciers in the head water of Urumqi River reported by Cui and Zhu (1989), Zhu (1992), Zhu et al (1992), Liu et al (1995), not mentioned by the authors.

Authors: It is true that we may have missed some small rock glaciers because the InSAR phase maps have a moderate spatial resolution of 15 m. We have highlighted this potential limitation of our method in Section 4 (see Lines 259–268).

Following the reviewer’s suggestion, we have added a discussion subsection to compare our inventory with the previous studies on rock glaciers in the Tien Shan (Cui and Zhu, 1989; Gorbunov et al. 1992; Zhu 1992; Zhu et al.1992; Liu et al. 1995). We have reviewed and compared the altitude distribution, geomorphic descriptions and surface velocities of the

rock glaciers that have been documented in these studies with our results (see Lines 409–435). However, we can only perform a qualitative, not quantitative, comparison. This is because (1) the exact locations of the rock glaciers documented in these studies are not available; (2) the method we adopted to map active rock glaciers is significantly different from the method used in these previous studies, making the comparison impractical.

- ARG 94 was recognized as TARG, in the same way, site a (43.6429°N, 85.4292°E ;) and b (43.6380°N, 85.4284°E) should be TARG. This greatly reduces the accuracy of number, regional and altitude distribution of the rock glaciers. This therefore, affect use of active rock glacier to determine permafrost lower limit.

Authors: We have investigated the two locations of site ‘a’ (43.6429°N, 85.4292°E) and ‘b’ (43.6380°N, 85.4284°E) on the Google Earth images. From the geomorphic features, we interpret that these two sites host two talus-derived rock glaciers. However, we found that our InSAR phase map suffers from severe de-correlation problem at these two locations, likely due to radar shadows. Therefore, we excluded them from our inventory. We have pointed out that the number of ARGs we compiled is a conservative estimation of all the ARGs in the NTS, and listed several reasons for this kind of conservative estimation (please see Lines 259–268). This is also why we gave a range of estimations rather than exact values of the lower altitude limits of permafrost, as described and justified in Section 4.4. Nevertheless, as discussed in Section 5.3, the distribution of the ARGs we inventoried are in good agreement with these three existing permafrost maps in the Northern Tien Shan.

(2) Identification of the initial line point (ILP) and front line point (FLP):

- It is not very clear how to determine ILP for talus derived rock glacier, for example ARG22, 51; and moraine derived rock glacier, for example ARG 50, view Google Earth, the ILP located at moraine covered glacier.

Authors: As described in the first paragraph in Section 3.3 (Lines 220–230), we determined the initial line point (ILP) and front line point (FLP) of a rock glacier based on both the InSAR phase measurements and the geomorphological features from the Google Earth images. The variations of the InSAR phase represent ground movements, and thus we can outline an active rock glacier (ARG) based on the InSAR phase variations. We interpreted the active boundary in the rooting zone of an ARG as the initial line where the permafrost starts to creep. We then used the central point of the initial line as the ILP. The FLP of the ARG can be determined based on the similar method. In addition, the FLP is generally located in the lowest place where a rock glacier can reach, which can be easily identified from the Google Earth images. For example, Fig. R1(a) below shows the outline of the ARG22 determined from the InSAR phase variations. Fig. R1(b) shows the corresponding

Google Earth image. We have rewritten the relevant text in Section 3.3 to clarify our method for identifying the ILP and FLP (see Lines 220–230).




Fig. R1 ARG22 and its ILP and FLP. (a) shows the InSAR phase map for the ARG22, and the blue line denotes the active boundary determined from the phase variations. (b) shows the Google image.

- Some MARG mentioned in the manuscript, both ILP and FLP might be misreading, for example, ARG 95, the ILP is at glacier, the FLP should be down slope at 43.6226°N , 85.4043°E .

Authors: Thank you very much for your corrections. We have corrected the ILP and FLP for the ARG 95 based on the interpretation of historical images from the Google Earth. We have also checked the ILP/FLP for each identified ARG to improve the accuracy of our inventory.

- ARG 219, 220, and 221 seem merge ARG 157, form a combined RG, the FLP might reach forest zone – see Google Earth.

Authors: Some nearby ARGs could merge and form a combined rock glacier. In such cases, we classified each sub-ARG as individual rock glacier for following two reasons. (1) The sub-ARGs have different aspect angles. (2) We want to include the identified ARGs as many as possible. We have added one sentence to explain the combined ARGs in our revised manuscript, please see Lines 187-189.

(3) Surface velocity: It is better offering surface velocity by several years' data. In the discussion, comparing with Cui and Zhu, Zhu.

Authors: It is a good idea to map the surface velocities of ARGs using several year's SAR

data. The multi-temporal observations would help us to reveal the seasonal fluctuations of the velocities of the ARGs (e.g., Liu et al., 2013). However, due to the rough topography in the NTS area, very few ALOS PALSAR interferometric pairs are suitable for InSAR time series analysis. The suitable interferometric pairs should have both small temporal span and perpendicular baseline, thus maintaining good phase coherence. In this study, we selected the PALSAR interferograms that have a temporal span of 46/92 days and perpendicular baselines smaller than 600 m. In addition, we only used the PALSAR images acquired in summer to form the interferograms except for the Path 503. We have added some descriptions about the SAR data we used in this study, please see Lines 136–140.

We have added a sub-section (Section 5.1) in the discussion part to compare our inventory with the previous studies. In which, we compared the surface velocities derived from our InSAR measurements with the field observations conducted by Cui and Zhu (1989) and Zhu (1992a). Please see Lines 409–435.

(4) Indication of lower limit of permafrost: Though the manuscript give detailed discussion, the estimated lower limit of permafrost is well below field survey (Qiu et al., 1983) might be caused by missing the small RGs, especially the talus derived ones.

Authors: Thank you for pointing out this possible reason that would influence the lower limit of permafrost we estimated. The InSAR phase maps we generated have a spatial resolution of about 15 meters. It is possible that some small rock glacier in the shadows of radar images cannot be identified. We have added this possible reason in the revised version (see Lines 259–268).

(5) References: Page 12, 4.4, references suggested: Jin HJ et al, 1993, Journal of Glaciology and Geocryology, 15(1). Qiu GQ, 1993, Journal of Glaciology and Geocryology, 15(1). Zhao L et al. 2010, Journal of Glaciology and Geocryology, 32(2).

Authors: Thanks for your suggestions. We have added these three references in the revised manuscript, please see Lines 372, 405–407.

3. Technical corrections: a compact listing of purely technical corrections, typing errors, typographical corrections, etc.

(1) Page 3 line 77-78: “There is a lack of studies on surface velocities of rock glaciers or an inventory containing the locations of the surveyed rock glaciers.” See Cui and Zhu 1989.

Authors: The relevant sentence has been rewritten as: “The studies on surface velocities of rock glaciers are very limited except for the field surveys at several specific sites conducted by Cui and Zhu (1989) and Zhu (1992a).”

(2) Page 4 line 97 (Zhu et al., 1992b) should add now reference. Zhao L. et al. 2010.

Authors: The reference of Zhao et al., (2010) has been added, please see Line 110.

(3) Page 4 line 102 “(Cui and Zhu, 1989; Zhu et al., 1992a, 1992b; Liu et al., 1995)” - (Cui and Zhu, 1989; Zhu, 1992a, Zhu et al, 1992b; Liu et al., 1995)

Authors: The citation format has been corrected in the revised version.

(4) Page 6 line 151-152 “Debris-covered glaciers are usually covered with uniformly thin debris layer, whereas rock glaciers debris cover is less homogenous and coarser.” – What is the basis?

Authors: Considering the comments of the referee 3 (Dr. Bolch), we have rewritten the relevant sentences as follows (please see Lines 174–180):

“We distinguished ARGs from debris-covered glaciers based on the de-correlation conditions of interferograms and their different visual features on the Google Earth images. Compared with rock glaciers, debris-covered glaciers generally move much faster (Janke et al., 2015), which results in large areas of de-correlation in our PALSAR interferograms. The surface of a debris-covered glacier is usually characterized by ice cliffs and supra-glacial lakes (Bolch et al., 2007). And the rooting zones of debris-covered glaciers are continuous with clean glacier ice (Davies et al., 2013; Lukas et al., 2007).”

(5) Page 10 line 239 “IPLAs”? Not mentioned in the text.

Authors: This is a typo. “IPLAs” has been changed to “ILPAs”.

(6) Page 11-12, 4.3 Surface velocities of the active rock glaciers Reference Cui and Zhu, 1989.

Authors: We have added a discussion subsection (Section 5.1) to compare our inventory with the previous studies of rock glaciers in the Tien Shan (Lines 409–435).

(7) Page 17 line 442-444. The paragraph (4) seems could be omitted.

Authors: We have removed the paragraph (4).

(8) Page 17 line 446-448 “This inventory offers a baseline dataset for the further investigations on permafrost modeling, slope stability, and water resource, etc.” – why slope stability, and water resource? Not mentioned in the text.

Authors: We have added some descriptions and references in the “Introduction” part to clarify the significance of rock glaciers on the water resource in the semi-arid region and the potential hazards in the high mountains. Rock glaciers serve as important freshwater reserves because of the potential melting of their internal ice in the runoff season. Active rock glaciers

may cause mass waste hazards on steep slopes when they move downslope, and thus threatening the sensible infrastructures in some downhill areas (Schoeneich et al., 2015). Please see Lines 31–33, 37–38.

Reply to comments by T. Strozzi on “Mapping and inventorying active rock glaciers in the Northern Tien Shan (China) using satellite SAR interferometry”

1. General comments

The manuscript by Wang et al. nicely describes the application of SAR interferometry and Google Earth optical images to the mapping of active rock glaciers in the Northern Tien Shan. The structure of the paper is solid, the objectives of the work are clearly stated, the employed technology is well explained, the results are nicely described and illustrated, and the conclusions are well formulated. Although not completely novel, the approach of mapping and inventorying active rock glaciers from SAR interferometry and optical images is here throughout applied for the first time in the high mountains of western China, revealing new insights into rock glaciers as proxies of permafrost. The same approach presented by Wang et al. can be used for the systematic investigations of rock glaciers, and thus permafrost, over other remote mountainous locations. Considering the excellent work done by the authors in the redaction of the manuscript, I only have a few specific comments.

We thank Dr. T. Strozzi for his thoughtful consideration and helpful comments on our manuscript. We have addressed all the comments below. Our replies are highlighted in blue.

2. Specific comments

1. The short description about error sources in SAR interferograms at line 138 could be expanded a little bit to avoid the impression that their effects are not well considered or underestimated. In particular, the use of a coarse resolution DEM as SRTM results in uncompensated phase signals that scale up with the baseline of the interferograms. The authors state that maximum baselines considered in their studies are 600 m, but I would expect here a quantitative estimate of the phase disturbances that may occur in high mountains as a consequence of typical SRTM height errors or artefacts.

Authors: We have added a sentence copied below to demonstrate the contribution of topographic error to the interferometric phase (please see Lines 151–156):

“The contribution of the topographic error to the interferometric phase is proportional to the perpendicular baseline, the radar slant range, and the radar look angle (Rosen et al., 2000). The maximum perpendicular baseline of the interferograms produced in this study is about 600 m. Adopting the vertical accuracy of the digital elevation model as its nominal value of 16 m (Farr et al., 2004), we estimate that the residual topographic phase in the InSAR measurements would be about 0.92 radians, corresponding to 1.7 cm.

In addition, a statement about the different distortions that occur with respect to the slope

and orientation of the rock glaciers and the satellite line-of-sight direction should be included.

Authors: We have rewritten the relevant part in Section 4 (copied below) to include the influences of geometric distortions to the identification of ARGs (Please see Lines 259–268):

“The number of ARGs we compiled is a conservative estimation of all the ARGs in the NTS due to the following three reasons. First, we may have missed some ARGs due to the geometric distortions such as shadows and layovers in the SAR images. For the PALSAR images used in this study, the rock glaciers facing east with the slope angles larger than 51.3° are in the radar shadows, and the rock glaciers facing west with slope angles larger than 38.7° are in the layover regions. These distortions would result in significantly phase decorrelation and make it difficult to identify rock glaciers in these regions. Second, ARGs facing nearly north or south might also have been missed, as InSAR (using images from all space missions, including ALOS) is not sensitive to ground motions along these directions. Finally, some small ARGs could not be identified as the interferogram maps have a moderate resolution of about 15 m.”

Then, on the same paragraph, a short comment on the typical atmospheric disturbances at L-band with respect to the size of the active rock glaciers would be beneficial to strengthen the potential of the technology.

Authors: We have added two sentences (copied below) to describe the atmosphere disturbance on the L-band SAR interferometry, with concerning the size of the active rock glaciers, please see Lines 156–159.

“The atmospheric (including tropospheric and ionospheric) effects are generally manifest themselves as long wavelength signals on the order of 1–10 km in interferogram maps (Hanssen, 2001), thus can be assumed nearly constant over a specific rock glacier. By using a local reference point just outside each rock glacier (see section 3.2), we effectively removed these large-scale atmospheric errors.”

Finally, Barboux et al. (2015) found out that phase unwrapping is the major limiting factor to the use of SAR interferometry for monitoring active rock glaciers in the Swiss Alps. In their paper, Wang et al. are not discussing at all possible phase unwrapping errors. I did some rapid calculations with the active rock glaciers velocities reported by the authors in the Northern Tien Shan. If maximum down-slope velocities of active rock glaciers in this region are about 114 cm yr^{-1} , then the maximum line-of-sight velocities should be about 2/3 of the down-slope direction, i.e. about 76 cm yr^{-1} . Over a time period of 46 days this would correspond to about 10 cm or less than one fringe an L-band. Indeed, in the Northern Tien

Shan phase unwrapping is not a relevant issue, but this is not the case in many other mountain regions. Therefore, a short comment on phase unwrapping with respect to the velocities of the active rock glaciers should be included in the manuscript.

Authors: We have added some words to demonstrate the unwrapping issues for addressing the surface velocities of rock glaciers using InSAR (Please see Lines 205–210). We also carefully examined unwrapped interferograms and confirmed no unwrapping errors over the rock glaciers we mapped.

“Previous studies indicated that phase unwrapping is the major limiting factor to the use of SAR interferometry for monitoring active rock glaciers (Barboux et al., 2015). Phase unwrapping may fail in the areas where phase gradients are large due to fast slope movements. To minimize the phase unwrapping errors, we applied the phase filtering with a window of 8×8 pixels and masked out the decorrelated areas with a coherence threshold of 0.3 before phase unwrapping.”

2. At line 63, the paper by Strozzi et al. (2010) is about landslides, not rock glaciers. Use instead Strozzi et al. (2004): Strozzi T., A. Kääb and R. Frauenfelder, Detecting and quantifying mountain permafrost creep from in situ inventory, space-borne radar interferometry and airborne digital photogrammetry, *Int. J. Remote Sensing*, Vol. 25, No. 15, pp. 2919-2931, doi: 10.1080/0143116042000192330 2004.

Authors: We have changed the reference “Strozzi et al. (2010)” to “Strozzi et al. (2004)” in Line 65.

3. Figure 9a is nearly impossible to interpret, there is too much information and there are too many colors and symbols. An alternative representation of this image should be proposed by the authors.

Authors: We have broken Figure 9a into two sub-figures to improve interpretation. We also changed Figures 9(c–d) as a new Figure 10.

Reply to comments by Tobias Bolch on “Mapping and inventorying active rock glaciers in the Northern Tien Shan (China) using satellite SAR interferometry”

We thank Dr. Bolch for his detailed and insightful review of the discussion paper. We have addressed all the comments and made the suggested changes in the revised version of our manuscript. Our point-by-point replies (in blue) to the critical comments (in black) are listed below.

The most important issues are:

- The delineation of the rock glaciers seems to be unprecise as e.g. visible from the figures: The upper boundary may be hard to define precisely but some more efforts should be made; is it not clear from Fig. 2 which criteria was used and the boundary is very probably not rectangular. It seems to me from Figs. 2b and 4 that exposed rocks are included. The lower boundary visible in Fig. 2b is also too rough as parts which clearly belong to the rock glacier are missed while others (e.g. even a part of the river were included). This needs to be improved as good as possible for all outlines so that at least no obvious errors are made.

Authors: We are grateful for the suggestions. We have checked the outline of each inventoried ARG and re-delineated some of them based on the InSAR phase and optical images. We also re-plotted Fig. 2 and Fig. 4 to correct the errors pointed out by the reviewer.

- Rock glaciers provide a hint where the lower limit of permafrost is located. However, the lower limit of rock glaciers varies significant due to topographic factors and the blocky material favors cooler temperatures allowing rock glaciers to exist at elevations where permafrost is otherwise unlikely (see. e.g. modelling studies for the Northern Tien Shan in Kazakhstan/Kyrgyzstan, see references). I'd therefore suggest rather to discuss the suitability of rock glaciers for investigation the permafrost occurrence (as you partly already did) than presenting new results of a lower limit of the permafrost occurrence which is highly uncertain. The authors should also keep in mind that there is no clear limit but that especially in mountains the permafrost occurrence is very heterogeneous.

Authors: Thanks very much for the insightful suggestion.

The purpose of this study is to propose a new method for inventorying active rock glaciers in the periglacial high mountains. We chose the Northern Tien Shan in China as a study area and then used the inventoried talus-derived active rock glaciers (TARGs) to infer the lower limit of permafrost. As suggested by the reviewer, there may be a high uncertainty in the estimates of permafrost limits due to the heterogeneous thermal region in mountainous environment. Previous studies conducted by Gorbunov et al. (2004) in Transili Alatau Tien Shan revealed a significant difference between the thermal regime of coarse blocky materials

and adjacent fine-grained soils. Mean annual temperatures inside the coarse debris could be 2.5–4.0 °C cooler than mean annual air temperatures (MAATs), which could cause rock glaciers occur at elevations where permafrost is unlikely to exist. However, the statistics of our inventory show that the MAAT at the TARGs is -5.4 °C, and 83 % of the TARGs have the MAAT lower than -4 °C. This indicates that the TARGs we used for estimating the lower limit of permafrost are in a cold environment, and thus reinforcing the credibility of the estimates. We now include such analysis and interpretation in the revised manuscript (Lines 456–459).

It is indeed important to discuss the suitability of rock glaciers for investigating the permafrost occurrence. An accurate permafrost distribution map will be helpful as a reference for this kind of investigation. However, the available permafrost distribution maps in our study (the IPA permafrost map, the CAS permafrost map, and the PZI map) all have very coarse spatial resolutions. Therefore, it is difficult to assess what characteristics of rock glaciers should have that would be suitable for inferring the permafrost distribution based on our present dataset. We decide to keep the original discussion on the lower limit of permafrost in the region, which is a range, and leave the use of individual rock glaciers for permafrost occurrence in the future work.

- There is limited discussion where the authors put their approach and results into context. There are several studies existing which were presenting rock glacier inventories and further information in the neighbouring Kazakh and Kyrgyz Tien Shan. Several ones are in Russian which might be difficult to understand (but may still be considered as they provide valuable information) but there are also several published in English (especially by A. Gorbunov, see references). One more recent paper co-authored by Gorbunov presents also a topographic analysis which would be very interesting to compare your results too.

Authors: We have added an addition discussion section (Lines 409–435) to compare our inventory with the previous studies of rock glaciers in the NTS (Cui and Zhu, 1989; Zhu, 1992a; Zhu et al., 1992b; Liu et al., 1995) and the Kazakh/Kyrgyz Tien Shan (Gorbunov et al., 1992, 1998). The rock glacier studies in the NTS conducted by Cui and Zhu (1989) and Zhu (1992a) reported that most of the identified rock glaciers are tongue-shaped, located at altitudes between 3300 m and 3900 m, and facing north, which are generally consistent with our inventory. The studies of rock glaciers in the Kazakh/Kyrgyz Tien Shan are concentrated in the Djungar Ala Tau, the Ile Alatau, and the Kungöj Ala-Too mountain ranges. The rock glaciers in the Djungar Ala Tau lay lower altitudes comparing to the NTS, probably due to their more northerly latitude. While the altitudinal distributions of the rock glaciers in Alatau and Kungöj Ala-Too are similar with that in the NTS. These previous studies also revealed a homogeneity pattern of the ARGs surface velocities.

Additionally, we found that the topographic analysis conducted by Bolch and Gorbunov (2014) was focusing on the influences of the contributing area parameters (e.g. its area or the headwall height) on the area and minimum altitudes of the rock glaciers. However, such contributing area parameters were not investigated in our inventory. Therefore, it is impractical to compare our results with the topographic analysis of Bolch and Gorbunov (2014).

- Recent findings show a clear seasonal behaviour of the surface velocity (e.g. Wirz et al. 2016). Hence, it is important to mention the acquisition period of the data used to calculate the velocity also in the text. The seasonal effect should also be considered in the discussion.

Authors: The ALOS PALSAR data we used were acquired between 2007 and 2009, and all the data were acquired in summer except for Path 503. Each interferogram spans 46 or 92 days. We state this explicitly in Section 3 (Lines 136 and 139) and Table 1. The seasonal effects may contribute to the spatial heterogeneity of the surface velocities due to the acquisition times for each image pairs are different. We state this point in the revised version in Lines 361–367.

- With respect to the discussion I suggest to include a separate discussion section where you put your approach and results into context of the existing literature. Currently, the discussion section contains only a comparison of the rock glacier derived permafrost estimates to permafrost distribution estimates.

Authors: Following this suggestion, we have added a separated discussion section (Section 5.1) to compare our approach and results with the previous studies of rock glaciers in Tien Shan (Lines 409–435).

- A minor but important point: The Tien Shan is a large mountain range stretching from Uzbekistan into China. Considering the Tien Shan as a whole using “Northern Tien Shan” is not correct. You should rather use “Northeastern Tien Shan” in the title and elsewhere.

Authors: The study area is completely located in China and is termed as “Northern Tien Shan” traditionally. We find very few papers use “Northeastern Tien Shan” in the literature. To avoid confusion, we have changed the “Northern Tien Shan” to “Northern Tien Shan of China” in the title through the paper.

Specific comments:

Abstract:

L10: Rock glaciers are widespread not only in western China but in the whole of Tien Shan.

Authors: We have revised the sentences (copied below), please see Lines 10–11.

“Rock glaciers are widespread in the Tien Shan Mountain. However, rock glaciers in the Chinese part of the Tien Shan have not been systematically investigated for more than two decades.”

L11: There are few recent studies, but your statement is true for Western China. Please correct.

Authors: Done. Please see Line 11.

L24: The approach is interesting for global rock glacier mapping and not only for western China.

Authors: We have rewritten the sentence to “...to map rock glaciers over mountain ranges globally” (Line 22).

1. Introduction:

L28: I am missing the classical monography by Barsch (Barsch 1996) in the references. I'd rather cite Haerberli et al. (2006) instead of 2010 here.

Authors: We have added the reference ‘Barsch (1996)’ in Line 28 and changed the reference ‘Haerberli (2010)’ to ‘Haerberli et al. (2006)’.

L36: Not sure if a definition of inactive and relict rock glaciers are needed here. This is a cryospheric journal where the readers should know such basic knowledge.

Authors: We removed the definitions of inactive and relict rock glaciers.

L45: Brenning (2005) already mentioned it and used the term intact for active and inactive rock glaciers. You would be able to distinguish which is a major advantage.

Authors: Here we want to express that it is difficult to differentiate active rock glaciers from inactive ones directly through visual interpretation. Thus we chose to keep the original wording, i.e. “active and inactive” (Line 46).

L47ff: You should mention also somewhere that multi-temporal optical images can and were also used to investigate rock glacier velocity (e.g. Gorbunov et al. 1992, by visual interpretation; Käab et al. 1997 by feature tracking).

Authors: We have added some words to include the use of multi-temporal optical images for investigating the surface velocities of ARGs (Lines 61–63).

L68: Mention here at least one of the existing studies.

Authors: We have added two references (Kenyi and Kaufmann, 2003; Liu et al., 2013) (Line 69).

L74: Include here few of the existing studies of the Kazakh and Kyrgyz Tien Shan.

Authors: We have added several sentences (copied below) (Lines 75–80).

“Rock glaciers are abundant in the high mountains of Central Asia and western China, such as the Tien Shan, Hindu Kush Himalayan, Kunlun Shan and Hengduan Shan. More than 1500 rock glaciers have been inventoried in the Kazakh and Kyrgyz Tien Shan (Gorbunov et al., 1992;1998). Schmid et al. (2015) compiled an inventory with 702 rock glaciers identified from Google Earth images in the vast Hindu Kush Himalayan region. Using these inventories, recent studies have further discussed the rock glaciers-permafrost interactions (Bolch and Gorbunov, 2014; Schmid et al., 2015).”

L80: You may mention here the study by Bolch and Gorbunov for the nearby Northern Tien Shan in Kazakhstan and Kyrgyzstan and maybe also Schmid et al. (2015)

Authors: We have added citations to these papers (Line 80).

2. Study area:

General: Provide more details about the general characteristics of the Tien Shan, the subdivision, and especially the climatic conditions.

Authors: Thanks for your suggestions. We have added detail descriptions of the subdivision and climatic conditions of the Northern Tien Shan of China (Lines 97–99, 111–117).

L97: Be more specific about the annual precipitation amount. The west is wetter than the east but not really wet. The climate is only relative humid for the continental conditions.

Authors: We have added several sentences (copied below) in this section (Lines 111–117).

“There are two long-term meteorological stations in the NTS. The Daxigou meteorological station (86.84° E, 43.11° N) in the eastern part is at about 3540 m above sea level. The Snow-cover and Avalanche Research Station (SARS, 84.40° E, 43.26° N) lies in a valley in the western part of the NTS, at an elevation of about 1776 m. The mean annual air temperature (MAAT) and the mean annual precipitation around the Daxigou station were about –6 °C and 405 mm in 1990–2008, respectively (Sun et al., 2013). The MAAT near the SARS was about 1.3 °C, falling to –9.4 °C at 3580 m. The mean annual precipitation at the SARS was about 830 mm (Shi et al., 2009).”

L102: You cite here almost the same studies than in L75 in the introduction. I suggest to cite in the intro those which provide information about the larger regions or from surrounding

ranges and in the study region section the specific ones.

Authors: Following the suggestion, we have rewritten the relevant sentences in introduction (Lines 75–80) and cited the studies of rock glaciers in the Kazakh and Kyrgyz Tien Shan and the Hindu Kush Himalayan region, such as Bolch and Gorbunov, 2014, Gorbunov et al., 1992, and Schmid et al., 2015.

3. Methodology:

General: The other reviewers are more experts in SAR processing. Therefore I will not comment on the technical aspects here. However, I'd like to see a better figure where the authors present the identification of moving surface based on the wrapped interferometric phase. Fig. 3 is interesting in this regard but I find it hard to understand how you identified surface displacements based on this image. Maybe a larger image or a zoom would help. Provide a short information about the quality and suitability of the images available at the time of the study in google earth, e.g. had all images good snow conditions and where all of very high resolution?

Authors: We have added two figures (Figure 3c and 3d) for showing the identification of the moving surface based on wrapped interferometric phase. Fig. 3c shows the outline of the rock glacier determined from the variations of phase map, and Fig. 3d shows the corresponding optical image for the rock glacier in Google Earth. Additionally, we selected the Google Earth images that were cloud free and were taken in summer season as the limited extent of snow cover to readily identify ARGs. We state this in the revised version in Lines 164–166.

L148: I agree that active rock glaciers have usually little or no vegetation, but not always (e.g. there are trees on rock glaciers in other parts of the Tien Shan). Hence, write “have usually little or no vegetation” or similar.

Authors: Corrected, please see Line 172.

L151: It is not true that debris-covered glaciers “are usually covered with uniformly thin debris layer”. There are manifold studies which show that the debris thickness usually increases towards the terminus and that the surface of a debris-covered glacier is characterised by ice cliffs and supraglacial lakes. It is partly hard to distinguish clearly between debris-covered glaciers and rock glaciers as gradual transitions to moraine-derived rock glaciers exist especially in continental conditions.

Authors: Thanks for pointing out this. We have rewritten the relevant descriptions for the debris-covered glacier (copied below). Please see Lines 174–180.

“We distinguished ARGs from debris-covered glaciers based on the de-correlation conditions of interferograms and their different visual features on the Google Earth images. Compared with rock glaciers, debris-covered glaciers generally move much faster (Janke et al., 2015), which results in large areas of de-correlation in our PALSAR interferograms. The surface of a debris-covered glacier is usually characterized by ice cliffs and supra-glacial lakes (Bolch et al., 2007). And the rooting zones of debris-covered glaciers are continuous with clean glacier ice (Davies et al., 2013; Lukas et al., 2007).”

L161: What is a rooting zone “Z” and where is it in Fig. 1?

Authors: We have enlarged the denotation of the zone “Z” in Fig. 1.

L184f: I find the abbreviations ILP (initial line point) and FLP (front line point) not intuitive as you are mainly interested in the altitude. Humlum (1998) uses RILA (rock glacier initiation line altitude). As you are referring to the maximum and minimum altitude I’d use h_{\min} and h_{\max} . But you may decide.

Authors: Thanks for your suggestion. We define the ILP and FLP to conveniently determine the parameters of the inventoried ARGs, such as length, aspect, and slope. Therefore, we decide to keep the abbreviations.

L197f: It can be quite difficult to identify the upper boundary. Hence, InSAR is quite promising. Provide more details and examples for how you identified the upper boundary. Ridges and furrows are typical for compressive flow in the lower rock glacier area and, hence, you might have missed parts if you only use these criteria.

Authors: We now added one more example to show the InSAR-determined outline of an ARG, please see Figs. 3c and 3d and Lines 198–200.

L200: Should be Gruber (2012).

Authors: Corrected.

L206: Why do you mention “lower limit of the permafrost distribution” here. You are quantifying the rock glacier parameters.

Authors: We have rewritten the sentence (copied blow), please see Lines 243–244.

“The uncertainties of the FLP and ALP altitudes are determined by the errors of the SRTM digital elevation model, which has a nominal vertical accuracy of less than 16 m (Farr et al., 2004).”

4. Results:

L222f: Move to methods.

Authors: We have moved the sentence to the Method section (Lines 204–205).

L250ff: What is about the influence of the general topography? It could be worth to compare the aspects of the rock glaciers to the aspect distribution of the investigated mountain ranges. The aspect distribution of the rock glaciers in Kazakh and Kyrgyz Northern Tien Shan is clearly influences by the topography.

Authors: We calculated the aspect distribution of the mountain ranges in the study area and found the number of north-facing and south-facing slopes are similar. Therefore, we inferred that the topography in the NTS is not the controlling factoring for the aspect distributions of the MARGs. We presumed that the larger percentage of north-facing MARGs in their quantum than the TARGs is due to aspect distribution of the glaciers. While we noted that the majority of glaciers in the NTS are north-facing.

L252: Kaldybayev et al. (2016) investigate glaciers in Dzhungar Alatau which is close to you study region but is not the Northern Tien Shan.

Authors: Thanks for pointing out this. The reference “Kaldybayev et al., 2016” has been removed in the revised version.

L266: Be careful with the statement about the lower limit. See my general comment above.

Authors: We have added some word to declare the lower limit, please see our replies to the general comments.

L270: What is about the precipitation? I would assume the precip is also of importance.

Authors: Agree. We have rewritten the relevant sentences to state the possible influence of precipitation on the altitude distribution of rock glaciers (Lines 319–322).

L278f: This is an important point and should be discussed more in detail.

Authors: We have further discussed the different altitude distributions of ARGs in the western and eastern parts of the NTS. The paragraph reads now as follows (Lines 323–331):

“Additionally, we found different altitude distributions of ARGs in the western and eastern parts (Fig. 7a). The nearly linear pattern in the east is more apparent than that in the west. We performed a T-test on the similarity of the FLP altitude distributions in the two sub-regions. The P-value is smaller than 0.01 at the 95 % confidence level. Therefore, we conclude that the altitude distribution in the western NTS is significantly different from those located in the eastern part. The relatively scattering pattern in the west indicates that the altitude distributions of the ARGs there are not dominantly controlled by the geographical location

(i.e., longitudes). There could be other factors influencing the altitude distributions in the west. For instance, Bolch and Gorbunov (2014) revealed that the characteristics of the contributing area (e.g., the area, slope, and elevation range) of rock glaciers would influence their altitude distribution as well.”

L280ff: You need to consider that the temperature in the blocky material of the rock glaciers can be significantly colder than the MAAT (e.g. Gorbunov et al. 2004).

Authors: Thanks for pointing out this. We have rewritten the relevant sentences (copied below) to declare this (Lines 333–336).

“The highest correlation coefficient is found at the factor “MAAT”. Although the blocky material of the rock glaciers may lower the ground temperature from the air temperature, this high correction implies that the MAAT may influence ground thermal conditions and thus the formation, evolution, and survival of the ARGs.”

L287ff: The first lines of this section describe methods and should be presented in the methods section. In addition, as mentioned information about the time of the year when the velocity was measured are required.

Authors: We move the first sentence into the method section (Line 202). The times of the measured velocities are provided in Table 1.

L302f: The presence of water (e.g. from snow melt or heavy rain fall) has a strong influence on the short term variation. This should be mentioned and discussed along with the acquisition period of the data.

Authors: We now state the influences of precipitations on the derived ARGs surface velocities (Lines 365–367).

5. Discussion

I suggest a separate discussion section where you put your results into context. I suggest to slightly shorten the discussion about the comparison to the existing permafrost maps.

Authors: As suggested, we have shortened the discussion on the comparison of our results to the IPA and CAS permafrost maps.

6. Conclusions

Readers often read the abstract and look at the figures conclusion only before they decide to read the entire paper. I would therefore not use non-common abbreviations in the conclusions and figure captions.

Authors: We have removed the non-common abbreviations in the conclusions and figure captions, such as ARGs and NTS.

L434: Use one decimal digit only. The exact area is quite uncertain.

Authors: We have changed the area of ARGs with one decimal digit.

L442ff: Conclusion 4 is hard to understand.

Authors: Considering the comment of the first referee, we removed the Conclusion 4.

L446: The methods cannot only applied in China.

Authors: We have stated the method can be applied to the other high mountains globally (Line 532).

L450: Reformulate the last sentence. Rock glaciers provide a hint for permafrost occurrence but should be used with care when modelling permafrost distribution.

Authors: We have rewritten the sentence to be rigorous, please see Lines 532–533.

Mapping and inventorying active rock glaciers in the Northern Tien Shan of China using satellite SAR interferometry

Xiaowen Wang^{1,2}, Lin Liu¹, Lin Zhao³, Tonghua Wu³, Zhongqin Li³, Guoxiang Liu²

¹ Earth System Science Programme, Faculty of Science, The Chinese University of Hong Kong, Hong Kong, China

² Department of Remote Sensing and Geospatial Information Engineering, Southwest Jiaotong University, Chengdu, China

³ Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou, China

Correspondence to: Lin Liu (liulin@cuhk.edu.hk)

Abstracts: Rock glaciers are widespread in the Tien Shan. However, rock glaciers in the Chinese part of the Tien Shan have not been systematically investigated for more than two decades. In this study, we propose a new method that combines SAR interferometry and optical images from Google Earth to map active rock glaciers (ARGs) in the Northern Tien Shan (NTS) of China. We compiled an inventory that includes 261 ARGs and quantitative information about their locations, geomorphic parameters, and down-slope velocities. Our inventory shows that most of the ARGs are moraine-derived (69 %) and facing north-east (56 %). The altitude distribution of ARGs in the western NTS is significantly different from those located in the eastern part. The down-slope velocities of the ARGs vary significantly in space, with a maximum of about 114 cm yr⁻¹ and a mean of about 37 cm yr⁻¹. Using the ARG locations as a proxy for the extent of alpine permafrost, our inventory suggests that the lowest altitudinal limit for the presence of permafrost in the NTS is about 2500–2800 m, a range determined by the lowest ARG in the entire inventory and by a statistics-based estimation. The successful application of the proposed method would facilitate effective and robust efforts to map rock glaciers over mountain ranges globally. And this study provides an important dataset to improve mapping and modeling permafrost occurrence in vast western China.

Keywords: Rock Glacier, Tien Shan, InSAR, Mountain Permafrost, Remote Sensing

25 **1 Introduction**

Rock glaciers are lobate or tongue-shaped landforms consisting of mixtures of unconsolidated rock debris and ice in an alpine environment. They are commonly regarded as visible expressions of permafrost presence in periglacial areas (Barsch 1996; Humlum, 2000; Haeberli et al., 2006; Brenning et al., 2007; Boeckli et al., 2012). Rock glaciers can take several thousand years to develop, so a knowledge of their
30 distribution can provide valuable information about past incidences of permafrost and related climatological characteristics (Humlum, 1998; Konrad et al., 1999; Millar et al., 2015; Sorg et al., 2015). Additionally, due to the potential melting of internal ice in the runoff season, rock glaciers serve as important freshwater reserves for the downstream water use in semi-arid regions (Brenning, 2005; Janke et al., 2015).

According to their kinematic status and the presence of internal ice, rock glaciers can be classified into three
35 groups: active, inactive, and relict. Active rock glaciers (ARGs) typically creep downslope cohesively under gravity by a few centimeters to a few meters per year (Berger et al., 2004; Kääb et al., 2003; Necsoiu et al., 2016). The downslope movement of ARGs may cause mass waste hazards on steep slopes, threatening the sensible infrastructures in some downhill areas (e.g., Schoeneich et al., 2015).

Vertically distributed within a narrow band just above the lower limit of permafrost, active and inactive rock
40 glaciers (collectively termed as “intact” rock glaciers) are indicators of alpine permafrost. Many studies have used active and inactive rock glaciers to map local-scale indicators of permafrost occurrence in major mountain ranges such as the European Alps, the Rocky Mountains, the Andes, and the Hindu-Kush-Himalayan region (e.g., Barsch 1996; Janke 2005; Brenning and Trombotto, 2006; Boeckli et al., 2012; Bollmann et al., 2015; Schmid et al., 2015). However, it is difficult to assess the kinematic status of rock
45 glaciers directly (either from the field or remote sensing) and therefore challenging to distinguish active and inactive rock glaciers (Onaca et al., 2016; Sattler et al., 2016). If one only uses ARGs, it would be more accurate and robust to infer the permafrost distribution (Sattler et al., 2016).

It is frequent to use geomorphological features to identify ARGs. Rock glacier movements commonly
50 manifest in viscous flow morphologies on the ground surface, such as the presence and development of ridges, furrows, and steep fronts (Clark et al., 1998; Ikeda and Matsuoka, 2006; Berthling, 2011; Janke et al., 2015). Based on the indications of these geomorphological features, one can map ARGs by visual

interpretation through field surveying or aerial/satellite images (Angillieri, 2009; Brenning et al., 2012; Lilleøren et al., 2013; Scotti et al., 2013; Schmid et al., 2015). However, field surveys are usually scarce because most rock glaciers are located in the remote areas and are difficult to access. Because optical images
55 are often affected by clouds and shadows, it is not always feasible to use them to distinguish ARGs from inactive and relict rock glaciers.

In addition to visually interpreting geomorphological features, measuring surface displacements can also be used to identify active rock glaciers. Various terrestrial geodetic techniques such as differential GPS and laser scanning have been used (Lambiel and Delaloye, 2004; Avian et al., 2009; Bollmann et al., 2015).
60 However, these terrestrial techniques are expensive and only offer limited spatial coverage. They have therefore mostly been used in detailed case studies. Multi-temporal optical images have also been used to investigate surface displacements of ARGs through visual interpretation and feature tracking (Gorbunov et al., 1992; Kääb et al. 1997).

In recent years, satellite synthetic aperture radar interferometry (or InSAR) has emerged as a unique
65 technique to investigate the dynamics of rock glaciers (Kenyi and Kaufmann, 2003; Strozzi et al., 2004; Lilleøren et al., 2013; Liu et al., 2013; Barboux et al., 2015; Necsoiu et al., 2016). Compared with optical remote sensing techniques, InSAR provides an all-weather imaging capability for measuring ground surface deformation with centimeter to sub-centimeter precision. Thus, the InSAR measurements can be used to determine the kinematic status of rock glaciers (especially their velocities, the upper and lower boundaries),
70 which complement the limitations of optical images (Kenyi and Kaufmann, 2003; Liu et al., 2013). However, the combined use of InSAR measurements and optical images for mapping ARGs in a vast, regional area is still rare. In this study, we proposed an integrative method to identify and map ARGs with improved accuracy, effectiveness, and efficiency by (1) detecting moving landforms using InSAR and (2) interpreting geomorphic features from optical images in Google Earth.

75 Rock glaciers are abundant in the high mountains of Central Asia and western China, such as the Tien Shan, Hindu Kush Himalayan, Kunlun Shan and Hengduan Shan. More than 1500 rock glaciers have been inventoried in the Kazakh and Kyrgyz Tien Shan (Gorbunov et al., 1992;1998). Schmid et al. (2015) compiled an inventory with 702 rock glaciers identified from Google Earth images in the vast Hindu Kush Himalayan region. Using these inventories, recent studies have further discussed the rock glaciers-

80 permafrost interactions (Bolch and Gorbunov, 2014; Schmid et al., 2015). However, rock glaciers in western China have not been systematically investigated except for a small number of studies conducted more than two decades ago. Based on the interpretation of aerial photos and field surveys, Cui and Zhu (1989) and Zhu (1992a) identified hundreds of rock glaciers in the NTS of China, while Cui (1984) and Li and Yao (1987) identified dozens of them in the Kunlun Shan and the Hengduan Shan, respectively.

85 The previous studies of rock glaciers in western China mainly focused on their shapes, classification, and topo-climatic conditions. The studies on surface velocities of rock glaciers are very limited except for the field surveys at several specific sites conducted by Cui and Zhu (1989) and Zhu (1992a). Additionally, there has not been any discussion on the permafrost distribution for these high mountains in western China based on locations of rock glaciers. In this study, we applied the new integrative method to identify and map the ARGs in the Northern Tien Shan (NTS) region in western China as an example study area. We determined the geographical locations of ARGs based on the proposed method. Then, we systematically summarized their geomorphic and climatic attributes, as well as their surface velocities. Finally, we inferred the lower limit of permafrost in the NTS from the inventoried ARGs, offering a modern, comprehensive, and rigorous update from the previous field-based estimates by Qiu et al. (1983). We also compared our inventory with the previous studies and existing permafrost maps.

2 Study area

The Tien Shan is a major mountain system in Central Asia, with a mean altitude of more than 4000 m above sea level. The Tien Shan extends 2500 km by 250–300 km, spanning regions of Uzbekistan, Kyrgyzstan, southeastern Kazakhstan, and the Xinjiang province of China. The Chinese portion of Tien Shan divides Xinjiang into two large basins, the Tarim Basin in the south and the Junggar Basin in the north, with the Gurbantunggut and the Taklamakan Deserts lying in hinterland basins (Fig.1). Due to its geographical location and rugged topography, Tien Shan possesses the ideal climatic conditions for the creation and preservation of mountain permafrost. Qiu et al. (1983) estimated that permafrost in the Chinese part of the Tien Shan covers approximately 6.3×10^4 km². In this study, we focus on the Northern Tien Shan (NTS) of China (82°–87° E, 42°–44° N), a large branch of the Tien Shan striking N20°W from northwest to southeast (Fig. 1). We note that some literature also refers to the part of Tien Shan in Kazakhstan and Kyrgyzstan as

“Northern Tien Shan”. If not explicitly stated, ‘NTS’ refers to our study area in China in the remainder of this paper.

110 The eastern and western parts of the NTS divide along longitude 86° E, and possess distinct climatic characteristics (Zhu et al., 1992b; Zhao et al., 2010). The west is relatively humid and cold. In contrast, the east is dry and cold. There are two long-term meteorological stations in the NTS. The Daxigou meteorological station (86.84° E, 43.11° N) in the eastern part is at about 3540 m above sea level. The Snow-cover and Avalanche Research Station (SARS, 84.40° E, 43.26° N) lies in a valley in the western part of the NTS, at an elevation of about 1776 m. The mean annual air temperature (MAAT) and the mean 115 annual precipitation around the Daxigou station were about -6 °C and 405 mm in 1990–2008, respectively (Sun et al., 2013). The MAAT near the SARS was about 1.3 °C, falling to -9.4 °C at 3580 m. The mean annual precipitation at the SARS was about 830 mm (Shi et al., 2009). The east-west contrast is also evident in the number and size of mountain glaciers. The west hosts more and larger glaciers than the east.

3 Methodology

120 We identified ARGs and measured their surface velocities by using InSAR and optical images. In this section, we first present the details of the InSAR processing for detecting the moving targets in the interferogram and the rules used to identify ARGs in Google Earth by visually interpreting their geomorphic features (Section 3.1). We then describe the method to derive the down-slope velocities of the ARGs (Section 3.2). Finally, we list and explain the criteria for compiling an inventory that contains the 125 geomorphic and dynamic attributes of the identified ARGs, followed by a description of the uncertainty analysis (Section 3.3).

3.1 Identifying and mapping active rock glaciers

3.1.1 Identifying ground movements using SAR interferometry

130 The InSAR technique detects the ground movements by exploiting the phase difference between SAR images acquired at different times (Rosen et al., 2000). The phases in the differential interferogram are wrapped within $-\pi$ to π . One phase cycle corresponds to one-half wavelength (e.g., 11.8 cm for L-band) of ground displacements along the line-of-sight direction (Rosen et al., 2000; Wang et al., 2015). We used

135 wrapped interferograms rather than unwrapped displacement maps to detect ARGs because wrapped phases in the interferogram show apparent variations visually even if the ground movements are small, for instance, at centimeter level.

We used seven orbital paths of L-band ALOS PALSAR images acquired from 2007 to 2009 to cover the entire study area. All images were collected at a radar off-nadir angle of 38.7° in HH polarization and have a pixel spacing of 4.7 m in slant range and 3.2 m in azimuth. For each orbital path, we used one image pair to form one interferogram. All the PALSAR images we used, except for the Path 503, were acquired in summer seasons. To achieve high interferometric coherence, we selected the images pairs with temporal spans of either 46 or 92 days and critical perpendicular baselines smaller than 600 m for further processing. Table 1 supplies detailed information on the interferometric pairs. With the availability of PALSAR images, we implemented the interferometric processing by using GAMMA software (Wegmüller et al. 1998). The topographic phases were estimated and removed by using the 1-arc-second SRTM digital elevation model data with a spatial resolution of about 30 m. We applied multi-looking operation (2 looks in range and 5 looks in azimuth) and filtered interferometric phase using the adaptive Goldstein filter with a window size of 8 by 8 pixels. Finally, we obtained the interferogram maps with a spatial resolution of about 15 m.

150 However, we were unable to map the ARGs solely based on the phase variations in the interferograms. First, the phase variations can be caused by several kinds of error sources in SAR interferograms such as decorrelation noise, residual topography contribution, and atmospheric delay (Liu et al., 2013; Strozzi et al., 2004). The contribution of the topographic error to the interferometric phase is proportional to the perpendicular baseline, the radar slant range, and the radar look angle (Rosen et al., 2000). The maximum perpendicular baseline of the interferograms produced in this study is about 600 m. Adopting the vertical accuracy of the digital elevation model as its nominal value of 16 m (Farr et al., 2004), we estimate that the residual topographic phase in the InSAR measurements would be about 0.92 radians, corresponding to 1.7 cm. The atmospheric (including tropospheric and ionospheric) effects are generally long wavelength (1–10 km) signals (Hanssen, 2001), thus can be assumed nearly constant over a specific rock glacier. By using a local reference point just outside each rock glacier (see Section 3.2), we effectively removed these large-scale atmospheric errors. Second, the moving targets may correspond to other periglacial landforms such as debris flow, solifluction, and protalus lobes (Scotti et al., 2013). Therefore, we decided to overlay the

interferograms on optical satellite images in Google Earth to identify the ARGs by visually interpreting their morphological features.

3.1.2 Identifying active rock glaciers using Google Earth images

165 The Google Earth images covering our study area were acquired by the SPOT satellite with spatial resolution generally smaller than 5 m (Visser et al., 2014). We selected the images that were cloud free and taken in summer seasons with limited snow cover, which facilitates our visual identification. ARGs are characterized by distinct flow features and structural patterns. Transversal or longitudinal flow features (ridges and furrows) are common on ARGs due to the deformation of internal ice (Clark et al., 1998; Humlum, 2000; Haeberli et al., 2006; Berthling, 2011). Many ARGs also have structural patterns such as the steep frontal slopes and side slopes with swollen bodies. Due to the constant supply of talus or debris, the surface textures of ARGs are usually different from the surrounding slopes, and their surface slopes usually have little or no vegetation. Based on these criteria, we visually examined the landforms in the Google Earth image that correspond to the moving targets in the interferograms and identified the ARGs. We marked the geographical location of each identified ARG and delineated its outline using Google Earth. We distinguished ARGs from debris-covered glaciers based on the de-correlation conditions of interferograms and their different visual features on the Google Earth images. Compared with rock glaciers, debris-covered glaciers generally move much faster (Janke et al., 2015), which results in large areas of de-correlation in our PALSAR interferograms. The surface of a debris-covered glacier is usually characterized by ice cliffs and supra-glacial lakes (Bolch et al., 2007). And the rooting zones of debris-covered glaciers are continuous with clean glacier ice (Davies et al., 2013; Lukas et al., 2007).

180 Based on the sources of rock materials that input into the surfaces and internal bodies of rock glaciers, we further classified the inventoried ARGs into two categories: moraine-derived active rock glaciers (MARGs) and talus-derived active rock glaciers (TARGs). MARGs develop beneath the end moraines of glaciers and transports mainly reworked glaciogenic materials (Lilleøren et al., 2013; Scotti et al., 2013). TARGs develop below talus slopes in which there is no visible ice upslope of the landform body, and hence transport mainly rock fragments generated from the adjacent rock walls (Lilleøren and Etzelmüller, 2011). For example, Figs. 2a and 2b show a typical MARG and TARG, respectively. In addition, we found some rock glaciers merge

to form a complex, and we classified the sub-branches as different rock glaciers in our inventory based on their different aspect angles.

190 We demonstrate the method described above using the identified ARGs in the zone “Z” (see its location in Fig. 1) which has an area of about 90 km². Fig. 3a shows the wrapped interferometric phase for this zone. The interferogram was made using images taken 46 days apart in 2007, on September 2 and October 18, respectively. This interferogram has good coherence with a mean value of 0.82, which makes it easy to identify the moving targets based on the phase variations. In the zone “Z”, we found ten moving targets. We
195 then visually checked the surface geomorphic features of each of the moving targets on the Google Earth images (Fig. 3b) and identified seven of ten as ARGs. For the other three moving targets, two are artifacts due to phase errors, and the other one is an alluvial fan other than rock glaciers, we grouped all these three as none rock glaciers. For a closer inspection, we zoom into the rock glacier in the northwest corner in Fig. 3a. Fig. 3c shows that outline determined from the variations of interferometric phases. Fig. 3d shows the
200 corresponding Google Earth image.

3.2 Deriving the surface velocity

We further estimated ground displacements using the InSAR measurements. We unwrapped the interferograms using the minimum cost flow method by selecting a reference point in the flat area with high coherence. For any ARG located in the overlapping regions of two adjacent paths, we selected the
205 interferogram with a higher coherence on the ARG to derive the surface velocity. Previous studies indicated that phase unwrapping is a major limiting factor to the use of SAR interferometry for monitoring active rock glaciers (Barboux et al., 2015). Phase unwrapping may fail in the areas where phase gradients are large due to fast slope movements. To minimize the phase unwrapping errors, we applied the phase filtering with a
window of 8×8 pixels and masked out the de-correlated areas with a coherence threshold of 0.3 before phase
210 unwrapping. We chose a local reference patch (3×3 pixels) near but outside each ARG to calibrate the unwrapped phase. We then calculated the surface velocities of the ARGs by dividing the displacements by the temporal span of the interferogram. We then converted the InSAR-estimated velocities in the line-of-sight direction (V_{LOS}) into the down-slope direction (V_{slp}) using the following equation (Liu et al., 2013):

$$V_{slp} = \frac{V_{LOS}}{\sin(\alpha-\beta) \sin \theta_{inc} \cos \theta_{slp} + \cos \theta_{inc} \sin \theta_{slp}} \quad (1)$$

215 where α is the flight direction of the SAR satellite measured from the north; θ_{inc} is the local incidence angle which can be calculated based on the SAR looking geometry and local topography (i.e., digital elevation model) data; β and θ_{slp} are the aspect angle and slope angle of the ARG, respectively. Finally, we calculated the mean value of all pixels within each ARG to represent its overall velocity.

3.3 Characterizing the geomorphic and climatic parameters of active rock glaciers

220 To characterize the geomorphic features of the ARGs, we first determined the initial line point (ILP) and front line point (FLP) of each ARG, respectively. According to the definition given by Humlum (1998) and Sattler et al. (2016), the ILP is usually located in the rooting zone of a rock glacier and represents the place where permafrost creep starts. Given that the interferometric phase variations represent ground movements, we can outline an ARG by identifying where interferometric phase starts to vary from the local background. We interpreted the boundary in the rooting zone of an ARG as the initial line where the permafrost starts to creep. If InSAR coherence was low in the rooting zone, we used morphologic features such as ridges and furrows instead to determine the initial line. We then used the central point of the initial line as the ILP for an ARG. The FLP, the lowest place where a rock glacier can reach, could be easily identified from InSAR and Google Earth images. Fig. 4 gives an example showing the determined ILP and FLP for a typical TARG.

We also quantified the geomorphic (including altitude, area, length, shape, slope angle, and aspect angle) and climatic parameters (Mean Annual Air Temperature, MAAT and potential income of solar radiation, PISR) of each ARG. We determined the altitudes of the ILP and FLP from the SRTM digital elevation model. We then calculated the slope distance (L), mean slope angle (θ_{slp}) and the mean aspect angle (β) based on the coordinates of ILP and FLP. Dividing the area by the square of length gives the aspect ratio, which distinguishes tongue-shaped (aspect ratio <1) from lobate-shaped (aspect ratio >1) ARGs. We extracted the MAAT at the FLP from the NCAR-NCEP global climate reanalysis and interpolated station measurements released by Gruber (2012). We calculated the PISR over one year (units: W/m^2) using the method described by Kumar et al. (1997) with the assistance of the digital elevation model.

240 With some assumptions, we performed a simple error analysis about the rock glacier parameters we estimated. First, with the high-resolution interferogram maps and Google Earth images, we estimated that the location of each ARG and FLP/ILP we determined is accurate within 50 m (i.e., three times of the

InSAR spatial resolution). The uncertainties of the FLP and ALP altitudes are 16 m, the nominal vertical accuracy of the SRTM digital elevation model. The uncertainties of the other geometric parameters can be estimated by error propagation. For instance, given a rock glacier with a length of 1000 m, an aspect angle of 80°, and a slope angle of 12°, we estimated that the uncertainties for length as 67.2 m, aspect as 3.8 °, and slope as 1.6 °. We calculated that the mean coherence of each ARG and consequently estimated the uncertainty of the mean InSAR measurements based on the empirical method of Hanssen (2001). We then estimated the uncertainties of surface velocities by using the Equation (1). The uncertainties of other geometric parameters (i.e., area and aspect ratio) are difficult to quantify due to the irregular shapes of the ARGs.

Finally, we compiled a spreadsheet to summarize the characteristics of the inventoried ARGs (see the supplementary files). Our inventory lists the geographical locations (including longitude, latitude, and altitude), the geomorphic attributes (including area, length, aspect ratio, slope angle, and aspect angle), the climatic factors (i.e., PISR and MAAT), and the surface velocity and its associated uncertainty of each rock glacier.

4 Results

We identified and mapped a total of 261 ARGs across the NTS (see Fig.1). Table 1 lists the number of ARGs identified in each satellite path. The number of ARGs we compiled is a conservative estimation of all the ARGs in the NTS due to the following three reasons. First, we may have missed some ARGs due to the geometric distortions such as shadows and layovers in the SAR images. For the PALSAR images used in this study, the rock glaciers facing east with the slope angles larger than 51.3° are in the radar shadows, and the rock glaciers facing west with slope angles larger than 38.7° are in the layover regions. These distortions would result in significantly phase de-correlation and make it difficult to identify rock glaciers in these regions. Second, ARGs facing nearly north or south might also have been missed, as InSAR (using images from all space missions, including ALOS) is not sensitive to ground motions along these directions. Finally, some small ARGs could not be identified as the interferogram maps have a moderate resolution of about 15 m.

In the following subsections, we first present some general statistics of the inventory (Section 4.1). We then
270 characterize the altitude distribution of the ARGs in Section 4.2 and describe of the surface velocities of the
ARGs in Section 4.3.

4.1 General statistics of the inventory

Table 2 provides the overall information of the inventoried ARGs in the NTS. Among the 261 ARGs we
mapped, 179 (69 %) are MARGs and 82 (31 %) are TARGs. The MARGs are generally longer, steeper, and
275 larger than the TARGs. The mean FLP altitudes for three groups, namely, all the ARGs, the MARGs, and
the TARGs, are 3174 m, 3208 m, and 3099 m, respectively. Their mean ILP altitudes are 3486 m, 3516 m,
and 3420 m, respectively. These figures indicate that the favorable altitude for ARGs in the NTS is above
3000 m, which is consistent with the previous reports of Liu et al. (1995) and Zhu et al. (1992b). Most of the
ARGs are tongue-shaped, with a mean aspect ratio of about 0.25. The mean area of the MARGs is nearly
280 twice as large as that of the TARGs. The ARGs cover a total area of about 91.5 km², 85 % of which is
occupied by MARGs.

Taking all ARGs as a single group, their ILP altitudes and FLP altitudes both show nearly norm distribution
(Figs. 5a and 5d). The majority of ARGs have the ILP altitudes higher than about 3300 m (Fig. 5a) and the
FLP altitudes higher than about 3000 m (Fig. 5d). ILPAs and FLP altitudes show different statistic
285 distributions between the MARGs and TARGs. The histogram of ILP altitude for the MARGs shows a sharp
increase at about 3300 m and a sharp decrease at about 3800 m (Fig. 5b). In contrast, the histogram of ILP
altitude for the TARGs shows a nearly even distribution, though with some degree of randomness (Fig. 5c).
The FLP altitudes for MARGs and TARGs both show nearly norm distributions (Figs. 5e–f). The minimum,
median, and maximum FLP altitudes of the MARGs are all higher than those of the TARGs, indicating that
290 the MARGs develop on higher slopes than the TARGs.

In the inventory, 70 % of the ARGs (number: 182) are located on north-facing (defined between NW and
NE) slopes (Fig. 6a), with the NE aspect predominating (56 %), indicating the north-facing slopes are the
most suitable slopes for rock glacier development and formation in the NTS. The number of north-facing
MARGs and TARGs are 134 and 48, respectively, accounting for the 75 % and 59 % of the total numbers of
295 MARGs and TARGs, respectively. The difference is probably because the aspect distribution of the
glaciers, which influence the aspect of MARGs as the evolution of MARGs are controlled by the input

glacier moraines. We also noted that the majority of glaciers in the NTS are north-facing, controlled by topography, climate, and geometric factors (Aizen et al., 1995; Li and Li, 2014). Compared with the MARGs, the TARGs are more evenly distributed across all aspects. This pattern is consistent with that found in rock glaciers in the central Italian Alps and Southern Carpathians (Scotti et al., 2013; Onaca et al., 2016).

The subsurface thermal condition is another major factor controlling the development of ARGs, which is influenced by solar radiation. Fig. 6b shows the variations of PISR with the aspect angles. The mean value of PISR for the north-facing and south-facing slopes are $8.83 \times 10^5 \text{ W/m}^2$ and $10.46 \times 10^5 \text{ W/m}^2$, respectively. The south-facing ARGs typically receive higher PISR than their north-facing counterparts. As most of the ARGs are north-facing, we conclude that slopes with lower PISR favor the presence of ARGs. This is probably because lower PISR helps to preserve ground ice inside ARGs.

4.2 Altitudinal distribution of the active rock glaciers

To explore how the factors would impact the altitude distribution of the ARGs, we calculated the Pearson correlation coefficients between the FLP altitudes and the environmental factors listed in Section 3.3. FLP altitude denotes the lowest altitude that an ARG can reach and is frequently used as the lower boundary for the presence of mountain permafrost locally (Lilleøren et al., 2013; Schmid et al., 2015). For the ARGs in our inventory, the correlation coefficients between the FLP altitudes and the longitude, latitude, MAAT, PISR, slope angle, and aspect angle are 0.47, -0.51, -0.86, 0.29, -0.35, and -0.08, respectively.

The simple correlation analysis confirms the strong influence of the geographical locations on the altitude distribution of the ARGs (Fig. 7a and 7b). The FLP altitudes generally increase from west to east and decrease from south to north. This spatial pattern could be mainly because the NTS lays with a strike of about $N20^\circ W$ extending from northwest to southeast. Along this direction, the air temperature increases with lower latitude, thus increasing the lowest altitude for the development of rock glaciers. Moreover, glaciers would also influence the altitude distribution of ARGs as MARGs generally develop beneath the end moraines of glaciers. The western part of the NTS is relatively more humid. Thus, the larger and longer glaciers would develop and cause the MARGs in the west having lower FLP altitudes than the eastern ones.

325 Additionally, we found different altitude distributions of ARGs in the western and eastern parts (Fig. 7a).
The nearly linear pattern in the east is more apparent than that in the west. We performed a T-test on the
similarity of the FLP altitude distributions in the two sub-regions. The P-value is smaller than 0.01 at the
95 % confidence level. Therefore, we conclude that the altitude distribution in the western NTS is
significantly different from those located in the eastern part. The relatively scattering pattern in the west
indicates that the altitude distributions of the ARGs there are not dominantly controlled by the geographical
location (i.e., longitudes). There could be other factors influencing the altitude distributions in the west. For
330 instance, Bolch and Gorbunov (2014) revealed that the characteristics of the contributing area (e.g., the area,
slope, and elevation range) of rock glaciers would influence their altitude distribution as well.

The correlation analysis also reveals the relative importance of various factors to the rock glacier
distributions. The highest correlation coefficient is found at the factor “MAAT”. Although the blocky
material of the rock glaciers may lower the ground temperature from the air temperature, this high correction
implies that the MAAT may influence ground thermal conditions and thus the formation, evolution, and
335 survival of the ARGs. A moderate positive correlation is observed at the factor “PISR”. Receiving a higher
PISR, the ground temperature is likely higher and the internal ice is more likely to melt. The correlation
between the slope angles and FLP altitudes are associated with the favorable presence of TARGs on the
steep slopes. We found no obvious correlation between the aspect angles and FLP altitudes.

340 **4.3 Surface velocities of the active rock glaciers**

We were conservative in conducting statistical analyses on the surface velocities by only including the
ARGs with a mean coherence higher than 0.3 and a mean velocity larger than 5 cm yr⁻¹. Finally, the surface
velocities of 170 ARGs (including 111 MARGs and 59 TARGs) were documented (Fig. 8). The surface
velocities of the 170 ARGs show a spatially heterogeneous pattern over the NTS. Most of ARGs (84 %)
345 have surface velocities lower than 60 cm yr⁻¹. The maximum surface velocities in the eastern and western
parts are about 89 and 114 cm yr⁻¹, respectively. The mean velocity and the standard deviation of all ARGs
are 37 cm yr⁻¹ and 21 cm yr⁻¹, respectively. The maximum velocity uncertainty is 28.8 cm yr⁻¹, and the mean
uncertainty is 7.5 cm yr⁻¹.

We selected two pairs of ARGs to inspect the spatial variations of surface velocities closely (Figs. 8a–b).
350 The two ARGs in each pair are adjacent to each other. Therefore, we assumed that they developed in similar

topo-climatic environments (e.g., similar altitude, MAAT, and precipitation). The surface velocities of ‘ARG1’ show a spatial homogeneity pattern while these of the other three all show a spatially heterogeneous pattern. The fastest moving zones are located near the center, the rooting zone, and the front of ‘ARG2’, ‘ARG3’, and ‘ARG4’, respectively. This significant heterogeneity of surface velocities within ARGs indicates that the kinematics of surface movements are complex. Previous studies have demonstrated that the surface velocities ARGs are not only influenced by topo-climatic factors but also the ice condition and internal structure of ARGs (Haeberli et al., 2006; Serrano et al., 2010; Schoeneich et al., 2015). However, it is difficult to determine and quantify the ground ice conditions or ground thermal conditions of the ARGs in this study. Nevertheless, the spatial variabilities of velocities at both the local and regional scales suggest that the dynamics of ARGs are not controlled by any single factor, but a combination of many factors.

In addition, the SAR images we used were acquired at different times, which may also contribute to the spatial heterogeneity of the surface velocities. We used seven pairs of PALSAR images, one from each path, to cover the study area. However, the acquisition times for each image pairs are different (Table 1). Recent studies show that the surface velocity of rock glacier may vary seasonally (Liu et al., 2013; Wirz et al., 2016). Moreover, strong precipitations or snowmelt may also influence the surface velocities. In this study, we simply chose these PALSAR images for the inventory exercise and ignored the temporal dynamics of rock glaciers.

4.4 Lower limit of permafrost occurrence in the Northern Tien Shan

We used the inventoried ARGs as a proxy of permafrost to infer lower limit of permafrost occurrence in the NTS. The previous estimates of the lower limit of permafrost presence in the NTS were mainly based on field surveys near the Urumqi River source region (87.00° E, 43.07° N) in the east and near the Haxilegen Pass (84.45°E, 43.77° N) region in the west (Qiu et al., 1983; Jin et al., 1993; Qiu, 1993). However, these site-based estimations may not be representative of the whole study area. Therefore, the ARGs we inventoried are valuable for re-estimating the permafrost distribution for this region. Permafrost is unlikely to develop below the FLP, and the proportion and extent of permafrost usually increase towards higher altitudes. Because our inventory shows that MARGs generally develop on higher slopes than TARGs (see Section 4.1), here we only used the FLP altitudes of TARGs to estimate the lower limit of permafrost.

We grouped the inventoried TARGs into different categories based on their locations (i.e., east and west of 86°E) and aspect angles (i.e., north-facing and south-facing) to infer the lower limit of permafrost under different environmental conditions. The first row of Table 3 summarizes their altitudinal limits: 2490 m and 2583 m for the north-facing and south-facing slopes in the west, respectively; and 2895 m and 3088 m for the north-facing and south-facing slopes in the east, respectively. Overall, the altitudes of the inventoried ARGs increase from west to east as shown in Fig. 7a.

However, these estimates may not give an entirely accurate picture of the permafrost limit for the whole study area, because they are based on a few specific ARGs that have the lowest FLP altitudes in the inventory. For example, the TARG with the lowest FLP altitude (i.e., 2490 m) is located on a north-facing slope in the western part of the NTS (shown in Fig. 9c, outlined by the blue polygon). We cannot rule out the possibility that this particular ARG may be a local anomaly and therefore cannot fully represent the permafrost presence in the broader NTS. Thus, we adopted an alternative method in a statistical sense, which approximates the lower limit of permafrost by subtracting the standard deviation of all FLP altitudes in the inventory from their mean values. The second row of Table 3 lists these statistics-based estimates: 2772 m and 2828 m for the north-facing and south-facing slopes in the west, respectively; and 3005 m and 3041 m for the north-facing and south-facing slopes in the east, respectively. The largest difference between the statistics-based and lowest-altitude-based is 282 m, indicating that both estimates are consistent, although the statistics-based values are normally higher. Therefore, we suggest the lower limit for the permafrost distribution in the NTS is about 2500–2800 m.

The third row of Table 3 summarizes the field-based estimates of the lowest altitude of permafrost given by Qiu et al. (1983). We note that the results of the field surveys are all higher than these inferred from our lowest-altitude-based estimates. We also note that the field-based results are all lower (and higher) than our statistics-based estimates for the north-facing (and south-facing) slopes. The largest differences between the lowest-altitude-based and the field-based estimates are 467 m and 162 m for the western and eastern parts, respectively. The largest discrepancies between the statistics-based and field-based estimates are 222 m and 209 m, respectively. One possible reason for these discrepancies is that our results based on ARG inventory give regional estimates, while the field surveys give site-specific estimates. Additionally, the ARGs move downslope and the fronts of those low-lying ARGs may reach places free of permafrost. Furthermore, the

permafrost conditions in the NTS may have changed in the past two decades. A recent study shows strong permafrost degradation in the Urumqi River Source region since 1992 (Zhao et al., 2010).

5 Discussion

5.1 Comparisons with the previous studies of rock glaciers in the Tien Shan

410 Previous studies of rock glaciers in the NTS are limited with most of them concentrating in the Urumqi River source region. Interpreting aerial photos acquired in the late 1980s and early 1990s, several researchers (Cui and Zhu, 1989; Zhu, 1992a; Zhu et al., 1992b; Liu et al., 1995) identified hundreds of rock glaciers in the NTS. They found that most of the identified rock glaciers are tongue-shaped, located at altitudes between 3300 m and 3900 m, and facing north. These findings are generally consistent with our inventory. These previous studies also revealed a homogeneity pattern of the ARG surface velocities. Cui and Zhu (1989) and Zhu (1992a) measured the movements of the marked boulders on the front slopes of five rock glaciers near the Urumqi River source region, and estimated that their surface velocities varied from 1 cm yr⁻¹ to 75 cm yr⁻¹. Moreover, they found that one rock glacier (referred to as 'RG4' in their study) moved at about 49 cm yr⁻¹ between 1985 and 1986. However, they estimated that its mean velocity was about 6 cm yr⁻¹ from 1960 to 1986. Our measurements show that the largest velocity is located the western part of the NTS, which is also consistent with the findings of Cui and Zhu (1989) and Zhu et al. (1992b). Yet these previous studies did not provide any detailed information such as geographical and geomorphic attributes of the identified rock glaciers. Thus, we cannot compare these field-based estimates with our measurements in further detail.

425 Recent studies of rock glaciers in the Tien Shan have been focusing in two sub-regions: (1) the Djungar Ala Tau in Kazakhstan at about 400 km northwest of our study area and (2) the Ile Alatau and Kungöj Ala-Too mountain ranges (42.5° to 43.5° N, 75° to 79°E) at the border between Kazakhstan and Kyrgyzstan. Gorbunov and Titkov (1992) identified about 850 ARGs in the first region. Comparing their inventory with ours, we find that the rock glaciers in Djungar Ala Tau are at lower altitudes than that in the NTS, which likely results from their more northerly latitude. Gorbunov et al. (1998) inventoried 1034 rock glaciers in the second region, of which at least 851 are identified to be active. Similar to our results, they found that the MARGs occupy a larger area than the TARGs. We also find the altitudinal distributions of the rock glaciers

435 there are similar to that in the NTS. Studies revealed that the annual displacement rates of the ARGs in these regions range from less than 1 m yr⁻¹ to about 11 m yr⁻¹ (Gorbunov and Titkov, 1992; Bolch and Gorbunov, 2014), faster than the ones we mapped in the NTS.

5.2 Limitations of using active rock glaciers to determine the lower limit of permafrost

440 The moderate discrepancies between our estimates and the field surveys of the altitudinal limits for permafrost occurrence in the NTS demonstrated the feasibility of our method to infer the permafrost distribution. However, several factors still exist and limit the accuracy of our method for quantifying the lower limit of permafrost. These factors include (1) the ambiguity about the genesis of rock glaciers, (2) the high heterogeneity of ground thermal regime, and (3) the uncertainties in the use of FLP altitudes of the ARGs to infer permafrost limit.

445 First, it is still debatable to use active rock glaciers as representatives of the exact permafrost boundaries. There are two contradictory viewpoints regarding the genesis of rock glaciers. One holds rock glaciers are permafrost creep features and form in a permafrost environment. The other one holds that rock glaciers form through a continuum of glacial to periglacial processes and thus they may also be related to glaciers in non-permafrost environments (Clark et al., 1998; Berthling, 2011). To avoid this ambiguity, we only used the talus-derived ARGs in this study to infer the lower limit of permafrost in the NTS.

450 Second, as shown in Section 4.2, the altitudinal distributions of the ARGs in the NTS indicate that the presence of rock glaciers is primarily governed by the ground thermal regime. However, the ground thermal regime in the alpine region can be highly heterogeneous in space. In Transili Alatau Tien Shan, the mean annual temperatures inside the coarse debris could be 2.5–4.0 °C cooler than the MAATs, as the blocky debris favors cooler temperatures allowing rock glaciers to exist at elevations where permafrost is otherwise unlikely (Gorbunov et al., 2004). Thus, individual rock glacier only serves as a local indicator of permafrost occurrence. This is why we estimated the lower limit of permafrost occurrence in the regional scale in lowest-altitude-based and statistics-based perspectives. Nevertheless, the statistics show that the mean MAAT at the TARGs is -5.4 °C, and 83 % of the TARGs have the MAAT lower than -4 °C. This indicates that the TARGs we used for estimating the lower limit of permafrost are located in a cold environment, and thus reinforcing the credibility of our estimates.

460 Lastly, several sources of uncertainty exist in the use of FLP altitudes of ARGs to infer the lowest altitude of
permafrost occurrence in the NTS. On one hand, ARGs move downslope and may export permafrost into
lower altitudes that are climatically unfavorable for permafrost formation. Thus, our estimates may give a
bias towards a lower than the true permafrost limit. On the other hand, we did not identify inactive rock
glaciers, which are also indicators of permafrost occurrence. Since inactive rock glaciers generally develop
465 at lower altitudes than active ones (Sollid and Sørbel, 1992), our estimates may bias the permafrost limit
towards a higher value. Additionally, we cannot rule the possibility that some extremely low-lying ARGs are
not included, which may lower our estimated altitude limits. These three kinds of uncertainties justify the
expression of a range of estimations rather than exact values of the altitude limits as given in Section 4.4.

5.3 Comparison with the existing permafrost distribution maps

470 We cross-validated the inferred permafrost distribution by comparing our inventoried ARGs with the
existing permafrost distribution maps. We selected three permafrost maps that covers the NTS region: (1)
the Map of Snow, Ice and Frozen Ground in China released by Cold and Arid Regions Environmental and
Engineering Research Institute, Chinese Academy of Sciences (Shi and Mi, 1988), which we refer to as the
CAS map, (2) the Circum-Arctic Map of Permafrost and Ground Ice Conditions released by the International
475 Permafrost Association (Brown et al., 1998), referred to as the IPA map, and (3) the global permafrost
zonation Index (PZI) map developed by Gruber (2012). The frozen ground in the CAS map is classified into
high-latitude, plateau, and mountain permafrost with a scale of 1:4000000. The IPA map adopts the classes
of continuous, discontinuous, sporadic, and isolated patches of permafrost extent with a scale of 1:1000000.
The PZI map is generated based on the relationships between permafrost extent and long-term MAAT data
480 with a ground resolution of 1 km. The PZI values range from 0 to 1. Values close to 1 represent permafrost
existing in nearly all conditions, while values close to 0 represent permafrost presenting only in the most
favorable conditions. The PZI uncertainty is expressed with “fringe of uncertainty”, referring to a
conservative or an anti-conservative variant than the “normal” scenario that considers the possibility of
<10 % permafrost area as well (Gruber, 2012).

485 Fig. 9 compares the ARGs we inventoried in the NTS with the three permafrost maps described above. The
dotted green and black lines in Fig. 9a outline the mountain permafrost extent from the CAS map and the
discontinuous permafrost from the IPA map, respectively. The background color map in Fig. 9b shows the

PZI values. Most ARGs are located within the permafrost extents of the CAS and IPA maps (256 out of 261 are within the CAS; 249 are within the IPA). We also found that most of the ARGs have high PZI values. As the PZI map has the highest spatial resolution among these three permafrost maps, we further compared our inventoried ARGs with the PZI map.

We randomly selected two sub-regions 'Z1' and 'Z3' (see Fig. 9b) in the western and eastern of the NTS, respectively, to examine the distributions of ARGs in the PZI map. It is evident that most of the MARGs are located above the PZI contour line of 0.8 (Figs. 10a and 10c). We selected the zone 'Z2' (see Fig.9b) as another example as it contains both the MARG and TARG that have the lowest FLP altitude in their corresponding category. The MARG developing in the narrow valley (location: 84.2957°E, 44.0021°N, outlined by the red polygon in Fig. 10b) has an FLP altitude of 2575 m. The TARG (location: 84.2883°E, 44.0228°N, outlined by the blue polygon in Fig. 10b) has an FLP altitude of 2490 m. The PZI values at the front points of these two ARGs are still higher than 0.2.

The PZI value at the FLP of an ARG shows how favorable the lowest altitude that the rock glacier reaches would for permafrost to exist. The statistical distribution of PZI at FLPs shows that most ARGs (73 %) are located in the PZI interval of 0.5 to 0.9 (Fig. 11a). The PZI density peaks at about 0.8 for the MARGs and 0.5 for the TARGs (Figs. 11b–c). The PZI density decreases towards the lower PZI values for the MARGs, while decreases towards both ends of the distribution (Fig. 11c) for the TARGs. This difference in PZI distribution is probably because the TARGs are located at relatively lower altitudes than the MARGs. The PZI values at lower altitudes show a larger variability as the air temperature increases. The mean PZI values at the FLPs are 0.64, 0.67, and 0.58 for all ARGs, MARGs, and TARGs, respectively, indicating the spatial distribution of the ARGs agrees well with the PZI map.

6 Conclusions

In this study, we successfully mapped and inventoried 261 active rock glaciers in the Northern Tien Shan of China by combining the use of the SAR interferometry and optical images from Google Earth. Based on the creep boundaries of the active rock glaciers determined by this novel method, we estimated the lower limit of the permafrost presence in the Northern Tien Shan. From the inventory and analyses, we draw the following conclusions:

- 515 (1) Most of the active rock glaciers in the Northern Tien Shan are moraine-derived, tongue-shaped, and
northeast-facing. The moraine-derived active rock glaciers are generally longer, steeper, and larger than
the talus-derived active rock glaciers. The total area of the active rock glaciers is about 91.5 km² and
moraine-derived ones occupy most of this area (85 %).
- 520 (2) The latitude/longitude locations have a strong influence on the altitude distribution the active rock
glaciers. We found the altitudes of active rock glaciers generally increase from west to east and decrease
from south to north. We also found distinct altitude distributions of active rock glaciers in the western
and eastern parts of the Northern Tien Shan (divided by 86°E), probably due to the influence of distinct
climates of these two sub-regions.
- 525 (3) We found that the lower limit of permafrost in the eastern part of the Northern Tien Shan is higher than
in the western part. The lower altitudinal limit for the permafrost distribution throughout the Northern
Tien Shan is about 2500–2800 m.

The study presented here provides the first comprehensive and modern documentation of the characteristics
of the active rock glaciers in the Northern Tien Shan of China. It updates the lower limit of permafrost
occurrence for this vast area. This inventory offers a baseline dataset for the further investigations on
530 permafrost modeling, slope stability, and water resource, etc. The successful application of the proposed
method in the Northern Tien Shan promises that this method can be used to the other high mountains
globally, thus will help to close the gaps in our knowledge of rock glaciers worldwide. This new knowledge
will be useful to infer alpine permafrost occurrence in high mountains.

Data availability

535 The locations and characteristic parameters of the 261 inventoried active rock glaciers for downloading and
visualizing are available in the supplementary files. Both the KMZ file and shapefiles are provided.

List of Acronyms

ALOS: Advanced Land Observing Satellite

ARG: Active Rock Glacier

540 CAS: Chinese Academy of Sciences
FLP: Front Line Point
GPS: Global Positioning System
ILP: Initial Line Point
InSAR: Synthetic Aperture Radar Interferometry
545 IPA: International Permafrost Association
MAAT: Mean Annual Air Temperature
MARG: Moraine-derived Active Rock Glacier
NTS: Northern Tien Shan
PALSAR: The Phased Array type L-band Synthetic Aperture Radar
550 PISR: Potential Income of Solar Radiation
PZI: Permafrost Zonation Index
SRTM: Shuttle Radar Topography Mission
TARG: Talus-derived Active Rock Glacier

Acknowledgements

555 We thank Tobias Bolch, Gengnian Liu, Tazio Strozzi, and the Editor for their constructive and insightful reviews, which have helped to improve the quality of this paper. The ALOS PALSAR data are copyrighted by the Japan Aerospace Exploration Agency and provided by the Alaska Satellite Facility, University of Alaska Fairbanks. The CAS permafrost map is provided by Environmental and Ecological Science Data Center for West China, National Natural Science Foundation of China (<http://westdc.westgis.ac.cn>). The
560 IPA permafrost map is provided by the Frozen Ground Data Center at the National Snow and Ice Data Center (NSIDC), Boulder, Colorado, USA (<http://nsidc.org/data/GGD318/versions/2>). The MAAT and permafrost map zonation data are downloaded from http://www.geo.uzh.ch/microsite/cryodata/pf_global/. Some figures in this paper were plotted using the Generic Mapping Tools (Wessel et al., 2013). Work at The Chinese University of Hong Kong was supported by Hong Kong Research Grants Council Grants
565 CUHK24300414 and CUHK14300815.

References

- Angillieri, M. Y. E.: A preliminary inventory of rock glaciers at 30°S latitude, Cordillera Frontal of San Juan, Argentina, *Quat. Int.*, 195, 151–157, doi:10.1016/j.quaint.2008.06.001, 2009.
- 570 Aizen, V.B., Aizen, E.M. and Melack, J.M.: Climate, snow cover, glaciers, and runoff in the Tien Shan, central Asia, *Journal of the American Water Resources Association*, 31(6), 1113–1129, 1995.
- Avian, M., Kellerer-Pirklbauer, A., and Bauer, A.: LiDAR for monitoring mass movements in permafrost environments at the cirque Hinteres Langtal, Austria, between 2000 and 2008, *Nat. Hazards Earth Syst. Sci.*, 9, 1087–1094, doi:10.5194/nhess-9-1087-2009, 2009.
- 575 Barboux, C., Strozzi, T., Delaloye, R., Wegmüller, U., and Collet, C.: Mapping slope movements in alpine environments using TerraSAR-X interferometric methods, *Int. J. Photogramm. Remote Sens.*, 109, 178–192, doi:10.1016/j.isprsjprs.2015.09.010, 2015.
- Barsch, D.: *Rock Glaciers: Indicators for the Present and Former Geoecology in High Mountain Environments*. Berlin: Springer, 1996.
- Berger, J., Krainer, K., and Mostler, W.: Dynamics of an active rock glacier (Otzal Alps, Austria), *Quat. Res.*, 62, 580 233–242, doi:10.1016/j.yqres.2004.07.002, 2004.
- Berthling, I.: Beyond confusion: rock glaciers as cryo-conditioned landforms, *Geomorphology*, 131, 98–106, doi:10.1016/j.geomorph.2011.05.002, 2011.
- Boeckli, L., Brenning, A., Gruber, S., and Noetzli, J.: Permafrost distribution in the European Alps: calculation and evaluation of an index map and summary statistics, *The Cryosphere*, 6, 807–820, doi:10.5194/tc-6-807-2012, 2012.
- 585 Bolch, T., Buchroithner, M. F., Kunert, A., Kamp, U.: Automated delineation of debris-covered glaciers based on ASTER data. In *Geoinformation in Europe (Proc. of 27th EARSeL Symposium)*, Bozen, Italy, pp: 403-410, 2007.
- Bolch, T., & Gorbunov, A.P. (2014). Characteristics and Origin of Rock Glaciers in Northern Tien Shan (Kazakhstan/Kyrgyzstan). *Permafrost and Periglacial Processes*, 25, 320–332.
- Bollmann, E., Girmair, A., Mitterer, S., Krainer, K., Sailer, R., and Stotter, J.: A rock glacier activity index based on 590 rock glacier thickness changes and displacement rates derived from airborne laser scanning, *Permafr. Periglac. Proc.*, 26, 347–359, doi:10.1002/ppp.1852, 2015.

- Brenning, A.: Geomorphological, hydrological and climatic significance of rock glaciers in the Andes of Central Chile, *Permafr. Periglac. Proc.*, 16, 231–240, doi:10.1002/ppp.528,, 2005.
- 595 Brenning, A., Grasser, M., and Friend, D. A.: Statistical estimation and generalized additive modeling of rock glacier distribution in the San Juan Mountains, Colorado, United States, *J. Geophys. Res-Earth Surface*, 112, doi:10.1029/2006jf000528, 2007.
- Brenning, A., Long, S. L., and Fieguth, P.: Detecting rock glacier flow structures using Gabor filters and IKONOS imagery, *Remote Sens. Environ.*, 125, 227–237, doi:10.1016/j.rse.2012.07.005, 2012.
- 600 Brenning, A. and Trombotto, D.: Logistic regression modeling of rock glacier and glacier distribution: Topographic and climatic controls in the semi-arid Andes, *Geomorphology*, 81, 141–154, doi:10.1016/j.geomorph.2006.04.003, 2006.
- Brown, J., Ferrians, O., Heginbottom, J. A., and Melnikov, E.: *Circum-Arctic Map of Permafrost and Ground-Ice Conditions*, USA Natl. Snow Ice Data Center, Boulder, Colorado, 1998.
- Clark, D. H., Steig, E.J., Potter, N. Jr and Gillespie, A.R.: Genetic variability of rock glaciers, *Geogr. Ann.*, 80A, 175–
605 182, 1998.
- Cui Z. J.: The discovery of the Kunlunshan-type rock glacier and the latest classification of rock glaciers. *Chin. Sci. Bull.*, 810–813, 1984. (in Chinese)
- Cui, Z. J. and Zhu, C.: The structure pattern of temperature and mechanism of the rock glaciers in Urumqi River region in Tien Shan, *Chin. Sci. Bull.*, 134–137, 1989. (in Chinese)
- 610 Davies, B. J., Glasser, N. F., Carrivick, J. L., Hambrey, M. J., Smellie, J. L., and Nyvlt, D.: Landscape evolution and ice-sheet behaviour in a semi-arid polar environment: Geological Society, London, Special Publications, 381, 353–395, doi:10.1144/Sp381.1, 2013.
- Farr, T. G., Rosen, P. A., Caro, E., Crippen, R., and Duren, R.: The Shuttle Radar Topography Mission, *Rev. Geophys.*, 45, RG2004, doi:10.1029/2005RG000183, 2004.
- 615 Gorbunov, A. P., Marchenko, S. S., and Seversky, E. V.: The thermal environment of blocky materials in the mountains of Central Asia. *Permafrost and Periglacial Processes*, 15(1), 95-98, 2004.

- Gorbunov, A. P., Seversky, E. V., Titkov, S. N., Marchenko, S. S. and Popov, M.: Rock glaciers, Zailiysiky Range, Kungei Ranges, Tienshan, Kazakhstan. National Snow and Ice Data Center/World Data Center for Glaciology: Boulder, CO. Digital media, 1998.
- 620 Gorbunov, A.P., Titkov, S.N., Polyakov, V.: Dynamics of the Rock Glaciers of the Northern Tien Shan and the Djungar Alatau, Kazakhstan. *Permafrost and Periglacial Processes*, 3, 29–39, 1992.
- Gruber, S.: Derivation and analysis of a high-resolution estimate of global permafrost zonation, *The Cryosphere*, 6, 221–233, doi:10.5194/tc-6-221-2012, 2012.
- 625 Haeberli, W., Hallet, B., Arenson, L., Elconin, R., Humlum, O., Kääh, A., Kaufmann, V., Ladanyi, B., Matsuoka, N., Springman, S., and Vonder Muhll, D.: Permafrost creep and rock glacier dynamics, *Permafr. Periglac. Proc*, 17, 189–214, doi:10.1002/ppp.561, 2006.
- Hanssen, R. F.: *Radar interferometry: data interpretation and error analysis*, Springer Science & Business Media, 2001.
- Humlum, O.: The climatic significance of rock glaciers, *Permafr. Periglac. Proc*, 9, 375–395, doi:10.1002/(Sici)1099-1530(199810/12)9:4<375::Aid-Ppp301>3.3.Co;2-S, 1998.
- 630 Humlum, O.: The geomorphic significance of rock glaciers: estimates of rock glacier debris volumes and headwall recession rates in West Greenland, *Geomorphology*, 35, 41–67, doi:10.1016/S0169-555x(00)00022-2, 2000.
- Ikeda, A. and Matsuoka, N.: Pebbly versus bouldery rock glaciers: Morphology, structure and processes, *Geomorphology*, 73, 279–296, doi:10.1016/j.geomorph.2005.07.015, 2006.
- 635 Janke, J. R.: The occurrence of alpine permafrost in the Front Range of Colorado, *Geomorphology*, 67, 375–389, doi:10.1016/j.geomorph.2004.11.005, 2005.
- Janke, J. R., Bellisario, A. C., and Ferrando, F. A.: Classification of debris-covered glaciers and rock glaciers in the Andes of central Chile, *Geomorphology*, 241, 98–121, doi:10.1016/j.geomorph.2015.03.034, 2015.
- Jin, H. J., Qiu, G. Q., Zhao, L., Wang, S. J., and Zeng, Z. G.: Thermal Regime of Alpine Permafrost in the Upper Reach of Urumqi River, Tian Shan, *Journal of Glaciology and Geocryology*, 15, 110-114, 1993. (in Chinese)
- 640 Kääh, A., Haeberli, W., and Gudmundsson, G. H.: Analysing the creep of mountain permafrost using high precision aerial photogrammetry: 25 years of monitoring Gruben rock glacier, Swiss Alps, *Permafrost Periglac.*, 8, 409–426, 1997.

- Kääb, A., Kaufmann, V., Ladstadter, R., and Eiken, T.: Rock glacier dynamics: implications from high-resolution measurements of surface velocity fields, In Eighth International Conference on Permafrost, 1, 501–506, 2003.
- 645 Kenyi, L. W. and Kaufmann, V.: Estimation of rock glacier surface deformation using SAR interferometry data, *IEEE Trans. Geosci. Remote Sens.*, 41, 1512–1515, doi:10.1109/Tgrs.2003.811996, 2003.
- Konrad, S. K., Humphrey, N. F., Steig, E. J., Clark, D. H., Potter, N., and Pfeffer, W. T.: Rock glacier dynamics and paleoclimatic implications, *Geology*, 27, 1131–1134, doi:10.1130/0091-7613(1999)027<1131:Rgdapi>2.3.Co;2, 1999.
- Kumar, L., Skidmore, A. K., and Knowles, E.: Modelling topographic variation in solar radiation in a GIS
650 environment, *International Journal of Geographical Information Science*, 11, 475–497, doi:10.1080/136588197242266, 1997.
- Lambiel, C. and Delaloye, R.: Contribution of real-time kinematic GPS in the study of creeping mountain permafrost: Examples from the Western Swiss Alps, *Permafr. Periglac. Proc*, 15, 229–241, doi:10.1002/ppp.496, 2004.
- Li, S. and Yao, H.: Preliminary study of rock glaciers in the Gongga Mt. Area, *Journal of Glaciology and
655 Geocryology*, 9, 55–59, 1987. (in Chinese)
- Li, Y. A. and Li, Y. K.: Topographic and geometric controls on glacier changes in the central Tien Shan, China, since the Little Ice Age, *Ann. Glaciol.*, 55, 177–186, doi:10.3189/2014AoG66A031, 2014.
- Lilleøren, K. S. and Etzelmüller, B.: A Regional inventory of rock glaciers and ice-cored moraines in Norway, *Geografiska Annaler Series a-Physical Geography*, 93a, 175–191, doi:10.1111/j.1468-0459.2011.00430.x, 2011.
- 660 Lilleøren, K. S., Etzelmüller, B., Gartner-Roer, I., Kääb, A., Westermann, S., and Gudmundsson, A.: The distribution, thermal characteristics and dynamics of permafrost in Trollaskagi, Northern Iceland, as inferred from the distribution of rock glaciers and ice-cored moraines, *Permafr. Periglac. Proc*, 24, 322–335, doi:10.1002/ppp.1792, 2013.
- Liu, G. N., Xiong, H. G., Cui, Z. J., and Song, C. Q.: The morphological features and environmental condition of rock glaciers in Tianshan mountains, *Scientia Geographica Sinica*, 15, 226–233, 1995. (in Chinese)
- 665 Liu, L., Millar, C. I., Westfall, R. D., and Zebker, H. A.: Surface motion of active rock glaciers in the Sierra Nevada, California, USA: inventory and a case study using InSAR, *The Cryosphere*, 7, 1109–1119, doi:10.5194/tc-7-1109-2013, 2013.

- 670 Lukas, S., Nicholson, L. I., and Humlum, O.: Comment on Lønne and Lysa (2005): “Deglaciation dynamics following the Little Ice Age on Svalbard: Implications for shaping of landscapes at high latitudes”, *Geomorphology*, 84, 145–149, doi:10.1016/j.geomorph.2006.07.003, 2007.
- Millar, C. I., Westfall, R. D., Evenden, A., Holmquist, J. G., Schmidt-Gengenbach, J., Franklin, R. S., Nachlinger, J., and Delany, D. L.: Potential climatic refugia in semi-arid, temperate mountains: Plant and arthropod assemblages associated with rock glaciers, talus slopes, and their forefield wetlands, Sierra Nevada, California, USA, *Quat. Int.*, 387, 106–121, doi:10.1016/j.quaint.2013.11.003, 2015.
- 675 Necsoiu, M., Onaca, A., Wigginton, S., and Urdea, P.: Rock glacier dynamics in Southern Carpathian Mountains from high-resolution optical and multi-temporal SAR satellite imagery, *Remote Sens. Environ.*, 177, 21–36, doi:10.1016/j.rse.2016.02.025, 2016.
- Onaca, A., Ardelean, F., Urdea, P., and Magori, B.: Southern Carpathian rock glaciers: Inventory, distribution and environmental controlling factors, *Geomorphology*, doi: 10.1016/j.geomorph.2016.03.032, 2016.
- 680 Qiu, G. Q.: Development condition of alpine permafrost in the Mt. Tianshan, China, *Journal of Glaciology and Geocryology*, 15, 96-103, 1993. (in Chinese)
- Qiu, G. Q., Huang, Y.Z. and Li, Z.F.: Basic Characteristics of permafrost in Tian Shan, China, *Proceedings of Second National Conference on Permafrost*, Lanzhou, 21–29. 1983. (in Chinese)
- 685 Rosen, P. A., Hensley, S., Joughin, I. R., Li, F. K., Madsen, S. N., Rodriguez, E., and Goldstein, R. M.: Synthetic aperture radar interferometry, *Proceedings of the IEEE*, 88, 333–382, doi:10.1109/5.838084, 2000.
- Sattler, K., Anderson, B., Mackintosh, A., Norton, K., and de Róiste, M.: Estimating permafrost distribution in the Maritime Southern Alps, New Zealand, based on climatic conditions at rock glacier sites, *Frontiers in Earth Science*, 4, doi:10.3389/feart.2016.00004, 2016.
- 690 Schmid, M. O., Baral, P., Gruber, S., Shahi, S., Shrestha, T., Stumm, D., and Wester, P.: Assessment of permafrost distribution maps in the Hindu Kush–Himalayan region using rock glaciers mapped in Google Earth, *The Cryosphere*, 8, 2089–2099, doi:10.5194/tcd-8-5293-2014, 2015.
- Schoeneich, P., Bodin, X., Echelard, T., Kaufmann, V., Kellerer-Pirklbauer, A., Krysiecki, J. M., and Lieb, G. K.: Velocity changes of rock glaciers and induced hazards, *Engineering Geology for Society and Territory*, Vol 1: Climate Change and Engineering Geology, 1, 223–227, doi:10.1007/978-3-319-09300-0_42, 2015.

- 695 Scotti, R., Brardinoni, F., Alberti, S., Frattini, P., and Crosta, G. B.: A regional inventory of rock glaciers and protalus ramparts in the central Italian Alps, *Geomorphology*, 186, 136–149, doi:10.1016/j.geomorph.2012.12.028, 2013.
- Serrano, E., de Sanjose, J. J., and Gonzalez-Trueba, J. J.: Rock glacier dynamics in marginal periglacial environments, *Earth Surf. Processes Landforms*, 35, 1302–1314, doi:10.1002/esp.1972, 2010.
- Shi, Z. W., Xu, J. R., Chen, Z. S., and Li, M. Y.: Analysis on climatic changes under global climatic change: a case
700 study of Tianshan Snow and Avalanche Research Station, *Journal of Mountain Science*, 27, 41–48, 2009. (in Chinese)
- Shi Y.F. and Mi D.S.: Map of Snow, Ice and Frozen Ground in China, SinoMaps Press, Beijing, 1988. (in Chinese)
- Sollid, J. L. and Sørbel, L.: Rock glaciers in Svalbard and Norway, *Permafr. Periglac. Proc.*, 3, 215–220, doi:10.1002/ppp.3430030307, 1992.
- Sorg, A., Kääh, A., Roesch, A., Bigler, C., and Stoffel, M.: Contrasting responses of Central Asian rock glaciers to
705 global warming, *Sci. Rep.*, 5, 8228, doi:10.1038/srep08228, 2015.
- Strozzi, T., Kääh, A., and Frauenfelder, R.: Detecting and quantifying mountain permafrost creep from in situ inventory, space-borne radar interferometry and airborne digital photogrammetry, *Int. J. Remote Sens.*, 25, 2919–2931, doi:10.1080/0143116042000192330, 2004.
- Sun, C., LI, C., Chen, Y., Zhang, M., Nie, Z., and Liu, G.: A long term monitoring of frost heaving in periglacial
710 environment in the Tianshan mountains, *Journal of Glaciology and Geocryology*, 35, 272–279, 2013. (in Chinese)
- Visser, V., Langdon, B., Pauchard, A., Richardson, D. M.: Unlocking the potential of Google Earth as a tool in invasion science. *Biological invasions*, 16(3), 513–2534, 2014.
- Wang, X. W., Liu, G. X., Yu, B., Dai, K. R., Zhang, R., Ma, D. Y., and Li, Z. L.: An integrated method based on
715 DInSAR, MAI and displacement gradient tensor for mapping the 3D coseismic deformation field related to the 2011
Tarlai earthquake (Myanmar), *Remote Sens. Environ.*, 170, 388–404, doi:10.1016/j.jse.2015.09.024, 2015.
- Wegmüller, U., Werner, C., and Strozzi, T.: SAR interferometric and differential interferometric processing chain, *International Geoscience and Remote Sensing Symposium*, 2, 1106–1108, doi: 10.1109/IGARSS.1998.699687, 1998.
- Wessel, P., Smith, W. H. F., Scharroo, R., Luis, J. F., and Wobbe, F.: Generic Mapping Tools: improved version released, *Eos Trans. AGU*, 94, 409–410, 2013.

- 720 Wirz, V., Gruber, S., Purves, R.S., Beutel, J., Gärtner-Roer, I., Gubler, S., and Vieli, A.: Short-term velocity variations at three rock glaciers and their relationship with meteorological conditions. *Earth Surf. Dynam.*, 4, 103–123, 2016.
- Zhao, L., Liu, G. Y., Jiao, K. Q., Li, R., Qiao, Y. Q., and Chien, L. P.: Variation of the Permafrost in the Headwaters of the Urumqi River in the Tianshan Mountains Since 1991, *Journal of Glaciology and Geocryology*, 32, 223-230, 2010. (in Chinese)
- 725 Zhu, C.: Research on the feature of rock glaciers on the central Tian Shan Mountain, *ACTA Geographica Sinica*, 47, 233–241, 1992a. (in Chinese)
- Zhu, C, Cui Z. J., Yao Z.: Some problems from the slope periglacio-landform on middle Tian Shan Mountain, *Mountain Research*, 10, 65–64, 1992b. (in Chinese)

Table 1. List of interferograms made from the seven paths of ALOS PALSAR data. Interferogram names are the conjunction of acquisition dates of the two images. The last column gives the number of identified active rock glaciers (ARGs) in each interferogram.

Path/Frame	Interferogram	Time span (days)	Perpendicular baselines (m)	Number of identified ARGs
501/850	20070902-20071018	46	314	55
502/850	20070619-20070919	92	365	20
503/860	20070706-20071006	92	340	3
504/860	20080424-20080609	46	-12	36
505/860	20080209-20080326	46	211	57
506/870	20070711-20070826	46	256	78
507/870	20090802-20090917	46	586	51

Table 2. Statistical summary of the inventoried active rock glaciers. Each column gives the mean values and the standard deviations (the values in brackets) of the parameters.

	Number	FLP altitude (m)	ILP altitude (m)	Length (m)	Aspect Ratio (-)	Slope (°)	Area (km ²)	Total area (km ²)
All ARGs	261	3174 (262)	3486 (208)	1341 (899)	0.25 (0.19)	14 (6)	0.35 (0.33)	91.5
MARGs	179	3208 (264)	3516 (180)	1521 (988)	0.24 (0.14)	11 (4)	0.43 (0.36)	77.4
TARGs	82	3099 (243)	34205 (247)	947 (465)	0.25 (0.18)	19 (4)	0.17 (0.14)	14.1

Table 3. The lower altitudinal limits of permafrost distribution in the NTS inferred from this study using the inventoried TARGs. ‘This study (Min)’ denotes the altitudinal limits inferred from specific TARGs with the lowest FLP altitudes; ‘This study (Stat)’ denotes the altitudinal limits inferred from the statistics-based calculations of FLP altitudes. The limits from ‘Field surveys’ are based on the work of Qiu et al. (1983). The rows ‘Difference (Min)’ and ‘Difference (Stat)’ list the differences between the results inferred from this study and the field surveys, respectively. All the altitudes given by this study are derived from the SRTM digital elevation model data with an uncertainty of about 16 meters.

	West of 86°E		East of 86°E	
	North-facing	South-facing	North-facing	South-facing
This study (Min)	2490 m	2583 m	2895 m	3088 m
This study (Stat)	2772 m	2828 m	3005 m	3041 m
Field surveys	2730 m	3050 m	2900 m	3250 m
Difference (Min)	-240 m	-467 m	-5 m	-162 m
Difference (Stat)	42 m	-222 m	105 m	-209 m




Figure 1. Topography map of the Northern Tien Shan (NTS) of China. The black dashed boxes outline the footprints of seven frames of PALSAR frames along Path 501–507. The red and blue circles represent moraine-derived active rock glaciers (MARGs) and talus-derived active rock glaciers (TARGs), respectively. The circle size represents rock glacier area. The white dashed box (labeled as ‘Z’) is shown in Figs. 3a and 3b.




Figure 2. Google Earth images of (a) a typical moraine-derived active rock glacier (MARG) and (b) a talus-derived active rock glacier (TARG). The white dots represent the central location of the ARGs with their longitude and latitude given at the bottom of the figures. The inset images show the topographic and morphologic features surrounding the same rock glaciers.




Figure 3. Examples of the identified active rock glaciers in the zones ‘Z’ shown in Fig. 1. (a) shows the wrapped interferometric phase (in radians) for the pair 20070902-20071018 (Path 501). The solid white polygons denote the identified moving targets. (b) shows the corresponding Google Earth image, in which the red and blue polygons denote the identified MARGs and TARGs, respectively. The dashed white polygons denote moving targets that are not rock glaciers (labeled as ‘Non-RGs’). Figs. 3c and 3d show the interferometric phase and optical image for the rock glacier in the northwest corner of Fig. 3a, respectively.




Figure 4. Google Earth image of a rock glacier, with labels showing its initial line point (ILP), front line point (FLP), length (L), and aspect angle (β). The longitude and latitude give the location of the central point (white dot) of the ARG.




Figure 5. Histograms of the ILP altitudes (a–c) and FLP altitudes (d–f) for all ARGs, MARGs, and TARGs, respectively. The minimum (Min), median (Med), and maximum (Max) values for the ILP altitudes and FLP altitudes are denoted. Their units are meters. The box on the top of each sub-figure denotes the box-and-whisker symbol which gives the locations of minimum, the 25 % quantile, the 50 % quantile, the 75 % quantile, and the maximum values. The altitudes are derived from the SRTM digital elevation model data.




Figure 6. (a) Histogram of the aspect angles for the MARGs (red) and TARGs (blue), respectively. (b) Wind rose plot shows the variations of potential income of solar radiation (PISR) with aspect angles.




Figure 7. Scatter plots between the FLP altitudes and (a) the longitude, (b) latitude.




Figure 8. The bottom map shows the down-slope velocities of 170 ARGs measured from SAR interferometry, with the histogram shown in the inset. The top two figures show the down-slope velocities of two pairs of ARGs adjacent to each other. Note that different velocity color bars are used.




Figure 9. Comparison between the mapped ARGs and existing permafrost maps. The green and black dashed lines in (a) shows the mountain permafrost boundaries of CAS permafrost map and the discontinuous permafrost boundaries of IPA map, respectively. The green-to-white colors in (b) show the global Permafrost Zonation Index (PZI) values. Areas without PZI values are shown in black. The PZI maps for the three selected zones 'Z1', 'Z2', and 'Z3' in (b) are further shown in Fig.10. The red and blue dots mark the locations of MARGs and TARGs, respectively.




Figure 10. Local PZI maps for the three selected regions ‘Z1’, ‘Z2’ and ‘Z3’ marked in Fig. 9b. (a) and (c) are selected randomly. (b) shows the MARG and TARG with the lowest FLP altitude within the entire study area. The thin yellow lines in (b-d) are contour lines of PZI values. The red and blue polygons outline the MARGs and TARGs, respectively.




Figure 11. The histograms of the PZI values at the front line point (FLP) for all ARGs (a), MARGs (b), and TARGs (c), respectively.