

This document contains replies to the referees' comments, together with marked text indicating (yellow highlight) all of the changes. The line numbers in the marked document differ from those in the original (due to revisions), and are given in parentheses below.

Reply to comments by Sakai

1) As you wrote 'Comparison of the drainage system structure in 2010 with evidence on Corona imagery from 1964 shows an upglacier expansion of the area occupied by closed depressions and perched lakes'(L623-625) I think analysis on the change of small basins (perched lake area) using Corona imagery (as past image) is useful to know the change of drainage system and to gain more insight of your synoptic view on the drainage system of debris-covered glaciers.

*Although it would be very interesting to analyse long-term basin evolution on the glacier, this is not possible. Our methods for defining basins use contoured DEMs, which cannot be constructed for the Corona data, which consist of mono images. Text has been added in L181-183 to explain this point.*

Iwata et al. (2000) have reported that high relief area expand from 1978 to 1995 at the middle ablation area of the Khumbu Glacier based on the geomorphic evidence. Although the target of Iwata's study was the Khumbu glacier (different from your target; the Ngozumpa Glacier), I think Iwata's result complement your result that lower limit of surface stream area has gone up to higher elevation recently. Therefore, Iwata et al. (2000) would be a nice reference of your manuscript.

*Reference to Iwata et al. (2000) has been added (L 789)*

2) In the section 4.2, 5.1 and 5.5, authors did not discuss on the start point (maximum altitude) of surface stream. The start point of surface stream strongly relates with altitude of ablation and it is significant for drainage system. I think discussion is required in the manuscript on the start point for example the difference between 1964 and 2010 (Fig. 13a and b) there is no solid evidence on the surface stream during LIA, the start point of surface stream during the LIA might (should?) be different from that of 1964. Please take into account the start point of surface stream in Fig. 13c.

*The upper limit of surface streams is not determined by the ELA, but the location of topographically controlled crevasse fields. The 1964 image shows these crevasse fields in the same location. Text has been added (L 747 ff.) to explain this point.*

L34-35 'since the 1970s'> Analysis period by Kääb et al., (2012) was 2003-2008.

*The position of Kääb reference has been changed to remove this problem (now L 40).*

L357-358 (411-413) 'The seasonal variations in ice velocities in the upper ablation zone are too large to be explained by changes in ice creep rates,' > Authors should write the reason

*justification of this statement has been added.*

L 375-385 (436) It's better to cite Fig. 13 in this section.

*A reference to Fig 14 (formerly 13) has been added.*

L487 'measured volume losses' is ambiguous expression. Please write specifically. If the measured volume loss is calculated from elevation change at ablation area, the value includes not only ablation but also emergence velocity.

*In the paper by Thompson et al. (2016) these volume losses were calculated from elevation changes on the stagnant part of the glacier, where complications from glacier flow do not arise. Rather than clutter the text with this information, we have added the phrase 'on the stagnant part of the glacier' on line 564.*

Fig 14 There is no symbols of a) and b) in the Fig. 14, although, authors used Fig. 14a, 14b in the text (L543-549)

*This Figure has been redrafted.*

L528 'where the overall gradient of the glacier is  $<3^\circ$  < reference?

*Statements about the glacier gradient have been clarified (L 121), and a new Figure 3 added.*

L543 'By 2010, this part of the glacier had been broken up into basins E-7, E-8 and E-9'.  
> In other word, you can estimate that basins E-7, E-8 and E-9 has coalesced in 1960s from the Corona image. I recommend if you can draw the basins boundary using Corona image. it would be great help to understand the geomorphic change of the Ngozumpa Glacier. (main comment 1))

*This Figure has been redrafted to make the changes clearer.*

L594 'On the Survey of Nepal map,' > Reference is necessary, here. I think following map is cited here. 'Nepal: Survey Department. 1997c. Namuche Bajar 1 : 50 000. Kathmandu, Ministry of Land Reform and Management. Survey Department. (Sheet No. 2786 03.)' This map was produced based on the aerial photography taken in 1992.

*Reference added.*

L629 'Such a drainage system might have existed during the Little Ice Age, and persisted into the early 20th Century.' > I recommend that the supraglacial channels during the Little Ice Age is not based on Satellite imagery or other evidences. Therefore, the line of the supraglacial channels should be drawn by dotted lines.

*This has been done.*

Please check whole references in the text and in the reference list (not only following comment).

L39 Reynolds, 2000 > I could not find the reference in the reference list.

*Reference deleted in text.*

L146 Thompson et al. (2016) > I could not find the reference in the reference list.

*Reference added.*

L471 Horodyskuj (2015) > I could not find the reference in the reference list.

*Reference added*

L730 The reference has no published year.( *Earth Science Reviews*)

*Year of publication added.*

L748 In the title of Gulley et al. 2009a, 'Mechanisms of ....' has been missed.

*Text corrected*

L800-805 There are two Quincey et al. (2005) but I could not find Quincey et al. (2005) in the body text.

*This has been reduced to one reference, with the correct year of 2007*

Reply to comments by Quincey

1. Part of the justification for the study (lines 63-71) is that we still know relatively little about englacial conduit formation, and specifically the relative importance of the three processes previously described in the two Gulley et al., 2009 papers and summarised in Benn et al., 2012. Although not stated explicitly, the subsequent analysis here suggests that cut-and-closure is the dominant, or even exclusive, mechanism, at least on Ngozumpa. Given that the argument against NG-01 to NG-03 being structurally controlled (lines 308-311) could be invoked for most conduits running parallel to flow, and that to my knowledge there have been no direct observations of hydrofracture here or in the wider region, some actual discussion of their relative importance would be an interesting addition to the manuscript. Based on their additional analysis, do the authors now believe that cut-and-closure is the dominant mechanism for these debris-covered glaciers, or does it just happen to prevail at Ngozumpa? Or is it paired, in that cut-and-closure forms the conduit in the first place, and then the relict channels provide the dominant structural control thereafter? Or some combination of these? Some clarification in the revised text would be a good addition.

*A paragraph has been added (L 691 ff.) to clarify these points.*

2. The least-well constrained element of the paper is the analysis of the existence or otherwise of a subglacial hydrology, understandably so. In the absence of any direct observations, the velocity proxy provides some evidence for subglacial water in the upper ablation area, but if the authors are correct in their interpretation of this, what happens to it then? I'd be interested to see some further discussion of the lower ablation area – if as stated (line 507) all of the water leaving the glacier passes through Spillway Lake, do the authors propose that the subglacial waters from the upper ablation area pass through the lower ablation area and are then elevated at the terminus under pressure? Some direct measurement of the discharge would give an indication of whether it is at least the correct order of magnitude for a glacier of this size, but in the absence of this some discussion of whether it might go to deep groundwater, or shallow groundwater and then emerge further down-glacier, would fill this gap. The hollow that drains the supraglacial channels (line 384) is also important in this regard – it hints at a direct surface-to-bed connection but there is no further information given – can any more light be shed on where these waters go?

*Although we lack sufficient data to answer these questions, we have added speculations (L427 & 572-575) to address the issue of water routing.*

3. The interpretation that hummocky closed basins cannot support a supraglacial hydrology is believable, but there is a spatial mismatch between the analysis shown in Figure 9 and the observations shown in Figure 8, which detracts from the argument. If this is the dominant control on supraglacial water (and the interpretation in Figure

13 hangs on it being so) then can the analysis in Figure 8 be extended up so we can see if it prevails in the upper ablation area too? Or at least see some morphological/topographic differences between c) and d) in Figure 8.

*There is indeed a gap between areas where channels are visible on the glacier surface and area where surface basins can be reliably mapped. We have addressed this mismatch by identifying a 'transitional zone', now discussed in the text (L 463 ff.), and shown on Fig. 9 (previously Fig. 8) and a new Figure 3.*

L86 (100): 7922 m here but 7952 m in Figure 1.

*>The true elevation is 7952 m, and this has been corrected in the text.*

L95 (109): I suggest stating that it is 'effectively' stagnant, since it is probably deforming at some rate, just not detectable by the satellite analysis.

*>The change has been made*

L130 and elsewhere: just a note that the terms pond and lake are used interchangeably – I'd have a preference for using the former for the majority, which are perched and relatively small, and saving the latter for Spillway Lake, which is not.

*> Done throughout, as suggested.*

L158-167: can you add some more detail on the masking/filtering that is evident in Figure 7? And what is the threshold of 'detectable flow' referred to in line 95 – I guess that should be mentioned here too.

*> No masking or filtering was applied. A note on the definition of 'detectable motion' has been added in L195-6.*

L331: Length of stagnant zone - this is 7 km earlier in the manuscript.

*> This has been changed to 6.5 km in both cases (L 110 & 379).*

L351: should these artifacts not be masked as per Fig 7a? As it stands, it looks like you have greater confidence in the speed-up data than the annual pair measurements, which can't be right since you derive the former from the latter.

*> Neither the absolute nor differenced velocity maps have been filtered, and the existence of velocity difference data in areas of no apparent absolute annual displacement is simply a result of the scales used. A note has been added to the caption of Fig. 7 to point out the lack of masking and filtering.*

L364-366 (L 421): there's some contradiction with the Figure caption here – the text says there are only crevasses in the upper part of the clean ice tongue (c), but the caption states they are only in (a)?

*> The text has been corrected.*

L371-373: this is the only hint at what happens to the subglacial water after it descends from the upper ablation area – can you offer any insight into where it may go then? Lines added to 375, and 491 ff, to speculate on fate of subglacial water.

> *We have added the speculation that the subglacial water is drained via the sub-marginal channels (L427 & 572-575).*

L382-385: this hollow is quite an important part of the picture, particularly if it shows the surface is connected to the bed directly – is it the same hollow that Horodyskuj monitored? Is it a moulin? Can you offer any further information on it?

> *This is not the same hollow, as is now made clear in the text (L 551). We have no evidence that direct surface to bed drainage occurs in any of these basins, and decline to speculate about this issue.*

L396: Figure 9 does not extend sufficiently far up-glacier for us to be able to verify this is true – can you extend the analysis to make this argument more robust?

> *Basins cannot be reliably delineated any further upglacier. Text and a new figure have been added to explain this in terms of a 'transitional zone' (L 463 ff.). (see reply to Point 3 above)*

L402-415: the number of basins etc is interesting here but the upper boundary of the analysis seems arbitrarily defined. Why not cover the whole of the debris-covered area? That way others can repeat the analysis for future time periods and quantify the change.

> *see previous comment*

L410: Figure 9 doesn't show any full drainage events, unless these are lumped into the < 1000 m<sup>2</sup> category? Shouldn't they be shown as 'empty'?

> *Empty basins in any one year are indicated by 'missing' coloured circles - The caption has been modified to explain this.*

L425 (692): do they have to be relict, necessarily?

> *'relict' has been changed to 'active or relict'*

L425-427: as above, I can't see the evidence for 35 drainage events – can you make this a bit clearer?

> *see comment on line 410.*

L424-427: can you add an acknowledgement that there is a seasonal signal in these data?

> *We do not have a good enough data series to reliably identify a seasonal signal.*

L439: they're probably underlain by thick sediment too, inhibiting bottom melt.

> *We do not have any data on pond bottom sediment thickness, so decline to comment.*

L456: this disparity between the number of basins on the west and the east sides merits some further comment – does it reflect the dominant englacial drainage pathway? Or debris-thickness? It's a stark contrast when looking at Figure 9.

> *This is indeed striking, but the reason is unknown. We think that speculation neither justified nor helpful.*

L460: missing 'the'

> *added*

L471: Horodyskuj (2015) is not in the reference list.

> *Added*

L507: can you be sure that all of the water passes through Spillway Lake?

> *A sentence has been added to qualify this statement (L 594-6).*

L518-521: can you bring this into line with the six elements stated in the abstract?

> *Done (L 606 ff.).*

L528: 3° gradient.

> *A statement about glacier gradients (line 121) and a new Figure 3 has been added to clarify this issue.*

L543-544: it'd be better to show these basins superimposed on Fig 14b, rather than repeating the channels on both figure panels.

> *The figure has been redrafted to make the evolution of this part of the glacier clearer.*

L545: can you indicate where these elongate ponds are on the figure for clarity?

*Done.*

L546-549: the figure referred to here doesn't relate to the text. Do you mean figure 10 instead? clarify

*This has been corrected.*

L553: where does this thickening debris cover come from?

> *Text has been added to explain this (L 650).*

L557: note cut-and-closure is hyphenated in some places but not in others.

> *Changed to 'cut-and-closure' throughout*

L589: section numbering jumps one here.

> *The numbering has been corrected.*

L626-627: remove the bracketed text since it is clear already?

> *Bracketed text has been removed.*

L662-664: interesting that the Khumbu ponds also follow what might be a sub-marginal channel.

*Agreed. This is something for others to investigate in detail.*

L666: this sentence implies that Ngozumpa is in a more advanced stage of recession than others in the region – is this what you mean? I'm not sure it is much different to others except for Spillway Lake?

> *Not all others - for example, Imja and Trakarding have large base-level lakes. Base level lakes do not yet exist on Khumbu Glacier. Spillway Lake is an important example of a transitional stage.*

L669: maybe 'interpretation' is better than 'view' here?

> *Changed as requested (L 806).*

Figure 7 (now 8): mentioned above too, but how can b) have more coverage than a) given that it is derived from a)?

> *The difference map also depends on the winter velocities (not shown). Also, the velocity scale in a) fades to grey, so small velocities (which can contribute to velocity differences) are not visible.*

Figure 8: what are the coloured dashes here?

> *They delineate the areas identified by the letters.*

Figure 11 (now 12): are there no better data than these TM images? Only because they're poor resolution. Do the 2010 and 2012 data you have not cover these areas? At least could you superimpose your interpreted pond boundaries?

> *Pond boundaries superimposed to make the figure clearer.*

Figure 12 (now 13): maybe add that the reader should see the text for explanation of the annotations?

> *Text added to caption*

## **Marked Text**

1     **Structure and evolution of the drainage system of a Himalayan**  
2            **debris-covered glacier, and its relationship with patterns of**  
3                            **mass loss**

4     Douglas I. Benn<sup>1</sup>, Sarah Thompson<sup>2</sup>, Jason Gulley<sup>3</sup>, Jordan Mertes<sup>4</sup>, Adrian  
5     Luckman<sup>2</sup> and Lindsey Nicholson<sup>5</sup>

6

7     <sup>1</sup> *School of Geography and Sustainable Development, University of St Andrews, UK*

8     <sup>2</sup> *Department of Geography, Swansea University, Swansea, UK*

9     <sup>3</sup> *School of Geosciences, University of South Florida, FL, USA*

10    <sup>4</sup> *Department of Geological and Mining Engineering and Sciences, Michigan Tech,*  
11    *MI, USA*

12    <sup>5</sup> *Institute for Atmospheric and Cryospheric Sciences, University of Innsbruck,*  
13    *Austria*

14

15    **Abstract**

16    This paper provides the first synoptic view of the drainage system of a  
17    Himalayan debris-covered glacier and its evolution through time, based on  
18    speleological exploration and satellite image analysis of Ngozumpa Glacier,  
19    Nepal. The drainage system has several linked components: 1) a seasonal  
20    subglacial drainage system below the upper ablation zone; 2) supraglacial  
21    channels allowing efficient meltwater transport across parts of the upper  
22    ablation zone; 3) sub-marginal channels, allowing long-distance transport of  
23    meltwater; 4) perched ponds, which intermittently store meltwater prior to  
24    evacuation via the englacial drainage system; 5) englacial cut-and-closure

25 conduits, which may undergo repeated cycles of abandonment and reactivation;  
26 6) a 'base-level' lake system (Spillway Lake) dammed behind the terminal  
27 moraine. The distribution and relative importance of these elements has evolved  
28 through time, in response to sustained negative mass balance. The area occupied  
29 by perched ponds has expanded upglacier at the expense of supraglacial  
30 channels, and Spillway Lake has grown as more of the glacier surface ablates to  
31 base level. Subsurface processes play a governing role in creating, maintaining  
32 and shutting down exposures of ice at the glacier surface, with a major impact on  
33 spatial patterns and rates of surface mass loss. Comparison of our results with  
34 observations on other glaciers indicate that englacial drainage systems play a  
35 key role in the response of debris-covered glaciers to sustained periods of  
36 negative mass balance.

37

## 38 **1. Introduction**

39 Debris-covered glaciers in many parts of the Himalaya have undergone  
40 significant surface lowering in recent times (Kääb et al., 2012), with net losses of  
41 several tens of metres since the 1970s (Bolch et al., 2008a, 2011). Glacier  
42 thinning and reduced surface gradients have resulted in lower driving stresses  
43 and ice velocities, and large parts of many glaciers are now stagnant or nearly so  
44 (Bolch et al., 2008b; Quincey et al., 2009). These morphological and dynamic  
45 changes have encouraged formation of supraglacial ponds and lakes and  
46 increased water storage within glacial hydrological systems (Quincey et al.,  
47 2007; Benn et al., 2012). Where lakes form behind dams of moraine and ice,  
48 volumes of stored water can be as high as  $10^8$  m<sup>3</sup>, in some cases posing

49 considerable risk of glacier lake outburst floods (GLOFs) (Yamada, 1998;  
50 Richardson and Reynolds, 2000; Kattelman, 2003).

51

52 Several studies have shown that the development and enlargement of englacial  
53 conduits play an important role in the evolution of debris-covered glaciers  
54 during periods of negative mass balance (e.g. Clayton, 1964; Kirkbride, 1993;  
55 Krüger, 1994; Benn et al., 2001, 2009, 2012; Gulley and Benn, 2007; Thompson  
56 et al., 2016). The collapse of conduit roofs can expose areas of bare ice at the  
57 glacier surface, locally increasing ablation rates. Additionally, areas of subsidence  
58 associated with englacial conduits create closed hollows (dolines) that can  
59 evolve into supraglacial ponds, further increasing ice losses by calving.  
60 Conversely, supraglacial ponds can drain if a connection is made with the  
61 englacial drainage system, provided the pond is elevated above hydrological base  
62 level ('perched lakes' in the terminology of Benn et al., 2001, 2012). Drainage of  
63 relatively warm water through the glacier leads to conduit enlargement, which in  
64 turn increases the likelihood of roof collapse, surface subsidence and ultimately  
65 new pond formation (Sakai et al., 2000; Miles et al., 2015). Because ablation  
66 rates around supraglacial pond margins are typically one or two orders of  
67 magnitude higher than those§ under continuous surface debris, ponds contribute  
68 disproportionately to overall rates of glacier ablation (Sakai et al., 1998, 2000,  
69 2009; Thompson et al., 2016). By controlling the location and frequency of  
70 surface subsidence and pond drainage events, englacial conduits strongly  
71 influence overall ablation rates, and the volume of water that can be stored in  
72 and on the glacier (Benn et al., 2012).

73

74 Speleological investigations in debris-covered glaciers in the Khumbu Himal  
75 have demonstrated that englacial conduits can form by three processes: 1) 'cut-  
76 and-closure' or the incision of supraglacial stream beds followed by roof closure;  
77 2) hydrologically assisted crevasse propagation, or hydrofracturing, which may  
78 route water to glacier beds; and 3) exploitation of secondary permeability in the  
79 ice (Gulley et al., 2009a, b; Benn et al., 2012). The relative importance of these  
80 processes in the development of glacial drainage systems, however, has not been  
81 investigated in detail. Furthermore, there are no data on the large-scale structure  
82 of englacial and subglacial glacial drainage systems in the Himalaya, or how they  
83 evolve during periods of negative mass balance. In this paper, we investigate the  
84 origin, configuration and evolution of the drainage system of Ngozumpa Glacier,  
85 using three complementary methods. First, speleological surveys of englacial  
86 conduits are used to provide a detailed understanding of their formation and  
87 evolution. Second, historical satellite imagery and high-resolution digital  
88 elevation models (DEMs) are used to identify past and present drainage  
89 pathways, glacier-wide patterns of surface water storage and release, and  
90 regions of subsidence. Finally, feature tracking on TerraSAR-X imagery is used to  
91 detect regions of the glacier subject to seasonal velocity fluctuations, as a proxy  
92 for variations in subglacial water storage. Taken together, these methods provide  
93 the first synoptic view of the drainage system of a large Himalayan debris-  
94 covered glacier, and its influence on glacier response to recent warming.

95

## 96 **2. Study area and methods**

97 Ngozumpa Glacier is located in the upper Dudh Kosi catchment, Khumbu Himal,  
98 Nepal (Fig. 1). It has three confluent branches: a western (W) branch flowing

99 from the flanks of Cho Oyu (8188 m); a north-eastern (NE) branch originating  
100 below Gyachung Kang (7952 m); and an eastern (E) branch (Gaunara Glacier)  
101 nourished below a cirque of 6000 m peaks. The NE and E branches are no longer  
102 dynamically connected to the main trunk, which is fed solely by the W branch  
103 (Thompson et al., 2016). The equilibrium line altitude (ELA) is not well known.  
104 Google Earth images from 3 November 2009 (after the end of the ablation  
105 season) and 9 June 2010 (at the beginning of the monsoon accumulation season)  
106 show bare ice up to ~5700 m above sea level (a.s.l.) on all three branches, and  
107 this value is adopted as an approximate value of the ELA.

108

109 The lower ablation zone of the glacier is effectively stagnant, with little or no  
110 detectable motion on most of the E branch, or on the main trunk for ~6.5 km  
111 upglacier of the terminus (Bolch et al., 2008b; Quincey et al., 2009; Thompson et  
112 al., 2016). The lowermost 15 km of the glacier (below ~5250 m a.s.l.) is almost  
113 completely mantled with supraglacial debris. The debris cover thickens  
114 downglacier, reaching  $1.80 \pm 1.21$  m near the terminus (Nicholson, 2004;  
115 Nicholson and Benn, 2012). In common with other large debris-covered glaciers  
116 in the region, Ngozumpa Glacier has undergone significant surface lowering in  
117 recent decades, and the glacier surface now lies >100 m below the crestlines of  
118 the late Holocene lateral moraines (Bolch et al., 2008a, 2011).

119

120 The lower tongue of the glacier has a concave surface profile, with the overall  
121 gradient declining from  $5.8^\circ$  to  $2.4^\circ$  between 5,300 and 4,650 m (Fig. 3). The ice  
122 surface also becomes increasingly irregular downglacier, and below 5,000 m it  
123 forms numerous closed basins separated by mounds, ridges and plateaux with a

124 relative relief of 50 - 60 m (Figs. 2 & 3). Most basins contain supraglacial ponds,  
125 which typically persist for a few years before draining (Benn et al., 2001; 2009,  
126 2012; Gulley and Benn, 2007). Near the terminus of Ngozumpa Glacier, a system  
127 of lakes is ponded behind the terminal moraine (informally named Spillway  
128 Lake; Fig. 1). This lake system increased in area by around 10% per year from  
129 the early 1990s until 2009, but between 2009 and 2015 experienced a reduction  
130 of area and volume as a result of lake level lowering and redistribution of  
131 sediment (Thompson et al., 2012, 2016; Mertes et al., 2016). This hiatus is likely  
132 to be temporary and continued growth of the lake is expected in the coming  
133 years, as has been the case with other 'base-level lakes' in the region (Sakai et al.,  
134 2009).

135

136 We surveyed 2.3 km of englacial passages in Ngozumpa Glacier, using standard  
137 speleological techniques modified for glacier caves (Gulley and Benn, 2007).  
138 Conduit entrances were identified during systematic traverses of the glacier  
139 surfaces. Within each conduit, networks of survey lines were established by  
140 measuring the distance, azimuth and inclination between successive marked  
141 stations using a Leica Distomat laser rangefinder and a Brunton Sightmaster  
142 compass and inclinometer. Scaled drawings of passages in plan, profile and  
143 cross-section were then rendered *in situ*, and include observations of  
144 glaciostructural and stratigraphic features exposed in passage walls, thereby  
145 allowing the origin and evolution of conduits to be reconstructed in detail. In this  
146 paper, we focus on five conduits, which exemplify different stages of conduit  
147 formation, abandonment and reactivation. Three of the conduits have been  
148 previously described by Gulley and Benn (2007), but in this paper we revise our

149 interpretation of their origin in some important respects. Some of the conduits  
150 drained water from or fed water into supraglacial ponds, and in some cases it  
151 was possible to relate phases of conduit development to specific pond filling or  
152 drainage events, identified in satellite images.

153

154 A range of optical imagery was used to map indicators of the large-scale  
155 structure of the drainage system (Table 1). The location of supraglacial channels  
156 and ephemeral supraglacial ponds were mapped using declassified Corona KH-4  
157 imagery from 1965, Landsat 5 TM (2009), GeoEye-1 (9 June 2010 and 23  
158 December 2012) and WorldView-3 (5 January 2015) imagery. The Corona and  
159 Landsat imagery was not co-registered or orthorectified beyond the standard  
160 terrain correction of the product, and was used to identify the presence /  
161 absence of larger ponds or channels and not to quantify rates of change.

162

163 Geo-Eye-1 imagery from June 2010 and December 2012, and Worldview-3  
164 imagery from January 2015 were acquired for a region covering 17.4 km<sup>2</sup> of the  
165 ablation area of the glacier. Three stereoscopic DEMs of 1 m resolution were  
166 constructed from the stereo multispectral imagery using the PCI Geomatica  
167 Software Package, and used to determine spatial patterns of elevation change.  
168 The construction and correction of the DEMs is discussed in detail in Thompson  
169 et al. (2016).

170

171 The 2010 DEM was used to define the extent of individual surface drainage  
172 basins on the glacier surface. This was achieved by identifying surface elevation  
173 contours that entirely surround other contours of a lesser height. Each

174 supraglacial catchment was then defined by the crestlines of ridges that separate  
175 the closed basins. Initially, we used 2 m contours but these produced a large  
176 number of very small 'basins', due to the high roughness of the bouldery glacier  
177 surface. Subsequently, we used 5 m contours that yielded a set of closed basins  
178 that closely matched the location of ephemeral supraglacial ponds on the glacier  
179 surface. The extent of many basins changed between 2010 and 2015 due to ice-  
180 cliff backwasting, although all basins persisted through the period covered by the  
181 DEMs. It was not possible to delineate basins on the historical Corona or Landsat  
182 imagery because our methods depend on the availability of DEMs and cannot be  
183 applied to mono images.

184

185 Glacier surface velocities were derived using feature tracking between synthetic  
186 aperture radar images acquired by the TerraSAR-X satellite on 19 September  
187 2014, 18 and 29 January 2015 and 5 January 2016. Feature tracking was done  
188 using the method of Luckman et al. (2007), which searches for a maximum  
189 correlation between evenly spaced subsets (patches) of each image giving the  
190 displacement of glacier surface features which are converted to speed using time  
191 delay between images. Image patches were ~400 m x 400 m in size and sampled  
192 every 40 m producing a spatial resolution of between 40 and 400 m depending  
193 on feature density. Based on feature matching precise to one pixel (2 m),  
194 precision of the measured velocities is 0.006 m day<sup>-1</sup> over the annual (341 day)  
195 period and 0.018 m day<sup>-1</sup> over the winter (111 day) period. These values are  
196 used to define the threshold for detectable motion on the lower glacier.

197

198 **3. Mechanisms of englacial conduit formation**

199 To provide an overview of processes of englacial conduit formation on the  
200 glacier, we describe two sites in detail (NG-04 and NG-05), then briefly describe  
201 and reinterpret three previously published sites (NG-01, NG-02 and NG-03;  
202 Gulley and Benn, 2007).

203

### 204 *3.1 NG-04*

205 *Description:* Conduit NG-04 (27°57'24"N, 86°41'55" E; 4805 m a.s.l.) was  
206 surveyed in November 2009, and consisted of a main passage (A) and two  
207 shorter side-passages (B and C) leading off to the west (Fig. 4). The main passage  
208 extended from a large hollow on the glacier surface (Basin C-63 on Figs. 2e and  
209 10a) for a distance of at least 473 m, where the survey was discontinued due to  
210 deep standing water on the cave floor. Side-passage B also connected with a  
211 basin on the glacier surface (Basin W-6, Figs. 2e and 10b). Side-passage C was at  
212 least 25 m long, but was not surveyed beyond this distance due to the evident  
213 instability of the highly fractured walls.

214

215 The main passage consisted of an upper level with a flat or gently inclined floor,  
216 and a lower narrow incised canyon. The passage was highly sinuous, with a  
217 sinuosity in the surveyed reach of 5.52. Near A4 (Fig. 4), there was a tight cutoff  
218 meander loop off the main passage (Fig. 5a). The base of the abandoned loop had  
219 a flat floor and lacked the incised lower level that was present elsewhere in the  
220 system. The upper floor level could also be traced along the walls of side  
221 passages B and C, which we interpret as twin remnants of a second meander  
222 cutoff. The floor of the upper level sloped gently downward from A1 to A14, rose  
223 from there to between A18 and A19, after which it descended once more. Sandy

224 bedforms on the floor and scallops on the ice walls of this upper level indicate  
225 that water flow was from A1 towards A21.

226

227 Passage morphology in the upper level was very variable, including tubular, box-  
228 shaped, triangular and irregular sections (Figs. 4 & 5b-d). Throughout most of  
229 the system, planar structures were visible in the ceiling or walls of the upper  
230 level, running parallel to the passage axis with variable inclination. The  
231 structures took the form of: (1) 'sutures' at the line of contact between opposing  
232 walls (S: Fig. 4; Fig. 5b, c), (2) intermittent narrow voids (V: Fig. 4; Fig. 5c), and  
233 (3) bands of sorted sand or gravel a few cm thick (SB: Fig. 4; Fig. 5d). Some of the  
234 voids increased in width inward, in some cases opening out into gaps tens of cm  
235 across. In some places, bands of sorted sediment could be traced laterally into  
236 open voids or sutures. At several points along the main passage, a pair of planar  
237 structures occurred on opposite walls of the passage. Side-passage B had a  
238 narrow, meandering seam of dirty ice running along its ceiling, and in Passage C  
239 the walls tapered upward to meet at a ceiling suture.

240

241 The floor of the incised lower level in both parts of the main passage sloped  
242 down towards side passages B and C (arrows, Fig. 4). A pair of incised channels  
243 was confluent at C1, whereas a single incised channel was present in passage B,  
244 where its lower (western) end was blocked by an accumulation of ice and debris.

245

246 *Interpretation:* The partially debris-filled structures in the walls and ceiling of the  
247 upper level are closely similar to many examples of canyon sutures we have  
248 observed in cut-and-closure conduits in the Himalaya and Svalbard, marking the

249 planes of closure where former passage walls have been brought together by ice  
250 creep and/or blocked by ice and debris (cf. Gulley et al., 2009a, b). Cut-and-  
251 closure conduits are typically highly sinuous and have variable cross sectional  
252 morphologies, ranging from simple *plugged canyons* (incised channels with roofs  
253 of névé), to *sutured canyons* (partially or completely closed by ice creep),  
254 *horizontal slots* (formed by lateral channel migration followed by roof closure),  
255 and *tubular passages* (where passage re-enlargement has occurred under pipe-  
256 full (phreatic) conditions (Gulley et al., 2009b). The tubular morphology of the  
257 upper passage in NG-04, combined with the sutures, voids and sediment bands in  
258 the walls and ceiling indicates that the passage has been re-enlarged under pipe-  
259 full conditions following an episode of almost complete closure. For example, the  
260 sub-horizontal bands of sorted sand on both conduit walls between A15 and A18  
261 (Fig. 5d) suggest complete suturing of a low, wide reach (horizontal slot) prior to  
262 formation of the surveyed passage.

263

264 The tubular and box-shaped cross-profiles and undulating long-profile of the  
265 upper passage are consistent with fluvial erosion under pipe-full or phreatic  
266 conditions (cf. Gulley et al., 2009b). This contrasts with the canyon-like form and  
267 consistent down-flow slope of channels incised under atmospheric (vadose)  
268 conditions, typical of simple cut-and-closure conduits. The dimensions of the  
269 upper passage (typically 2 m high and 3 m wide) are consistent with high  
270 discharges. We conclude that the upper passage formed when water draining  
271 from a supraglacial pond in Basin C-63 exploited the remnants of an abandoned  
272 cut-and-closure conduit (Fig. 2e).

273

274 Following formation of the upper passage, the lower level was incised under  
275 vadose (non pipe-full) conditions when the system accessed a new local base  
276 level via side-passages B and C. We infer that this occurred when a cutoff  
277 meander loop between B1 and C2 was exposed by ice-cliff retreat in Basin W-6.  
278 Water flow between A1 and B2 continued in the same direction as before, but  
279 between A14 and A21 flow was reversed and discharge much reduced.

280

281 Evolution of conduit NG-04 can be summarized as follows: 1) a cut-and-closure  
282 conduit was formed by incision of a supraglacial stream; 2) this conduit was  
283 abandoned and almost completely closed, presumably after it lost all or most of  
284 its source of recharge following downwasting of the overlying glacier surface; 3)  
285 the conduit remnants were exploited and enlarged by water draining from a  
286 supraglacial pond in Basin C-63; and 4) surface ablation in Basin W-6 broke into  
287 the conduit, creating a new base level and initiating floor incision. This  
288 remarkable cave illustrates how relict drainage systems can be reactivated when  
289 connected to new sources of recharge, and demonstrates how patterns of  
290 drainage can change dramatically within a single system in response to changing  
291 surface topography.

292

### 293 3.2 NG-05

294 *Description:* In December 2009 a conduit portal was exposed in an ice cliff at the  
295 margin of Spillway Lake (27°56'36" N, 86°42'46" E, 4670 m a. s. l.; Figs. 2a and 6).  
296 This portal was one of the two main efflux points that discharged water into the  
297 lake from upglacier (Thompson et al., 2016). Access to the conduit could be  
298 gained via the frozen lake surface (Fig. 7a), although the lake ice was broken up

299 each morning by debris falling from the melting glacier surface above, severely  
300 limiting the time available for survey. Consequently, only a short section could be  
301 mapped (Fig. 6). The conduit had two main levels, separated by a narrow,  
302 partially ice-filled canyon. The floor of the lower part was at lake level, and that  
303 of the upper level was 4.8 m higher, close to the summer monsoon level of the  
304 lake, as indicated by shorelines exposed around the lake margins. Several  
305 notches on the passage walls recorded intermediate water levels. The ice cliff  
306 above the upper level was obscured by a mass of icicles, but observations inside  
307 the cave showed that the roof tapered up into a narrow debris band or suture.

308

309 *Interpretation:* Although short, this passage is important for understanding the  
310 drainage system of Ngozumpa Glacier. The debris band and suture in the roof  
311 indicates that, like NG-04, the passage formed by a process of channel incision  
312 and roof closure. Additionally, the passage is graded to the seasonally fluctuating  
313 surface of Spillway Lake. We therefore conclude that the main drainage on the  
314 eastern side of the glacier consists of a cut-and-closure conduit graded to the  
315 hydrologic base level of the glacier. For several km upglacier of the portal, the  
316 debris-covered ice surface is highly irregular and broken into numerous closed  
317 basins, implying that the conduit evolved from a surface stream that predates  
318 significant downwasting of the glacier. The significance of these conclusions will  
319 be discussed later in the paper.

320

### 321 *3.3 NG-01, 02 and 03*

322 *Description:* NG-01, NG-02 and NG-03 (Fig. 2d) were mapped in December 2005,  
323 and described by Gulley and Benn (2007). NG-01 had carried water southward

324 into a large basin on the glacier surface (Basin C-25, Fig. 10a), whereas NG-02  
325 drained water southward out of the basin. NG-01 (27°57'58" N, 86°41'50" E)  
326 was a sinuous canyon passage with three main levels. Debris bands cropped out  
327 in the walls of the uppermost level throughout its length, either at the lateral  
328 margins of the passage or in the roof (Fig. 7b). The mid-level had a sub-  
329 horizontal floor, into which the canyon linking to the lower level had been  
330 incised (Fig. 7c). NG-02 (27°57'55" N, 86°41'51" E) was a sinuous canyon  
331 passage on two levels, extending in a southwesterly direction from the basin. The  
332 upper level had a circular cross profile, and an incised canyon beneath formed  
333 the lower level. A suture and debris band was exposed along the entire length of  
334 the ceiling of the upper passage, mirroring the planform of the passage (Fig. 7d).  
335 The lower level was an asymmetric flat-floored passage with a series of sills  
336 along the margins. NG-03 (27°57'52" N, 86°42'02" E) consisted of a single  
337 passage graded to a supraglacial pond in Basin E-19. Passage morphology varied  
338 between a low, wide semi-elliptical cross-section and a more complex form with  
339 an elliptical upper section separated by a narrow neck from a lower A-shaped  
340 section. At the top of the canyon, the ceiling narrowed to a narrow slot,  
341 terminating in a band of coarse, unfrozen sandy debris.

342

343 *Interpretation:* For much of their length, all three conduits follow the trend of  
344 debris bands in the walls or roof, leading Gulley and Benn (2007) to conclude  
345 that all were structurally controlled. The debris bands were originally  
346 interpreted as debris-filled crevasse traces that had been deformed during  
347 advection downglacier. When the original work was conducted, the cut-and-  
348 closure model had not been developed, and we had yet to learn how to recognize

349 the diverse forms such conduits can take, especially in the later stages of their  
350 development. It is now apparent that these conduits have all the hallmarks of  
351 cut-and-closure conduits. The continuity and sinuous planform of the debris  
352 bands is consistent with formation by the closure of incised canyons, rather than  
353 crevasse fills that had been deformed by ice flow. Crevasses in the upper part of  
354 the glacier ablation area tend to be short, discontinuous and oriented transverse  
355 to flow, unlike the observed debris bands in the conduit roofs, and ice  
356 deformation is unlikely to be capable of generating the highly sinuous patterns  
357 observed within the conduit debris bands.

358

359 We therefore reinterpret NG-01 – 03 as cut-and-closure conduits that have  
360 undergone cycles of incision, abandonment, partial closure and later reactivation  
361 in response to fluctuating patterns of recharge on the glacier surface. The  
362 circular and elliptical cross profiles observed in NG-02 and NG-03 are consistent  
363 with phases of phreatic passage enlargement, analogous to that in NG-04.  
364 Abandoned, incompletely closed conduits create hydraulically efficient flow  
365 paths, which can be readily exploited and enlarged when surface ablation brings  
366 them into contact with new sources of recharge.

367

#### 368 **4. Drainage system structure**

369 In this section, we present evidence for the large-scale structure of the drainage  
370 system and patterns of water storage and release, using X-band radar and optical  
371 satellite imagery and high resolution DEMs from 2010, 2012 and 2015.

372

##### 373 *4.1 Subglacial drainage system*

374 *Observations:* Direct observation of the subglacial drainage system was not  
375 possible. Instead, we use seasonal fluctuations in glacier surface velocity to infer  
376 areas subject to variable subglacial water storage. Mean daily ice velocities of the  
377 glacier between 29 January 2015 and 5 January 2016 are shown in Figure 8a.  
378 There is no detectable motion (i.e. greater than  $\sim 0.01$  m day<sup>-1</sup>) on the main trunk  
379 within  $\sim 6.5$  km of the terminus or on the lowermost 6 km of the E branch. The W  
380 branch is the most active, with velocities of  $\sim 0.16$  m day<sup>-1</sup> ( $\sim 60$  m yr<sup>-1</sup>) at 5300  
381 m a.s.l., declining to near zero at 4900 m. The NE branch is slower, although  
382 velocities in its upper part could not be determined due to image 'lay-over' in  
383 steep terrain. The active part of the NE branch does not extend as far down as  
384 the confluence with the W branch, and a strip of stagnant ice  $\sim 100$  - 200 m wide  
385 extends  $\sim 3$  km down the eastern side of the main trunk from the confluence  
386 zone. Thus, neither the E nor the NE branch is dynamically connected to the main  
387 trunk.

388

389 Evidence for seasonal velocity fluctuations is shown in Fig. 8b, which shows  
390 mean daily velocities between 29 January 2015 and 5 January 2016 (341 days)  
391 minus mean daily velocities from 19 September 2014 to 18 January 2015 (111  
392 days). Meteorological data from the Pyramid Weather Station, at 5050 m a.s.l. c.  
393 12 km east of Ngozumpa Glacier (available through the Ev-K2-CNR SHARE  
394 program), indicate that air temperatures were consistently below freezing  
395 between the 25th of September 2014 and the 28th of May 2015, defining a  
396 minimum winter period for the upper ablation zone. The 111 day interval lies  
397 almost entirely within the winter period but is less than half of its total duration,  
398 so Figure 8b yields minimum values for a summer speed-up on the glacier. Most

399 of the active parts of the glacier exhibit some speed-up, although it is much more  
400 pronounced in some areas than others. On the W branch, the greatest speed-up  
401 (by  $\sim 0.015$  m day<sup>-1</sup> or  $\sim 10\%$ ) occurs above the confluence with the NE branch.  
402 Areas of lesser speed-up also occur on the main trunk below this point, although  
403 these are discontinuous and less than the margin of error so are likely to be  
404 artifacts. Only the northern side of the NE branch is affected by a seasonal speed-  
405 up. This area coincides with the tongue of clean ice that descends through the  
406 icefall below Gyachung Kang (Fig. 1). Patchy areas of apparent speed-up and  
407 slow-down occur elsewhere on the NE branch but may be artifacts. A small  
408 speed-up also affects the active part of the E branch, above 5350 m a.s.l.

409

410 *Interpretation:* The seasonal variations in ice velocities in the upper ablation  
411 zone are too large to be explained by changes in ice creep rates, which would  
412 require fluctuations in driving stress that are inconsistent with the observed  
413 surface elevation changes on the glacier (Thompson et al., 2016). We interpret  
414 the velocity data as evidence for variations in basal motion (sliding and/or  
415 subglacial till deformation) in response to changing subglacial water storage.  
416 This interpretation is supported by the spatial distribution of areas affected by  
417 the seasonal speed-up, which coincide with, or occur downglacier of, heavily  
418 crevassed ice. Much of the upper ablation area of Ngozumpa Glacier consists of  
419 icefalls with surface gradients up to  $30^\circ$ , and fields of transverse crevasses occur  
420 across the entire width of the W branch down to an elevation of 5270 m (a, Fig.  
421 9). Below this zone, crevasses are largely absent, reflecting decreasing ice  
422 velocities and compressive flow (b, c, and d; cf. Fig. 8). Fields of transverse  
423 crevasses occur in the upper basin of the E branch, above  $\sim 5400$  m. Crevasses

424 allow meltwater to be routed rapidly to the bed, and the existence of multiple  
425 recharge points will encourage development of a distributed drainage system  
426 following the onset of the monsoon ablation season. The lack of a clear seasonal  
427 velocity response on the lowermost 10 km of the glacier suggests that subglacial  
428 water is transported along the main trunk in efficient conduits, possibly along  
429 the glacier margins (see Section 4.4).

430

#### 431 *4.2 Supraglacial channels*

432 *Observations:* Supraglacial stream networks are visible below the crevassed  
433 zones on all three branches of the glacier. The most extensive network occurs on  
434 the tongue of clean ice on the NE branch, where a set of sub-parallel channels  
435 descends from ~5180 m to the junction with the W branch at ~4990 m (Fig. 2b,  
436 c; Fig. 14a). There are several discontinuous supraglacial channels on the W  
437 branch between 5220 m and 5120 m a.s.l., including one along the eastern  
438 margin of the glacier. Supraglacial channels occur on both flanks of the E branch  
439 below ~5100 m a.s.l. The channels converge at the junction with the main trunk,  
440 and after flowing over the glacier surface for several hundred metres the  
441 combined stream sinks in a large hollow in Basin E-11. Patterns of water storage  
442 and release in this hollow are discussed in Section 4.4.

443

444 *Interpretation:* Perennial supraglacial channels can only persist if the annual  
445 amount of channel incision exceeds the amount of surface lowering of the  
446 adjacent ice (Gulley et al., 2009b). The rate at which ice-floored channels incise is  
447 controlled by viscous heat dissipation associated with turbulent flow, and  
448 increases with discharge and surface slope (Fountain and Walder, 1998; Jarosch

449 and Gudmundsson, 2012). Because supraglacial stream discharge is a function of  
450 surface melt rate and melt area, significant channel incision requires large  
451 catchment areas. Therefore, incised surface channels tend to occur only where  
452 potential catchments are not fragmented by crevasses or hummocky surface  
453 topography (Fig. 2). At present, these conditions are met in relatively limited  
454 areas of Ngozumpa Glacier, below crevassed areas and above hummocky debris-  
455 covered areas.

456

#### 457 *4.3 Hummocky debris-covered areas and perched ponds*

458 *Observations:* Most of the lower ablation zone of the glacier (below ~5,000 m)  
459 consists of hummocky debris-covered topography, where the glacier surface is  
460 broken up into distinct closed depressions, each of which forms a separate  
461 surface drainage basin (Fig. 2d, e). Not including the Spillway Lake basin that  
462 drains externally, we defined 111 surface basins in this zone in 2010 (Fig. 10).  
463 Some surface basins also occur between 5,000 and 5,100 m on the W Branch, but  
464 these are typically small, shallow and ill-defined (Fig. 2). This part of the glacier  
465 is steeper ( $3.4^\circ$ ) and has lower relative relief (~10 m) than the lower glacier, and  
466 appears to be a transitional zone between the channelized upper ablation area  
467 and the hummocky debris-covered zone (Fig. 3). Surface basins along the east  
468 and west margins of the glacier form a series of depressions within almost  
469 continuous lateral troughs, and are considered in Section 4.4. Here, we focus on  
470 the basins in the central part of the glacier (C1 - C69; Fig. 10a) and the terminal  
471 zone (T1 - 6, Fig. 10b).

472

473 Of the 70 basins in the central part of the glacier, 56 (80%) contained ponds in  
474 at least one of the three years covered by the Geo-Eye and Worldview imagery.  
475 Fifteen of the 42 ponds present in 2010 (36%) had disappeared by 2012 or  
476 2015, whereas 14 basins that were empty in 2010 contained ponds in one or  
477 more of the later years. Almost all of the remainder underwent partial drainage  
478 and /or refilling. In contrast, the 5 ponds in the terminal zone of the glacier  
479 (below Spillway Lake) exhibited great stability. Four showed no significant  
480 change in area between 2010 and 2015, while the other showed an increase in  
481 area.

482

483 *Interpretation:* Observations on and below the glacier surface show that drainage  
484 of perched ponds occurs when part of the floor is brought into contact with  
485 permeable structures in the ice (Benn et al., 2001; Gulley and Benn, 2007). The  
486 characteristics of NG-01 - 05 (which all occur within the hummocky debris-  
487 covered zone) show that relict cut-and-closure conduits are the dominant cause  
488 of secondary permeability in the glacier, providing pre-existing lines of weakness  
489 along which perched ponds can drain.

490

491 The spatial extent and high temporal frequency of perched pond drainage events  
492 on the glacier (Fig. 10a) imply a high density of active or relict conduits within  
493 the ice. A rough estimate can be obtained by dividing the number of complete  
494 and partial drainage events (35) by the total area of basins in the central part of  
495 the glacier (4.62 km<sup>2</sup>), yielding ~7.6 relict conduit reaches per km<sup>2</sup>. This is a  
496 minimum estimate, because additional conduit remnants could occur below and  
497 beyond the margins of observed ponds. Conversely, the number of pond filling

498 events (23 over the 5 ablation seasons spanned by the imagery) shows that  
499 drainage routes commonly become blocked. Conduit blockage processes have  
500 been described by Gulley et al. (2009b), and include accumulation of icicles or  
501 floor-ice at the end of the melt season and creep closure of opposing conduit  
502 walls. The interplay between drainage events and conduit blockage maintains a  
503 dynamic population of supraglacial ponds, which contribute significantly to  
504 ablation of the glacier, through absorption of solar radiation and ice melt, and  
505 calving (Thompson et al., 2016).

506

507 The stability of ponds in the terminal zone probably reflects a combination of  
508 factors. These ponds are flanked by stable slopes of thick debris, which inhibit  
509 pond growth by melt or calving. Furthermore, the ponds are located at or close  
510 to the hydrologic base level of the glacier, determined by the terminal moraine  
511 that encircles the glacier terminus, inhibiting drainage via relict conduits.

512

#### 513 *4.4 Sub-marginal drainage*

514 *Observations:* Elevation differences between successive DEMs indicate linear  
515 zones of enhanced surface lowering along both margins of Ngozumpa Glacier,  
516 forming troughs along the base of the bounding lateral moraines (Thompson et  
517 al., 2016; Fig. 11). The inner moraine slopes consist of unvegetated,  
518 unconsolidated till, and undergo active erosion by a range of processes including  
519 rockfall, debris flow and rotational landslipping (Benn et al., 2012; Thompson et  
520 al., 2016). Although debris eroded from the moraine slopes is transferred  
521 downslope into the troughs, the troughs underwent surface lowering of 6 – 9 m  
522 from 2010 to 2015, with a total annual volume loss in the moraine-trough

523 systems of  $\sim 0.4 \times 10^6 \text{ m}^3 \text{ yr}^{-1}$  (Thompson et al., 2016). This implies that a large  
524 volume of ice, debris or both is evacuated annually from below the lateral  
525 margins of the glacier.

526

527 The lateral troughs form a series of closed basins, 12 on the west side and 22 on  
528 the east (Fig. 10b). Eight of the basins in the west trough and 17 of those in the  
529 east contained a pond in 2010, 4 (W) and 7 (E) of which had completely drained  
530 by 2012 or 2015. Four new ponds appeared in the eastern trough in 2012 or  
531 2015, and 1 (W) and 7 (E) underwent partial drainage and/or refilling. Three  
532 basins on the western side and one on the eastern side showed no fluctuations in  
533 pond area.

534

535 Benn et al. (2001) provided detailed descriptions of pond filling and drainage  
536 cycles in basins W-7 and W-5 (Lakes 7092 and 7093 in their terminology). In  
537 October 1998, basin W-7 contained three shallow ponds, but by October 1999  
538 the basin was occupied by a single large pond and water level had risen by  $\sim 9$  m.  
539 Pond area had increased from  $17,890 \text{ m}^2$  to  $52,550 \text{ m}^2$ , with 36% of the increase  
540 attributable to backwasting and calving of the surrounding ice cliffs. By  
541 September 2000, the pond had almost completely drained and only shallow  
542 ponds remained. Pond drainage occurred via an englacial conduit, which had  
543 been exposed by retreat of the pond margin. A pond in basin W-5 also  
544 underwent fluctuations in area and depth between 1998 and 2000, but did not  
545 completely drain during that time. Horodyskuj (2015) used time-lapse  
546 photography and a pressure transducer to document rapid pond-level

547 fluctuations in basin W-5, including rises and falls of several metres within  
548 hours.

549

550 Short-term cycles of pond drainage and filling can also be demonstrated in other  
551 basins within the lateral trough systems using optical satellite imagery. Figure 12  
552 shows a series of images of the east side of the glacier close to the junction with  
553 the E branch, where a supraglacial stream (Section 4.2) flows into a closed  
554 depression in Basin E-11 (Fig. 10b). A pond occupying the basin expanded in  
555 area between March and May 2009, but drained between June and August. In  
556 2015 there is little evidence of the pond in January but a large pond is present in  
557 June.

558

559 *Interpretation:* Widespread, rapid subsidence along both margins of the glacier  
560 can be explained by enlargement and episodic collapse of sub-marginal conduits  
561 (Thompson et al., 2016). Potential internal ablation rates were calculated from  
562 energy losses associated with runoff and supraglacial pond drainage, and the  
563 resulting value of 0.12 to 0.13 x 10<sup>6</sup> m<sup>3</sup> yr<sup>-1</sup> is around 30% of the measured  
564 volume losses in the moraine-trough systems on the stagnant part of the glacier,  
565 the difference being at least partly attributable to sediment evacuation by  
566 meltwater.

567

568 The sub-marginal conduits are perennial features of the glacier drainage system,  
569 and discharge water into Spillway Lake during the winter months. Winter  
570 discharge may partly reflect slow release of water from supraglacial and  
571 englacial storage, but it may also partly consist of subglacial water from the

572 upper ablation zone (see Section 4.1). This hints at the possibility that the sub-  
573 marginal channels function as the downglacier continuations of the subglacial  
574 drainage system, in addition to carrying water transferred more directly from  
575 the glacier surface.

576

577 Much of the lower ablation zone appears to be bypassed by the sub-marginal  
578 conduits, as evidenced by widespread water storage in supraglacial ponds  
579 (Section 4.3). As noted above, water is intermittently discharged from ponds in  
580 the central part of the glacier into the lateral troughs via englacial conduits.  
581 Cycles of pond drainage and filling in lateral basins indicate intermittent  
582 connections between surface catchments and the sub-marginal meltwater  
583 channels (Fig. 10b). In some cases, drainage events can be directly attributed to  
584 exploitation of englacial conduits (Benn et al., 2001). The hourly changes in pond  
585 level recorded by Horodyskuj (2015) cannot be explained by conduit opening  
586 and blockage, and more likely reflect short-term fluctuations in recharge from  
587 surface melt and water release from storage.

588

#### 589 *4.5 Spillway Lake*

590 *Observations:* In 2010, the area of the Spillway Lake surface catchment was 0.8  
591 km<sup>2</sup>, of which 0.27 km<sup>2</sup> was occupied by the lake system. All of the water leaving  
592 the glacier passes through Spillway Lake, entering via portals or upwellings at or  
593 close to lake level, and leaving via a gap in the western lateral moraine ~1 km  
594 from the glacier terminus (1: Fig. 13). (It is possible that water also exits the  
595 glacier via groundwater flow, although no springs have been observed in the  
596 frontal moraine ramp.) In 2009, conduit NG-05 (Fig. 6; Section 3.2) entered the

597 NE corner of the Spillway Lake and is interpreted as the distal part of the eastern  
598 sub-marginal conduit. A second conduit portal visible at the NW lake margin in  
599 the same year is interpreted as the efflux point of the western sub-marginal  
600 stream. The evolution of the Spillway Lake system, and its implications for  
601 drainage system structure in this part of the glacier, are examined in Section 5.4  
602 below.

603

#### 604 *4.6 Summary*

605 The evidence presented above demonstrates that the drainage system of  
606 Ngozumpa Glacier comprises **six linked elements**: 1) a seasonal subglacial  
607 drainage system below the upper ablation zone; 2) supraglacial channels  
608 allowing efficient meltwater transport across parts of the upper ablation zone; 3)  
609 sub-marginal channels, allowing long-distance transport of meltwater; 4)  
610 perched ponds, which intermittently store meltwater prior to evacuation via the  
611 englacial drainage system; 5) englacial cut-and-closure conduits, which may  
612 undergo repeated cycles of abandonment and reactivation; 6) a 'base-level' lake  
613 system (Spillway Lake) dammed behind the terminal moraine. These elements  
614 have a distinct spatial distribution (Fig. 14a). Evidence for seasonal subglacial  
615 water storage is restricted to active parts of the glacier downglacier of crevasse  
616 fields, where surface water can be routed to the bed. Supraglacial channels occur  
617 where surface catchments and discharge are large enough to allow channel  
618 incision rates to outpace surface ablation rates. Thus, perennial channels only  
619 occur where the glacier surface is not broken up by crevasse fields or into small,  
620 closed basins. Perched ponds occur where the glacier surface is broken up into  
621 closed basins, where the overall gradient of the glacier is  $<2.4^\circ$ . The life cycle of

622 perched ponds is governed by the location of englacial 'cut-and-closure' conduits  
623 and the frequency of connection and blockage events. Sub-marginal conduits  
624 occur below both flanks of the glacier, and transport water from supraglacial  
625 channels, intermittent drainage from perched ponds, and possibly the subglacial  
626 drainage system, into Spillway Lake. The lake lies at the hydrologic base level of  
627 the glacier, and its extent reflects the surface elevation of the glacier relative to  
628 the spillway through the terminal moraine.

629

## 630 **5. Evolution of the drainage system**

631 In this Section, we present evidence for changes in drainage system structure  
632 through time, including features visible on Corona images from 1964 and 1965,  
633 speleological observations, and repeat surveys of Spillway Lake since 1999.

634

### 635 *5.1 Supraglacial channels*

636 In 1964, a connected supraglacial drainage stream network was present on the  
637 eastern side of the main trunk above the junction with the E branch (10 - 8 km  
638 from the terminus, 4950 m to 4920 m a.s.l.) (Fig. 15a). By 2010, this part of the  
639 glacier had been broken up into basins E-7, E-8 and E-9, part of the lateral trough  
640 systems described in Section 4.4. Stream channels were no longer present,  
641 although a number of isolated elongate ponds occupied depressions close to  
642 some of the original channel locations (Fig. 15b). The depressions have an  
643 overall reduction in elevation to the south, but in detail they have up-and-down  
644 long profiles. In cross profile, they are U-shaped and become wider and deeper  
645 through time (Fig. 15c).

646

647 We hypothesize that the supraglacial channels became deeply incised and  
648 transitioned into cut-and-closure conduits, which continue to evacuate  
649 meltwater below the glacier margins despite fragmentation of the surface  
650 topography. Channel incision may have been encouraged by thickening debris  
651 cover (from melt-out of englacial debris) that would have reduced glacier surface  
652 lowering rates.

653

654 At the distal end of the eastern lateral trough, conduit NG-05 (Fig. 6) emerges  
655 into Spillway Lake. Passage morphology indicates that at this point the conduit  
656 formed by cut-and-closure (Section 3.2). Thus, there is evidence for a cut-and-  
657 closure origin of subsurface conduits at both ends of the eastern lateral trough.  
658 We therefore infer that the sub-marginal conduits originated as supraglacial  
659 streams that became incised below the surface. Such a scenario would require a  
660 continuous slope along both glacier margins. We conclude that supraglacial  
661 streams occurred along both margins before development of the current  
662 irregular topography, but transition to cut-and-closure conduits allowed these  
663 drainage routes to persist after break-up of the glacier surface.

664

### 665 *5.2 Englacial conduits in the hummocky debris-covered zone*

666 Transition of drainages from supraglacial channels to cut-and-closure conduits  
667 appears to have been a widespread process on the glacier. The presence of  
668 sutures, planar voids and bands of sorted sediments in the ceilings and walls of  
669 conduits NG-01 - NG-05 record former episodes of channel incision. As was the  
670 case for the lateral channels, we infer that systems of supraglacial channels

671 existed in the central part of the lower tongue before the glacier surface was  
672 broken up into small closed basins (Fig. 14c).

673

674 Differential surface ablation can eventually cause fragmentation and  
675 abandonment of cut-and-closure conduits, cutting off downstream reaches from  
676 former water sources. In abandoned reaches, processes of passage closure  
677 dominate over those of enlargement, and systems gradually shut down. Because  
678 cut-and-closure conduits are generally located close to the glacier surface, shut-  
679 down is commonly incomplete. Zones of narrow voids or sutures with infills of  
680 unfrozen sediment may persist, forming meandering lines of high permeability  
681 through otherwise impermeable glacier ice.

682

683 Reactivation of abandoned conduits will occur if a new water source becomes  
684 available, and a conduit remnant connects this source with a region of lower  
685 hydraulic potential. These conditions are met on stagnant, low-gradient glacier  
686 surfaces. Supraglacial ponds in closed basins provide both reservoirs of water  
687 and regions of elevated hydraulic potential. Drainage is highly episodic, and  
688 water may be stored in supraglacial ponds for years before passing farther down  
689 the system.

690

691 Cut-and-closure is the dominant primary process of conduit formation on  
692 Ngozumpa Glacier, and active and relict cut-and-closure conduits create a  
693 secondary permeability that can be exploited by water from supraglacial ponds.  
694 Debris-filled crevasse traces may provide additional lines of weakness in some  
695 cases, although this is likely a minor process. We have not observed

696 hydrofracture-type conduits in the debris-covered area of Ngozumpa Glacier,  
697 although it is possible that they may form under compressive flow conditions as  
698 described on Khumbu Glacier by Benn et al. (2009). Hydrofracturing likely plays  
699 a dominant role in surface-to-bed drainage in the crevasse fields of the upper  
700 ablation zone.

701

### 702 *5.3 Spillway Lake*

703 The recent history of Spillway Lake was discussed in detail by Thompson et al.  
704 (2012, 2016), and is briefly reviewed here. The present spillway through the SW  
705 side of the terminal moraine has been in existence since at least 1965, when  
706 water emerged from the glacier and entered a small pond behind the lateral  
707 moraine (1: Fig. 13). In the following decades, the Spillway Lake system  
708 expanded upglacier from this point. On the Survey of Nepal map (Nepal Survey  
709 Department, 1997) based on aerial photographs taken in 1992, the lake has a  
710 ribbon-like form, extending NE for ~600 m from the spillway. The lake had  
711 essentially the same outline at the time of our first field survey in 1998, when  
712 water was observed to enter the lake via a subaerial portal and an upwelling  
713 point (Fig. 13; Benn et al., 2001; Thompson et al., 2012). Between 1998 and  
714 1999, several chasms and holes opened up on the glacier surface north of the  
715 western portal, and by 2001 these had evolved into linear ponds and lakes (2:  
716 Fig. 13). Between 2001 and 2009, the Spillway Lake system underwent  
717 considerable expansion to the north, accompanied by upglacier migration of the  
718 portal locations (3, 4: Fig. 13).

719

720 The predominantly linear patterns of lake expansion, and the location of  
721 meltwater portals and upwellings, indicate that evolution of the Spillway Lake  
722 system was strongly preconditioned by the locations of shallow englacial  
723 conduits (a, b: Fig. 13). Conduit NG-05 (Section 3.4 and Fig. 6) and other  
724 examples exposed around the lake margins show that the drainage system  
725 consists of cut-and-closure conduits graded to lake level. This near-surface  
726 englacial conduit system provided pre-existing lines of weakness in the ice  
727 which, when opened up to the surface by internal ablation and collapse, were  
728 exploited by ice-cliff melting and calving processes.

729

730 Spillway Lake was thus established on a template provided by two englacial  
731 conduits (a & b, Fig. 13), which were confluent prior to 1992. As it expanded  
732 upglacier, Spillway Lake encroached on areas formerly occupied by perched  
733 ponds and incorporated former supraglacial basins. A recent example is Basin C-  
734 33, which forms an inlier within the Spillway Lake catchment (Figs. 10a and 13).  
735 This basin contained a perched pond in 2009 and 2010, but this drained prior to  
736 December 2012 and has not reformed. It is likely that this basin will become  
737 entirely subsumed within the Spillway Lake catchment in the near future, as a  
738 consequence of ice-cliff backwasting.

739

#### 740 *5.4 Changing drainage patterns on the glacier*

741 Comparison of the drainage system structure in 2010 with evidence on Corona  
742 imagery from 1964 shows an upglacier expansion of the area occupied by closed  
743 depressions and perched ponds, and the formation and upglacier expansion of  
744 the base-level Spillway Lake (Fig. 14b). The widespread occurrence of cut-and-

745 closure conduits provides evidence of an even earlier stage in drainage  
746 evolution, when supraglacial channels extended along most of the glacier tongue  
747 and closed basins were absent or rare (Fig. 14c). The upglacier limit of  
748 supraglacial channels was similar in 1964 and 2010, due to the persistent  
749 location of crevasse fields in the upper ablation zone. The channels are likely to  
750 have had similar upglacier limits in earlier times, because of the strong  
751 topographic control of the crevasse fields. Figure 14c shows a hypothetical  
752 distribution of supraglacial channels on the glacier during the Little Ice Age and  
753 early 20th Century.

754

755 Ngozumpa Glacier has thus responded to a prolonged period of negative mass  
756 balance with a systematic reordering of its drainage system, characterized by  
757 less efficient evacuation of meltwater and greater amounts of storage. More  
758 recent elements of the drainage system retain a memory of older elements, and  
759 processes and patterns of ablation on the glacier continue to be influenced by  
760 active and relict channels and conduits. Former supraglacial channels  
761 preconditioned the location and density of cut-and-closure conduits, which in  
762 turn precondition the formation and drainage of perched ponds and provide  
763 templates for the expansion of Spillway Lake.

764

## 765 **6. Comparison with other debris-covered glaciers**

766 Observations on other debris-covered glaciers in the Himalaya indicate that their  
767 drainage systems share many of the characteristics described in this paper.  
768 Seasonal velocity fluctuations have been documented on other large glaciers in  
769 the Mount Everest region and on Lirung glacier, Nepal (Benn et al., 2012;

770 Kraaijenbrink et al., 2016), indicating surface-to-bed drainage and variations in  
771 subglacial water storage. Perennial supraglacial channels occur in the upper  
772 ablation zones of many glaciers, in places where catchments are not fragmented  
773 by crevasse fields or irregular surface topography (Gulley et al., 2009b; Benn et  
774 al., 2012). Continuity between a supraglacial channel and an englacial cut-and-  
775 closure conduit has been observed on Khumbu Glacier, clearly demonstrating the  
776 genetic relationship between the two features (Gulley et al. 2009b). Perched  
777 ponds are widespread on Himalayan debris-covered glaciers, and evidence for  
778 repeated filling and drainage (Watson et al., 2016; Miles et al., 2017) suggest that  
779 englacial conduits may play an important role in their life cycles. However,  
780 englacial conduits have only been explored in a few glaciers (Gulley and Benn,  
781 2007; Gulley et al. 2009b; Benn et al. 2009), and much research remains to be  
782 done. Similarly, very little is known about possible sub-marginal channels in  
783 Himalayan glaciers, and our few attempts to enter these highly dynamic  
784 environments have been repulsed.

785

786 The upglacier expansion of the area occupied by closed depressions and perched  
787 ponds on Ngozumpa Glacier (Fig. 14) also appears to have occurred on other  
788 glaciers in the Everest region during the current period of negative mass balance.  
789 Iwata et al. (2000) noted an increase in the area occupied by high-relief  
790 hummocky topography on Khumbu Glacier from 1978 to 1995. The presence of  
791 cut-and-closure conduits below hummocky terrain on that glacier shows that  
792 these areas formerly supported supraglacial streams (Gulley et al., 2009b).

793

794 There is strong evidence on many glaciers that growth of base-level lakes is  
795 preconditioned by englacial conduits. For example, upglacier expansion of the  
796 proglacial lake at Tasman Glacier, New Zealand, has repeatedly followed the  
797 locations former chains of sink holes on the glacier surface (Kirkbride, 1993;  
798 Quincey and Glasser, 2009). Recently formed chains of ponds on the lower  
799 ablation zone of Khumbu Glacier, strongly suggests that the same process is  
800 underway on that glacier (Watson et al., 2016). The integrated picture of  
801 drainage system structure and evolution presented in this paper provides a  
802 framework for predicting what the future may have in store for other debris-  
803 covered glaciers in the region.

804

## 805 **7. Summary and Conclusions**

806 This paper has provided the first **synoptic interpretation** of the drainage system  
807 of a Himalayan debris-covered glacier, including the spatial distribution of  
808 system components, their evolution through time, and their influence on  
809 processes and patterns of ablation. Our specific conclusions are as follows.

810 1) In the upper ablation zone, seasonal variations in ice velocity indicate routing  
811 of surface meltwater to the bed via crevasses, and fluctuations in subglacial  
812 water storage.

813 2) Systems of supraglacial channels occur where the glacier surface is  
814 uninterrupted by crevasses or closed depressions, allowing efficient evacuation  
815 of surface melt.

816 3) Active sub-marginal channels are evidenced by linear zones of subsidence  
817 along both margins of the glacier, and fluctuations in surface water storage and  
818 release. These channels likely formed from supraglacial channels by a process of

819 cut-and-closure, and permit long-distance transport of meltwater through the  
820 ablation zone. Transport of sediment via the lateral channels destabilizes inner  
821 moraine flanks and delivers debris to the terminal zone, where it modulates  
822 ablation processes.

823 4) In the lower ablation zone (below ~5,000 m) the glacier surface consists of  
824 numerous closed drainage basins. Meltwater in this zone typically undergoes  
825 storage in perched ponds before being evacuated via the englacial drainage  
826 system. Englacial conduits in this zone evolved from supraglacial channels by a  
827 process of cut-and-closure, and may undergo repeated cycles of abandonment  
828 and reactivation. Cut-and-closure is the dominant process of conduit formation  
829 on Ngozumpa Glacier, and is likely so on other debris-covered glaciers in the  
830 Himalaya.

831 5) Enlargement of englacial conduits removes ice mass that is not captured by  
832 surface observations until conduit collapse occurs, with the implication that  
833 observations of sudden surface lowering need not reflect sudden glacier mass  
834 loss over the same time period. Subsurface processes play a governing role in  
835 creating, maintaining and shutting down exposures of ice at the glacier surface,  
836 with a major impact on spatial patterns and rates of surface mass loss.

837 6) A large lake system (Spillway Lake) is dammed behind the terminal moraine,  
838 which forms the hydrologic base level for the glacier. Since the early 1990s,  
839 Spillway Lake has expanded upglacier, exploiting weaknesses formed by  
840 englacial conduits.

841 7) As part of the glacier response to the present ongoing period of negative mass  
842 balance, the structure of the drainage system has changed through time,  
843 characterized by decreasing efficiency and greater volumes of storage. Processes

844 and patterns of ablation on the glacier are strongly influenced by active and relict  
845 elements of the drainage system. Former supraglacial channels evolved into cut-  
846 and-closure conduits, which in turn precondition the formation and drainage of  
847 perched ponds and provide templates for the expansion of Spillway Lake. Thus  
848 drainage elements that initially formed during earlier active phases of the  
849 glacier's history continue to influence its evolution during stagnation.

850

851

852

853

854

855

856

857

858

859