

# Responses of sub-ice platelet layer thickening rate and frazil ice concentration to variations in Ice Shelf Water supercooling in McMurdo Sound, Antarctica

Chen Cheng<sup>1,2,3</sup>, Adrian Jenkins<sup>2</sup>, Paul R. Holland<sup>2</sup>, Zhaomin Wang<sup>4</sup>, Chengyan Liu<sup>4,1,3</sup>, and Ruibin Xia<sup>1,3</sup>

<sup>1</sup>Polar Climate System and Global Change Laboratory, Nanjing University of Information Science & Technology, Nanjing, 210044, China

<sup>2</sup>British Antarctic Survey, Cambridge, CB3 0ET, UK

<sup>3</sup>School of Marine Sciences, Nanjing University of Information Science & Technology, Nanjing, 210044, China

<sup>4</sup>College of Oceanography, Hohai University, Nanjing, 210098, China

*Correspondence to:* Zhaomin Wang (zhaomin.wang@hhu.edu.cn)

**Abstract.** Persistent outflow of supercooled Ice Shelf Water (ISW) from beneath McMurdo Ice Shelf creates a rapidly growing sub-ice platelet layer having a unique crystallographic structure under the sea ice in McMurdo Sound, Antarctica. A vertically-modified frazil-ice-laden ISW plume model that encapsulates the combined nonlinear effects of the vertical distributions of supercooling and frazil concentration on frazil ice growth is applied to McMurdo Sound, and is shown to reproduce the observed ISW supercooling and platelet layer distributions. Using this model, the dependence of platelet layer thickening rate and depth-averaged frazil ice concentration on ISW supercooling in McMurdo Sound is investigated, and found to be predominantly controlled by the vertical distribution of frazil concentration. The complex dependence on frazil concentration highlights the need to improve frazil ice observations within the sea ice-ocean boundary layer in McMurdo Sound.

## 1 Introduction

Ice shelf basal melting removes more mass from the Antarctic Ice Sheet than iceberg calving does, but the three largest ice shelves, Filchner-Ronne, Ross, and Amery, contribute only 18% of the net meltwater flux (Rignot et al., 2013). That is because the seawater-filled cavities beneath those ice shelves are dominated by High Salinity Shelf Water that has a potential temperature at or near the surface freezing point. Ice shelf basal melting occurs at depth, because the freezing point temperature is lower under elevated pressure, and results in the formation of Ice Shelf Water (ISW), characterized by temperatures below the surface freezing point. When the buoyant ISW ascends along the ice shelf base, the pressure relief causes it to become supercooled in situ, a necessary condition for ice crystals to persist in suspension. Those disk-shaped frazil ice crystals accumulate under the ice shelves, leading to the formation of marine ice that is thicker and more localized than would be possible through direct freezing at the ice shelf base (Morgan, 1972; Oerter et al. 1992; Fricker et al., 2001;

Holland et al., 2007, 2009). Occasionally, frazil ice crystals bathed in supercooled ISW are also carried out beyond the ice shelf front and precipitated under adjacent sea ice, forming an unconsolidated, porous, sub-ice platelet layer (Hunkeler et al., 2016; Langhorne et al., 2015; Leonard et al., 2006; Robinson et al., 2014). The platelet layer not only harbours some of the highest concentrations of sea ice algae on Earth (Arrigo et al., 2010) but also contributes to the sea ice thickness when the water within the pores of the platelet layer freezes to become incorporated ice (Smith et al., 2001). **Therefore**, the platelet layer should not be ignored when investigating sea ice thickness near an ice shelf front.

Owing to the paucity of direct observation, our understanding of the evolution of frazil-ice-laden ISW relies heavily on numerical models. Those models are mostly derived from plume theory (Holland and Feltham, 2006; Jenkins and Bombosch, 1995; Rees Jones and Wells, 2018; Smedsrud and Jenkins, 2004), **but include three-dimensional ocean circulation models** (Galton-Fenzi et al., 2012), and have been widely applied to assess the marine ice beneath Filchner-Ronne (Bombosch and Jenkins, 1995; Holland et al., 2007; Smedsrud and Jenkins, 2004), Larsen (Holland et al., 2009) **and Amery** ice shelves (Galton-Fenzi et al., 2012), and the **platelet layer** under the sea ice in McMurdo Sound (Hughes et al., 2014, hereinafter HU14). To date, all the ISW plume models **mentioned above** have been depth-integrated, and all the **scalar** quantities, **i.e.**, **water temperature, salinity, and frazil concentration** in those models are treated as vertically-uniform. **The well-mixed temperature and salinity have been validated by borehole observations beneath the Amery Ice Shelf (Herraiz-Borreguero et al., 2013) and under the sea ice in McMurdo Sound (Robinson et al., 2014; HU14). Although there are no observations of the vertical profile of frazil ice concentration, it is unlikely to be vertically uniform because the buoyant rise of the crystals will counteract the turbulent diffusion that tends to homogenise the other properties.** Recently, Cheng et al. (2017) showed that **adopting an approach in which the frazil ice growth is calculated using a vertically-uniform frazil concentration** results in substantial underestimation of marine ice production **underneath the western side of Ronne Ice Shelf. Idealized one dimensional models confirm that the vertical distribution of frazil concentration cannot remain well-mixed in the upper layers of the ocean (Svensson and Omstedt, 1998) and beneath ice shelves (Holland and Feltham, 2005).** Consequently, earlier assessments of **either marine ice or platelet layer** production in the aforementioned areas may need to be reevaluated.

McMurdo Sound, located in the southwestern Ross Sea (Fig. 1), is characterized by significant ISW outflow, **arguably one of the most comprehensively observed ISW plumes available** (HU14; Langhorne et al., 2015; Robinson et al. 2014). A prominent platelet layer forms in the central-western sound (Dempsey et al., 2010); the maximum (area-averaged) observational first-year sea ice and platelet layer thickness are 2.5 (2) and 8 (3) m as determined from ice cores recovered adjacent to McMurdo Ice Shelf front between late November and early December in 2011 (Fig. 9 in HU14). The thin (~20 m) McMurdo Ice Shelf front allows the ISW outflow to be delivered to the ocean surface without mixing with warmer ambient waters (Robinson et al., 2014). The study documented in HU14 was the first to apply the steady, one-dimensional frazil-ice-laden ISW plume model developed by Smedsrud and Jenkins (2004) to McMurdo Sound, although a constant ISW plume thickness was used. McMurdo Sound therefore seems an ideal setting in which to apply and evaluate the new vertically-

modified ISW plume model proposed by Cheng et al. (2017), which includes time dependence and two horizontal dimensions. The main objective is to quantify for the first time the response of the platelet layer thickening rate to variations in ISW supercooling. Establishing such a relationship is of significance to the assessment of total sea ice thickness, and thus the oceanic heat flux associated with the platelet layer, in McMurdo Sound and elsewhere.

5

Here we first analyze the combined nonlinear effects of the vertical distributions of supercooling and frazil concentration on the suspended frazil ice growth rate in a supercooled ISW plume, and compare results with those obtained with a commonly-used, depth-averaged formulation. Then, we evaluate the performance of the vertically-modified ISW plume model in reproducing the observed ISW supercooling and platelet layer distribution to show the importance of considering the combined nonlinear effects. Finally, we conduct 211 sensitivity simulations with the purpose of quantitatively establishing the response of the platelet layer thickening rate **as well as the frazil ice concentration** to variations in ISW supercooling in McMurdo Sound.

10

## 2 Physically-based formulation for frazil ice growth rate

The growth rate of suspended frazil ice controls both the dynamic and thermodynamic evolution of ISW **plumes** and **the accretion of ice crystals beneath ice shelves** (Cheng et al., 2017; Holland and Feltham, 2006; Smedsrud and Jenkins, 2004) and sea ice (HU14). The frazil ice growth rate is found to be proportional to the following integral expression once a number of physical parameters within the commonly-used formulation of Jenkins and Bombosch (1995) are merged:

15

$$I_{gr} = \int_0^1 T_{SC} c_i(\sigma) d\sigma, \quad T_{SC} = T_f(\sigma, S) - T, \quad (1)$$

where  $\sigma \in [0, 1]$  is the relative vertical coordinate, with 0 and 1 respectively corresponding to the upper ice-plume and lower plume-ambient water interfaces,  $T$  and  $S$  are respectively the plume's temperature and salinity, vertically well-mixed within the plume,  $c_i$  is the vertically-distributed, in this study, volumetric frazil concentration within the plume,  $T_{SC}$  and  $T_f$  are respectively the supercooling level (positive for supercooling) and local freezing point. Because of the well-known linear decrease in  $T_f$  with increasing water depth,  $T_{SC}$  also varies linearly with depth, transitioning from supercooling to overheating as  $\sigma$  increases (Figs. 2a and 3). The corresponding transition height at which  $T_{SC} = 0$  is defined by supercooled thickness  $D_{SC} = \sigma_{SC} D$  where  $\sigma_{SC}$  and  $D$  are respectively supercooled fraction and total ISW plume thickness.

20

25

In earlier ISW plume models, because  $c_i$  is treated as vertically-uniform, the integral of (1) can be represented by the product of the depth-averaged values  $T_{SC}^{0.5}$  (0.5 means at mid-depth) and  $C_i$ . Thus, we refer to these ISW plume models as vertically-uniform (VU). It is worth mentioning that in order to take the supercooling into account when  $\sigma_{SC} < 0.5$ , HU14 integrated  $T_{SC}$  over the supercooled part only without introducing any frazil ice melting. However, in this study, we will demonstrate that the important role of frazil ice melting in the lower, overheated part of the plume cannot be ignored.

30

The vertical distribution of frazil concentration, in reality, much like the concentration of suspended sediment (Cheng et al., 2013, 2016), should be vertically non-uniform, with higher concentrations near the ice shelf/sea ice base. Considering only the balance between the buoyant-rise-induced vertical advection and turbulent diffusion terms, the governing equation for frazil concentration can be written as

$$\frac{d}{d\sigma} \frac{K}{D} \frac{dc_i}{d\sigma} + w_i \frac{dc_i}{d\sigma} = 0,$$

where  $w_i$  is the frazil ice rise velocity, determined by ice crystal size,  $K$  is the vertical frazil concentration diffusion coefficient, which can be parameterized as vertically constant (Cheng et al., 2013, 2016):

$$K = \frac{1}{6} \kappa u_* D,$$

where  $\kappa = 0.4$  is von Karman's constant,  $u_* = \sqrt{C_d} U$  is the friction velocity, related to the turbulent intensity within the ISW plume,  $C_d$  is the basal drag coefficient,  $U = \sqrt{(U_p + U_a)^2 + (V_p + V_a)^2 + U_t^2}$  is the total flow speed,  $U_p(U_a)$  and  $V_p(V_a)$  are the depth-averaged ISW plume (ambient current) speed in the  $x$  and  $y$  directions respectively,  $U_t$  is the root-mean square tidal speed. Using a zero net flux condition in the equilibrium state at the lower boundary of the plume, i.e.,

$$\frac{K}{D} \frac{dc_i}{d\sigma} + w_i c_i = 0, \text{ for } \sigma=1$$

and a Dirichlet boundary condition at the upper boundary, i.e.,

$$c_i = c_{i,b}, \text{ for } \sigma=0$$

where  $c_{i,b}$  is the frazil concentration at the ice shelf/sea ice base, the vertical exponential profile for the equilibrium frazil concentration can be readily obtained (Cheng et al., 2017):

$$\frac{c_i(\sigma)}{c_{i,b}} = \exp(-6Z_*\sigma),$$

where  $Z_* = w_i / \kappa u_*$  is the suspension index, otherwise known as the Rouse number. Integrating this exponential profile from  $\sigma=0$  to 1, we finally obtain the relation between  $c_i(\sigma)$  and  $C_i$  as

$$\frac{c_i(\sigma)}{C_i} = \frac{6Z_* \exp(-6Z_*\sigma)}{1 - \exp(-6Z_*)}. \quad (2)$$

As shown in Fig. 2a, the vertical distribution of frazil concentration is strongly controlled by  $Z_*$ . The gradient of the vertical distribution becomes greater with increasing  $Z_*$ , and a vertically-uniform frazil concentration distribution can only be achieved as  $Z_*$  approaches 0. Accordingly, Cheng et al. (2017) introduced (2) into (1), and as a result significantly improved the simulated pattern of marine ice growth under the western side of Ronne Ice Shelf, compared with the VU and satellite-derived (Joughin and Padman, 2003) results. Hereinafter, we refer to this vertically-modified ISW plume model as VM. To conclude, the only difference between VM and VU models is whether the vertical distribution of frazil ice concentration is introduced.

The dependence of the integral value of  $I_{gr}$  on  $Z_*$  under specified conditions of supercooling (Fig. 2a) is shown in Fig. 2b, where  $D_{SC} = 50$  m (a value within the calculated range for the standard run, Fig. 1) in all the cases. It can be seen that the integral value increases nonlinearly with  $Z_*$ . The critical  $Z_*$  that represents the transition from frazil ice melting ( $I_{gr} < 0$ ) to freezing ( $I_{gr} > 0$ ) decreases as the supercooled part of ISW plume increases. In contrast, owing to the neglect of vertical variation in  $c_i$ , the integral values calculated using the VU formulation are constant, leading to transitions from overestimation of frazil ice growth to underestimation, compared with VM, as  $Z_*$  increases. Only if the ISW plume is fully supercooled ( $\sigma_{SC} = 1$ ) and  $Z_*$  is close to 0 are the integral values of  $I_{gr}$  calculated by VU and VM formulations equal (star in Fig. 2b). These features are illustrated in Fig. 2a: for given supercooling, if  $Z_*$  becomes larger, there is higher (lower) frazil concentration in the upper (lower), supercooled (overheated) part of the ISW plume. Owing to the assumption that thermohaline exchanges between frazil crystals and ambient water occur only at the crystal edge for freezing, but over the whole crystal surface for melting (Jenkins and Bombosch, 1995), the integral values of  $I_{gr}$  for the lower overheated part can be of much greater magnitude (Fig. 2b). It is therefore necessary to limit the mass loss due to frazil melting in one model time step such that it does not exceed the frazil concentration in the lower, overheated part of the plume. Overall, the frazil concentration and frazil growth rate distributions in the VM model show physically-reasonable and desirable characteristics that are absent from the VU model, and the impacts will be demonstrated by evaluation of the VM model in McMurdo Sound.

### 3 ISW model in McMurdo Sound

The unsteady VM and VU models used in this study are described in detail by Cheng et al. (2017). The governing equations for ISW properties and frazil concentration in both VM and VU models remain as they were in the depth-integrated, two-dimensional ISW plume model developed by Holland and Feltham (2006), except for the different treatments of the specific terms associated with the frazil ice growth rate, described above, in the frazil concentration and potential temperature transport equations of the VM model. Both VM and VU models combine the same commonly-used parameterizations of thermohaline exchanges across the ice–water interfaces, specifically a three-equation formulation (Holland and Jenkins, 1999) for the sea ice base and a two-equation formulation for frazil ice (Galton-Fenzi et al., 2012), with a multiple size–class frazil dynamics model (Smedsrud and Jenkins, 2004), to calculate basal freezing ( $f'$ ) and frazil melting/freezing ( $w'$ ), secondary nucleation ( $N'$ ), and precipitation ( $p'$ ). These processes are summarized in Fig. 3. Rather than repeat all the equations here, we recall some of them and present how we set up our ISW plume models on the McMurdo Sound domain.

The model domain (Fig. 1) is delimited by a 45×40 km rectangle in the  $x$ - $y$  plane with an ISW outflow from beneath McMurdo Ice Shelf. The base of the sea ice in McMurdo Sound is assumed to be horizontal and rough, owing to the presence of the platelet layer. The drag coefficient of the ice underside is therefore 6-30 times larger than that typically

applied in ice-ocean interaction models (Robinson et al, 2017). The parameterization of the sea ice thermodynamics, the assumption of no entrainment of ambient water into the ISW plume, and the boundary conditions at the ISW outflow follow HU14. The initial thickness of the ISW outflow from underneath McMurdo Ice Shelf is set equal to that of the supercooled layer, i.e.,  $D = D_{SC}$ , and the discharge per unit width is set to  $0.02 \text{ m}^2 \text{ s}^{-1}$ . The addition of both an ambient circulation and tides follow HU14: the former, which represented the only source of momentum in the study of HU14, is assumed to be parallel to the Victoria Land coast, in the negative  $y$  direction, and to be constant throughout the model domain; the latter is calculated using root-mean square tidal speeds from Padman and Erofeeva (2005). Because ISW persists in McMurdo Sound for at least the 8-9 months of the ice growth season (Robinson et al., 2014), all runs are integrated for 240 days. The model resolution and time step ( $\Delta t$ ) are 1 km and 25 s, respectively. The frazil ice size distribution is represented by 5 crystal size classes, and the transfer processes, induced by frazil freezing and melting, between different size classes are calculated using the scheme proposed by Smedsrud and Jenkins (2004). The ice concentration at the ISW outflow is evenly distributed among the classes (Holland and Feltham, 2005, 2006; Smedsrud and Jenkins, 2004).

We treat the frazil ice precipitation rate  $p'$  as inverted sedimentation and follow the parameterization of McCave and Swift (1976):

$$p' = w_i C_i \left(1 - \frac{U^2}{U_c^2}\right) \times He \left(1 - \frac{U^2}{U_c^2}\right), \quad (3)$$

where  $U_c$  is a critical velocity, above which precipitation cannot occur, determined by Jenkins and Bombosch (1995):

$$U_c^2 = \frac{\theta_i(\rho_0 - \rho_i)g^2 r_e}{\rho_0 C_D},$$

where  $\theta_i$  is the Shields criterion,  $\rho_0$  and  $\rho_i$  are reference seawater and ice densities, respectively,  $g$  is gravity,  $r_e$  is the equivalent radius of a sphere with the same volume as the frazil disk. The frazil ice rise velocity,  $w_i$ , is calculated by Morse and Richard (2009):

$$w_i = \begin{cases} 2.025 D_i^{1.621} & \text{if } D_i \leq 1.27 \text{ mm} \\ -0.103 D_i^2 + 4.069 D_i - 2.024 & \text{if } 1.27 < D_i \leq 7 \text{ mm} \end{cases}$$

where  $D_i = 2r_i$  is the diameter of a frazil crystal in mm. The inclusion of the Heaviside function  $He$  means that negative precipitation (i.e., erosion of previously deposited frazil ice) is not permitted. Because we have no idea about how cohesive the ice crystals are once they have settled, the estimation of an erosion rate would entail additional uncertainties.

The complex processes after the frazil ice precipitates onto the sea ice base are simplified in our model. In order to calculate platelet layer thickness  $D_p$  at the  $n^{\text{th}}$  time interval, we adopt the assumptions of HU14 that solid ice fraction within the platelet layer in McMurdo Sound is 0.25 based on the observational estimation from Gough et al. (2012) and that the ice crystals, on average, double in volume after precipitation:

$$D_p = \frac{1}{0.25} \times 2 \times \sum_{k=1}^n (p'_k \times \Delta t).$$

It should be noted that the volume change factor is a broad estimate, with almost no supporting evidence in the literature to guide it.

## 4 Results

### 4.1 Standard model run

- 5 The performance of the VU and VM models in reproducing the ISW supercooling and platelet layer pattern in McMurdo Sound is evaluated by comparing results with observational data. To our knowledge, the data reported by HU14 are the most comprehensive available to evaluate our model, including both oceanographic and ice core data in two horizontal dimensions adjacent to McMurdo Ice Shelf. As this study represents the first application of a two-dimensional ISW plume model to the McMurdo Sound region, extensive tuning of the least constrained model parameters, including the ISW outflow properties, platelet layer basal drag coefficient, frazil ice crystal size distribution, ambient current speed, and Shields criterion was required to produce the distributions of ISW properties and platelet layer thickness shown in Figs. 4a and 6, respectively. Table 1 gives values adopted for the key parameters. Model results are evaluated by means of skill metrics: Root-Mean-Square Error (RMSE), Correlation Coefficient (CC), and Skill Score (SS), respectively given by

$$RMSE = \left[ \frac{\sum (X_{cal} - X_{obs})^2}{M} \right]^{1/2},$$

$$15 \quad CC = \frac{\sum (X_{cal} - \bar{X}_{cal})(X_{obs} - \bar{X}_{obs})}{[\sum (X_{cal} - \bar{X}_{cal})^2 \sum (X_{obs} - \bar{X}_{obs})^2]^{1/2}},$$

$$SS = 1 - \frac{\sum (X_{cal} - X_{obs})^2}{\sum (X_{obs} - \bar{X}_{obs})^2},$$

where  $X$  is the variable being evaluated,  $M$  is the number of data points, and the overbar denotes the arithmetic mean. The performance of each model is indicated by SS as: >0.65 excellent; 0.65–0.5 very good; 0.5–0.2 good; <0.2 poor (Luo et al., 2017; Ralston et al., 2010; Song and Wang, 2013).

20

- It can be seen that at the end of the simulations both VM and VU models reproduce the observed reduction in ISW supercooling at the sea ice base ( $T_{SC}^0$ , superscript “0” denotes the sea ice base) in the cross- and long-sound directions, in spite of some evident model discrepancies (Fig. 4a) that may result from the limitations in our model setup: both the ambient current and tides are treated as temporally and spatially constant; there are no long-term observations of ISW outflow to provide reliable boundary conditions; we use a constant drag coefficient, ignoring the spatiotemporal evolution of the sea ice basal form characterized by the platelet layer. Nevertheless, the SS of  $T_{SC}^0$  calculated using VM and VU models are 0.56 and 0.58, respectively, and the CC and RMSE are also reasonable (Table 2). There are only small differences throughout the time series of  $T_{SC}^0$  simulated by the VM and VU models (Fig. 4) and the final distributions of both total ISW plume thickness and supercooled thickness are also very similar (see Fig. 5a-d). A comprehensive comparison of  $T_{SC}^0$  calculated by the VM and

VU models in an extensive set of sensitivity experiments will be discussed later. Finally, it can be seen that in both models the ISW plume flow is predominantly governed by a geostrophic balance (Fig 5a-d).

In contrast, the frazil concentration (red lines in Fig. 4b, Fig. 5e and f) and platelet layer thickness (green lines in Fig. 4b, Fig. 6b and c) are both underestimated by the VU model, compared with the results of the VM model, throughout the time series. Given the small differences in  $T_{SC}^0$  calculated by VM and VU models, this result demonstrates that the vertical distribution of frazil concentration within the ISW plume plays a critical role in determining the suspended frazil ice growth (Fig. 2), and thus the frazil concentration and platelet layer thickness distributions. The supercooling is utilized more efficiently in the VM model, giving a greater depth-averaged frazil concentration than is produced by the commonly-used VU model. Because the sea ice base is horizontal, there are no changes in the freezing point associated with pressure change, so supercooling is always highest at the ISW outflow (Fig. 5c and d). That results in the greatest frazil concentration (Fig. 5f) and platelet layer thickness (Fig. 6b) near the location of the outflow, with large changes from  $2.2 \times 10^{-4}$  to  $4 \times 10^{-5}$  and 15 to 6 m, respectively, in the case of the VM model within 5 km of the outflow. Such small-scale features in the platelet layer thickness distribution, if present, were not resolved by the relatively coarse spatial distribution of ice-core sampling (dots in Fig. 6). Nevertheless, the platelet layer thickness calculated by the VM model at drill sites agrees well with the measurements (Fig. 6a), being graded “excellent” in contrast with the “poor” performance of the VU model (Table 2). Despite efforts to tune the VU model to give a better match with the observed platelet layer thickness, even a limited expansion of the platelet layer can only be achieved with a considerable increase in the calculated  $T_{SC}^0$ , in disagreement with the observations.

For both VM and VU models, the time series of area-averaged  $T_{SC}^0$ ,  $C_i$  (hereafter  $T_{SC}^0$  and  $C_i$  denote their area-average values), and platelet layer thickness indicate respectively two near-constant values and one near-constant growth rate after about the 150<sup>th</sup> day (Fig. 4b). It is informative to explore how our various assumptions about the vertical distribution of frazil concentration influence the steady-state relationship between those variables in the McMurdo Sound region.

#### 4.2 Dependence of platelet layer thickening rate on ISW supercooling

The response of ice shelf basal melting to variations in ocean temperature has been investigated using satellite altimetry (Rignot and Jacobs, 2002; Shepherd et al., 2004) and numerical models (Grosfeld and Sandhäger, 2004; Holland et al., 2008; Payne et al., 2007; Walker and Holland, 2007; Williams et al., 1998, 2002). In contrast, we know of no studies to date of the response of marine ice (or platelet layer) thickening rate beneath ice shelves (or sea ice) to variations in supercooling, which is of potential significance for evaluating the mass balance of deep-draughting ice shelves in cold water environments and adjacent sea ice subject to climatic variability.

Owing to the number of poorly-constrained parameters in the frazil-ice-laden ISW plume model, we conducted 211 comparative sensitivity experiments between VM and VU models, varying both physical and input parameters, including



drag coefficient, frazil ice crystal size configuration, average number of frazil crystals, ambient current speed, width and thickness of the ISW outflow, and frazil concentration within the outflow (see Table 3). For all model runs, we plot the relationship between  $T_{SC}^0$  and thickening rate in the steady state, using output from the last 30 days of each run (Fig. 7).

- 5 In Fig. 7a, the results of the VM model are grouped by the prescribed supercooled layer thickness  $D_{SC}^{ini}$  in the ISW outflow. For  $D_{SC}^{ini} < 65$  m there is a relatively consistent increase in thickening rate with increasing  $T_{SC}^0$ , while for  $D_{SC}^{ini} \geq 65$  m the thickening rate tends to be much more variable. It is worth mentioning that  $D_{SC}^{ini} = 65$  m is the value estimated by HU14 based on the measurements conducted by Lewis and Perkin (1985) and Jones and Hill (2001). For  $D_{SC}^{ini} = 78$  m and greater, inflexions emerge separating a region of low thickening rate, where the thickening rate tends to decrease with increasing  $T_{SC}^0$ , from a region of high thickening rate, where there is a very rapid increase in thickening rate with increasing  $T_{SC}^0$ . This complex response of the VM model must result from the consideration of vertical structure in the frazil concentration, controlled by the frazil ice suspension index  $Z_*$  (Fig. 2), in the calculation of frazil ice growth.

We therefore calculated the weighted-average of  $Z_*$  at each grid point for the VM model using the following equation:

15 
$$\bar{Z}_* = \frac{\sum_{k=1}^n C_i^k Z_*^k}{\sum_{k=1}^n C_i^k} = \frac{\sum_{k=1}^n C_i^k Z_*^k}{C_i},$$

- where  $C_i^k$  is the frazil concentration of the  $k^{th}$  size class, and  $n$  is the number of size classes used. Then, we took the average of  $\bar{Z}_*$  over all the grid points occupied by the plume to give a representative suspension index for the VM runs (hereinafter  $\bar{Z}_*$  denotes its area-averaged value). We replotted the VM model results characterized by  $\bar{Z}_*$  in Fig. 7b. We find systematic changes in  $\bar{Z}_*$  with increasing thickening rate (along the coloured lines in Fig. 7b), particularly for  $D_{SC}^{ini} \geq 78$  m where the inflexions emerge. With decreasing  $\bar{Z}_*$ ,  $T_{SC}^0$  first decreases, and then increases. If  $\bar{Z}_*$  is sufficiently large, the suspended frazil crystals deposit out of the ISW plume so rapidly that they cannot efficiently use the ISW supercooling to grow, leading to the smallest platelet layer production for the VM model. For smaller  $\bar{Z}_*$ , the frazil crystals bathed in the supercooled layer of the ISW plume can remain in suspension and grow longer, resulting in a thicker platelet layer and less residual supercooling. However, if  $\bar{Z}_*$  decreases further, higher frazil concentration occurs within the lower, overheated part of the ISW plume, where melting of the crystals can mitigate the release of latent heat (Fig. 2b). That promotes further growth of frazil ice which can remain in suspension even longer, and thus lead to rapid platelet layer production. The thickening rate calculated by the VU model is also shown, and is discernibly smaller than that calculated by the VM model. In addition, the maximum values of  $T_{SC}^0$  were obtained within the VU model, because the supercooling is used less efficiently for producing the platelet layer in the VU than in the corresponding VM runs.

30

These arguments can be further illustrated by a more detailed comparison of  $T_{SC}^0$  calculated by the VM and VU models (Fig. 8). There are a number of runs, including the standard run, that have larger  $T_{SC}^0$  values in the VM than in the VU model. The trend from larger  $T_{SC}^0$  in the VM model to larger  $T_{SC}^0$  in the VU model is accompanied by increases in  $\bar{Z}_*$ . When  $\bar{Z}_*$  is relatively small, large frazil concentration exists within the lower overheated part of the ISW plume (Fig. 2b) where melting of frazil ice (causing cooling) counteracts the consumption of supercooling by frazil growth (causing warming) in the upper part of the plume. As  $\bar{Z}_*$  increases, the frazil concentration within the lower overheated part decreases, and finally vanishes, and the resulting release of supercooling in the upper part is more efficient in the VM model, giving larger  $T_{SC}^0$  values in the VU model.

In Fig. 7a, when  $D_{SC}^{ini} < 65$  m, ISW supercooling is insufficient to distinguish runs with different  $\bar{Z}_*$ . In other words, the relation between thickening rate and  $T_{SC}^0$  is independent of  $\bar{Z}_*$  for such small  $D_{SC}^{ini}$ . When  $D_{SC}^{ini}$  is within the range of 65 to 78 m, the VM model results are distinguishable, with data points having smaller thickening rate and larger  $T_{SC}^0$  corresponding to larger  $\bar{Z}_*$  (Fig. 7b). When  $D_{SC}^{ini} \geq 78$  m, the inflexions emerge, and the ISW supercooling revives when  $\bar{Z}_*$  decreases further. Therefore, we conclude that when  $D_{SC}^{ini}$  exceeds a critical value (about 65 m for these McMurdo Sound simulations), the efficiency of converting ISW supercooling into frazil ice growth is controlled by the suspension index.

### 4.3 Dependence of frazil concentration on ISW supercooling

In view of the correlation between platelet layer thickening rate and frazil concentration shown in Eq. (3) (also see Fig. 5e and f; Fig. 6b and c), we will explore the relationship between  $T_{SC}^0$  and  $C_i$  here. As expected, the complex response of  $C_i$  to variations in  $T_{SC}^0$  (Fig. 9) is similar to the relationship between  $T_{SC}^0$  and thickening rate (Fig. 7) in the VM model.

20

The magnitude of the difference in  $C_i$  calculated by VM and VU models (VM minus VU) is compared in Fig. 10a, where we find that  $C_i$  calculated by the VM model is always larger than that calculated by the VU model. In general, the difference increases with decreasing  $\bar{Z}_*$ , while the sensitivity grows with increasing  $D_{SC}^{ini}$ . The dependence on  $\bar{Z}_*$  is once again due to the impact of the combined thermodynamic processes, i.e., the efficient growth in the upper supercooled part of the plume together with the maintenance of supercooling by melting of frazil in the lower part, discussed above. We also see similar behavior for the difference in the thickening rate (Fig. 10b).

25

Fig. 7 (Fig. 9) suggests a possible relationship between platelet layer thickening rate (frazil concentration) and supercooling in McMurdo Sound, but observations of suspended frazil ice crystal sizes and turbulence within the ISW would be needed to calculate a representative suspension index. To date, there are limited observations of frazil ice in situ, and the majority of the observations made use of instruments not specifically designed for ice crystal detection (Leonard et al., 2006).

30

## 5 Summary and future works

In this study, we demonstrated how the vertical distributions of supercooling and frazil ice concentration within an ISW plume jointly determine the growth of suspended frazil ice, and thus the rate of platelet layer formation [under sea ice and marine ice beneath ice shelves](#). A vertically-modified, frazil-ice-laden, ISW plume model which encapsulates these combined nonlinear effects was applied to the McMurdo Sound region, and reproduced the observed ISW supercooling and platelet layer distributions in two horizontal dimensions. Using multiple model runs, the relationship of ISW supercooling to platelet layer thickening rate and frazil concentration in McMurdo Sound was explored, [and shown to be dependent](#) on the suspension index that controls the vertical distribution of frazil concentration within the ISW plume. Moreover, when the thickness of a supercooled layer of ISW is large enough, the efficiency of converting ISW supercooling into frazil concentration, and thus platelet layer growth is determined by the suspension index. [These findings highlight the need for further](#) observations in McMurdo Sound, particularly focused near the ISW outflow region in the western sound, where the supercooled ISW plume and platelet layer are prominent, and more general observations that help to constrain the frazil size spectrum within the sea ice-ocean boundary layer. [In addition](#), the performance of the VM model in providing reliable estimates of supercooling and frazil ice flux at the platelet layer base makes it an attractive tool for coupling with sea ice models focusing on microscale processes within the bottom layer of the ice (Buffo et al., 2018).

[It would be straightforward for the next step to investigate the relationship between supercooling and marine ice thickening rate underneath ice shelves using the VM model. Quantifying this relationship would be the key to parameterizing the process in more complex three-dimensional, primitive equation ocean models, which frequently neglect details of the ice shelf-ocean boundary layer and processes associated with an evolving suspension of frazil ice crystals \(Liu et al., 2017; Mueller et al., 2012, 2018\). The main difference between the process of marine ice accretion beneath ice shelves and the growth of the sub-ice platelet layer discussed here is likely to be the magnitude of the vertical heat flux that the deposited frazil ice is subjected to. At the base of an ice shelf, typically several hundred meters thick, the vertical temperature gradient is comparatively small, so the deposited crystals form a slushy layer \(Engelhardt and Determann, 1987\) that slowly consolidates, possibly as much through compaction as freezing. The ice-ocean interface and the associated drag coefficient are therefore likely to be very different to those observed in McMurdo Sound, where the platelet layer appears to comprise a more open matrix of ice and water that consolidates by freezing as heat is lost to the atmosphere. Therefore, the VM model would need to be re-evaluated against observations of sub-ice shelf ISW plumes and the ice shelf-ocean boundary layer. Finally, further process studies, including the influence of the vertical current structure within either the ice shelf or sea ice - ocean boundary layer \(Jenkins, 2016; Robinson et al., 2017\) could also contribute to improving our understanding of marine ice and platelet layer formation.](#)

*Data availability.* The data archive associated with this study can be found in the Global Change Master Directory under the keyword K063\_2011\_2012\_NZ\_1.

*Author contributions.* CC led the study. The simulations were designed by ZW and CC, implemented by CL and RX, and analyzed by CC, AJ, and PRH. The paper was written by CC, AJ, and PRH.

5 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We would like to thank two anonymous referees and Ken Hughes for their thorough review and helpful comments and suggestions. This work was funded by the National Natural Science Foundation of China (41406214, 41306208, 41606217). CC and CL were respectively supported by the China Scholarship Council (201708320046, 201504180026). ZW was supported by “the Fundamental Research Funds for the Central Universities” (2017B04814, 2017B20714).

## References

- Arrigo, K. R., Mock, T., and Lizotte, M. P.: Primary producers and sea ice, *Sea ice*, 2, 283-325, 2010.
- Bombosch, A. and Jenkins, A.: Modeling the formation and deposition of frazil ice beneath Filchner-Ronne Ice Shelf, *J. Geophys. Res.-Oceans*, 100, 6983-6992, <http://dx.doi.org/10.1029/94JC03224>, 1995.
- 15 Buffo, J. J., Schmidt, B. E., and Huber, C.: Multiphase Reactive Transport and Platelet Ice Accretion in the Sea Ice of McMurdo Sound, Antarctica, *J. Geophys. Res.-Oceans*, 123, 324-345, <https://doi.org/10.1002/2017JC013345>, 2018.
- Cheng, C., Song, Z. Y., Wang, Y. G., and Zhang, J. S.: Parameterized expressions for an improved Rouse equation, *Int. J. Sediment Res.*, 28, 523-534, [http://dx.doi.org/10.1016/S1001-6279\(14\)60010-X](http://dx.doi.org/10.1016/S1001-6279(14)60010-X), 2013.
- Cheng, C., Huang, H., Liu, C., and Jiang, W.: Challenges to the representation of suspended sediment transfer using a depth-averaged flux, *Earth Surf. Proc. Land*, 41, 1337-1357, <http://dx.doi.org/10.1002/esp.3903>, 2016.
- 20 Cheng, C., Wang, Z., Liu, C., and Xia, R.: Vertical Modification on Depth-Integrated Ice Shelf Water Plume Modeling Based on an Equilibrium Vertical Profile of Suspended Frazil Ice Concentration, *J. Phys. Oceanogr.*, 47, 2773-2792, <http://dx.doi.org/10.1175/JPO-D-17-0092.1>, 2017.
- Dempsey, D. E., Langhorne, P. J., Robinson, N. J., Williams, M. J. M., Haskell, T. G., and Frew, R. D.: Observation and modeling of platelet ice fabric in McMurdo Sound, Antarctica, *J. Geophys. Res.-Oceans*, 115, <https://doi.org/10.1029/2008JC005264>, 2010.
- 25 Engelhardt, H. and Determann, J.: Borehole evidence for a thick layer of basal ice in the central Ronne Ice Shelf, *Nature*, 327, 318-319, 1987.
- Fricker, H. A., Popov, S., Allison, I., and Young, N.: Distribution of marine ice beneath the Amery Ice Shelf, *Geophys. Res. Lett.*, 28, 2241-2244, <https://doi.org/10.1029/2000GL012461>, 2001.
- 30

- Galton-Fenzi, B. K., Hunter, J. R., Coleman, R., Marsland, S. J., and Warner, R. C.: Modeling the basal melting and marine ice accretion of the Amery Ice Shelf, *J. Geophys. Res.-Oceans*, 117, <http://dx.doi.org/10.1029/2012JC008214>, 2012.
- Gough, A. J., Mahoney, A. R., Langhorne, P. J., Williams, M. J., Robinson, N. J., and Haskell, T. G.: Signatures of supercooling: McMurdo Sound platelet ice, *J. Glaciol.*, 58, 38-50, <https://doi.org/10.3189/2012JoG10J218>, 2012.
- 5 Grosfeld, K. and Sandhäger, H.: The evolution of a coupled ice shelf–ocean system under different climate states, *Global Planet. Change*, 42, 107-132, <http://dx.doi.org/10.1016/j.gloplacha.2003.11.004>, 2004.
- Herraiz-Borreguero, L., Allison, I., Craven, M., Nicholls, K. W., and Rosenberg, M. A.: Ice shelf/ocean interactions under the Amery Ice Shelf: Seasonal variability and its effect on marine ice formation, *J. Geophys. Res.-Oceans*, 118, 7117-7131, <https://doi.org/10.1002/2013JC009158>, 2013.
- 10 Holland, P. R. and Feltham, D. L.: Frazil dynamics and precipitation in a water column with depth-dependent supercooling, *J. Fluid Mech.*, 530, 101-124, <https://doi.org/10.1017/S002211200400285X>, 2005.
- Holland, P. R. and Feltham, D. L.: The effects of rotation and ice shelf topography on frazil-laden ice shelf water plumes, *J. Phys. Oceanogr.*, 36, 2312-2327, <https://doi.org/10.1175/JPO2970.1>, 2006.
- Holland, P. R., Feltham, D. L., and Jenkins, A.: Ice shelf water plume flow beneath Filchner-Ronne Ice Shelf, Antarctica, *J. Geophys. Res.-Oceans*, 112, <https://doi.org/10.1029/2006JC003915>, 2007.
- 15 Holland, P. R., Jenkins, A., and Holland, D. M.: The response of ice shelf basal melting to variations in ocean temperature, *J. Climate*, 21, 2558-2572, <https://doi.org/10.1175/2007JCLI1909.1>, 2008.
- Holland, P. R., Corr, H. F., Vaughan, D. G., Jenkins, A., and Skvarca, P.: Marine ice in Larsen ice shelf, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2009GL038162>, 2009.
- 20 Hughes, K. G., Langhorne, P. J., Leonard, G. H., and Stevens, C. L.: Extension of an Ice Shelf Water plume model beneath sea ice with application in McMurdo Sound, Antarctica, *J. Geophys. Res.-Oceans*, 119, 8662-8687, <https://doi.org/10.1002/2013JC009411>, 2014.
- Hunkeler, P. A., Hoppmann, M., Hendricks, S., Kalscheuer, T., and Gerdes, R.: A glimpse beneath Antarctic sea ice: Platelet layer volume from multifrequency electromagnetic induction sounding, *Geophys. Res. Lett.*, 43, 222-231, <https://doi.org/10.1002/2015GL065074>, 2016.
- 25 Jenkins, A.: A simple model of the ice shelf–ocean boundary layer and current, *J. Phys. Oceanogr.*, 46, 1785-1803, <http://dx.doi.org/10.1175/JPO-D-15-0194.1>, 2016.
- Jenkins, A. and Bombosch, A.: Modeling the effects of frazil ice crystals on the dynamics and thermodynamics of ice shelf water plumes, *J. Geophys. Res.-Oceans*, 100, 6967-6981, <http://dx.doi.org/10.1029/94JC03227>, 1995.
- 30 Jones, S. J. and Hill, B. T.: Structure of sea ice in McMurdo Sound, Antarctica, *Ann. Glaciol.*, 33, 5-12, <http://dx.doi.org/10.3189/172756401781818347>, 2001.
- Joughin, I. and Padman, L.: Melting and freezing beneath Filchner-Ronne Ice Shelf, Antarctica, *Geophys. Res. Lett.*, 30, <http://dx.doi.org/10.1029/2003GL016941>, 2003.

- Langhorne, P. J., Hughes, K. G., Gough, A. J., Smith, I. J., Williams, M. J. M., Robinson, N. J., Stevens, C. L., Rack, W., Price, D., Leonard, G. H., Mahoney, A. R., Haas, C., Haskell, T. G.: Observed platelet ice distributions in Antarctic sea ice: An index for ocean-ice shelf heat flux, *Geophys. Res. Lett.*, 42, 5442-5451, <https://doi.org/10.1002/2015GL064508>, 2015.
- 5 Lewis, E. L. and Perkin, R. G.: The winter oceanography of McMurdo Sound, Antarctica, *Oceanology of the Antarctic continental shelf*, 145-165, <https://doi.org/10.1029/AR043p0145>, 1985.
- Leonard, G. H., Purdie, C. R., Langhorne, P. J., Haskell, T. G., Williams, M. J. M., and Frew, R. D.: Observations of platelet ice growth and oceanographic conditions during the winter of 2003 in McMurdo Sound, Antarctica, *J. Geophys. Res.-Oceans*, 111, <https://doi.org/10.1029/2005JC002952>, 2006.
- 10 Liu, C., Wang, Z., Cheng, C., Xia, R., Li, B., and Xie, Z.: Modeling modified Circumpolar Deep Water intrusions onto the Prydz Bay continental shelf, East Antarctica, *J. Geophys. Res.-Oceans*, 122, 5198-5217, <http://dx.doi.org/10.1002/2016JC012336>, 2017.
- Luo, Z., Zhu, J., Wu, H., and Li, X.: Dynamics of the sediment plume over the Yangtze Bank in the Yellow and East China Seas, *J. Geophys. Res.-Oceans*, 122, 10073-10090, <https://doi.org/10.1002/2017JC013215>, 2017.
- 15 McCave, I. N. and Swift, S. A.: A physical model for the rate of deposition of fine-grained sediments in the deep sea, *Geological Society of America Bulletin*, 87, 541-546, [https://doi.org/10.1130/0016-7606\(1976\)87<541:APMFTR>2.0.CO;2](https://doi.org/10.1130/0016-7606(1976)87<541:APMFTR>2.0.CO;2), 1976.
- Morgan, V. I.: Oxygen isotope evidence for bottom freezing on the Amery Ice Shelf, *Nature*, 238, 393, <https://doi.org/10.1038/238393a0>, 1972.
- 20 Morse, B. and Richard, M.: A field study of suspended frazil ice particles, *Cold Reg. Sci. Technol.*, 55, 86-102, <https://doi.org/10.1016/j.coldregions.2008.03.004>, 2009.
- Mueller, R. D., Hattermann, T., Howard, S. L., and Padman, L.: Tidal influences on a future evolution of the Filchner–Ronne Ice Shelf cavity in the Weddell Sea, Antarctica, *The Cryosphere*, 12, 453, <http://dx.doi.org/10.5194/tc-12-453-2018>, 2018.
- 25 Mueller, R. D., Padman, L., Dinniman, M. S., Erofeeva, S. Y., Fricker, H. A., and King, M. A.: (2012), Impact of tide-topography interactions on basal melting of Larsen C Ice Shelf, Antarctica, *J. Geophys. Res.-Oceans*, 117, C05005, <https://doi.org/10.1029/2011JC007263>, 2012.
- Oerter, H., Kipfstuhl, J., Determann, J., Miller, H., Wagenbach, D., Minikin, A., and Graft, W.: Evidence for basal marine ice in the Filchner–Ronne Ice Shelf, *Nature*, 358, 399, <https://doi.org/10.1038/358399a0>, 1992.
- 30 Padman, L. and Erofeeva, S.: Tide Model Driver (TMD) Manual, Version 1.2, Earth and Space Research, Seattle, Wash., 2005. [Available at [www.esr.org/polar\\_tide\\_models/README\\_TMD.pdf](http://www.esr.org/polar_tide_models/README_TMD.pdf), last accessed 7 Nov. 2012.]
- Payne, A. J., Holland, P. R., Shepherd, A. P., Rutt, I. C., Jenkins, A., and Joughin, I.: Numerical modeling of ocean-ice interactions under Pine Island Bay's ice shelf, *J. Geophys. Res.-Oceans*, 112, <http://dx.doi.org/10.1029/2006JC003733>, 2007.

- Ralston, D. K., Geyer, W. R., and Lerczak, J. A.: Structure, variability, and salt flux in a strongly forced salt wedge estuary, *J. Geophys. Res.-Oceans*, 115, C06005, <https://doi.org/10.1029/2009JC005806>, 2010.
- Rees Jones, D. W. and Wells, A. J.: Frazil-ice growth rate and dynamics in mixed layers and sub-ice-shelf plumes, *The Cryosphere*, 12, 25-38, <http://dx.doi.org/10.5194/tc-12-25-2018>, 2018.
- 5 Rignot, E. and Jacobs, S. S.: Rapid bottom melting widespread near Antarctic ice sheet grounding lines, *Science*, 296, 2020-2023, <http://dx.doi.org/10.1126/science.1070942>, 2002.
- Rignot, E., Jacobs, S., Mouginot, J., and Scheuchl, B.: Ice-shelf melting around Antarctica, *Science*, 341, 266-270, <https://doi.org/10.1126/science.1235798>, 2013.
- Robinson, N. J., Stevens, C. L., and McPhee, M. G.: Observations of amplified roughness from crystal accretion in the sub-ice ocean boundary layer, *Geophys. Res. Lett.*, 44, 1814-1822, <https://doi.org/10.1002/2016GL071491>, 2017.
- 10 Robinson, N. J., Williams, M. J., Stevens, C. L., Langhorne, P. J., and Haskell, T. G.: Evolution of a supercooled Ice Shelf Water plume with an actively growing subice platelet matrix, *J. Geophys. Res.-Oceans*, 119, 3425-3446, <https://doi.org/10.1002/2013JC009399>, 2014.
- Shepherd, A., Wingham, D., and Rignot, E.: Warm ocean is eroding West Antarctic ice sheet, *Geophys. Res. Lett.*, 31, <http://dx.doi.org/10.1029/2004GL021106>, 2004.
- 15 Smedsrud, L. H. and Jenkins, A.: Frazil ice formation in an ice shelf water plume, *J. Geophys. Res.-Oceans*, 109, <http://dx.doi.org/10.1029/2003JC001851>, 2004.
- Smith, I. J., Langhorne, P. J., Haskell, T. G., Trodahl, H. J., Frew, R., and Vennell, M. R.: Platelet ice and the land-fast sea ice of McMurdo Sound, Antarctica, *Ann. Glaciol.*, 33, 21-27, <http://dx.doi.org/10.3189/172756401781818365>, 2001.
- 20 Song, D. and Wang, X. H.: Suspended sediment transport in the Deepwater Navigation Channel, Yangtze River Estuary, China, in the dry season 2009: 2. Numerical simulations, *J. Geophys. Res.-Oceans*, 118, 5568-5590, <http://dx.doi.org/10.1002/jgrc.20411>, 2013.
- Stern, A. A., Dinniman, M. S., Zagorodnov, V., Tyler, S. W., and Holland, D. M.: Intrusion of warm surface water beneath the McMurdo Ice Shelf, Antarctica, *J. Geophys. Res.-Oceans*, 118, 7036-7048, <https://doi.org/10.1002/2013JC008842>, 2013.
- 25 Svensson, U. and Omstedt, A.: Numerical simulations of frazil ice dynamics in the upper layers of the ocean, *Cold Reg. Sci. Technol.*, 28, 29-44. [http://dx.doi.org/10.1016/S0165-232X\(98\)00011-1](http://dx.doi.org/10.1016/S0165-232X(98)00011-1), 1998.
- Walker, R. T. and Holland, D. M.: A two-dimensional coupled model for ice shelf-ocean interaction, *Ocean Model.*, 17, 123-139, <http://dx.doi.org/10.1016/j.ocemod.2007.01.001>, 2007.
- 30 Williams, M. J. M., Warner, R. C., and Budd, W. F.: The effects of ocean warming on melting and ocean circulation under the Amery Ice Shelf, East Antarctica, *Ann. Glaciol.*, 27, 75-80, <http://dx.doi.org/10.3189/1998AoG27-1-75-80>, 1998.
- Williams, M. J. M., Warner, R. C., and Budd, W. F.: Sensitivity of the Amery Ice Shelf, Antarctica, to changes in the climate of the Southern Ocean, *J. Climate*, 15, 2740-2757, [http://dx.doi.org/10.1175/1520-0442\(2002\)015<2740:SOTAIS>2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(2002)015<2740:SOTAIS>2.0.CO;2), 2002.

**Table 1: List of parameters used in standard model run.**

Parameter	Value	Description
$f$	$-1.4244 \times 10^{-4} \text{ s}^{-1}$	Coriolis parameter
$D_{ini}(D_{SC}^{ini})$	78 m	<b>Constant</b> ISW plume outflow thickness ( <b>constant</b> outflow supercooled layer thickness)
$W_{ini}$	3 km	ISW plume outflow width <b>with constant</b> $D_{ini}(D_{SC}^{ini})$
$C_t^{ini}$	$1 \times 10^{-6}$	Depth-averaged volumetric frazil concentration in outflow
$N_{ice}$	5	Number of frazil ice sizes
$r_{i,1}, r_{i,2}, r_{i,3}, r_{i,4}, r_{i,5}$	0.2, 0.6, 0.9, 1.2, 1.5 mm	Frazil ice radii for each class
$a_r$	0.02	Aspect ratio of frazil discs
$\bar{n}$	$1 \times 10^3 \text{ m}^{-3}$	Average number of frazil crystals in all size classes per unit volume
$C_d$	0.02	Platelet layer basal drag coefficient
$V_a$	$-0.01 \text{ m s}^{-1}$	<b>Ambient</b> flow speed
$A_H$	$100 \text{ m}^2 \text{ s}^{-1}$	Horizontal eddy viscosity
$K_H$	$20 \text{ m}^2 \text{ s}^{-1}$	Horizontal turbulent diffusivity
$S_{ini}$	34.59 psu	ISW plume outflow salinity
$T_{ini}$	$-0.0573 \times S_{ini} + 0.0832 - 7.61 \times 10^{-4} D_{ini}$	ISW plume outflow temperature
$\theta_i$	0.075	Shields criterion number



**Table 2: List of calculated skill metrics for the results of VM and VU standard model runs.**

Variable	RMSE		CC		SS	
	VM	VU	VM	VU	VM	VU
$T_{SC}^0$	0.0070 °C	0.0069 °C	0.83	0.84	0.56	0.58
Platelet layer thickness	1.034 m	2.928 m	0.91	0.01	0.79	-0.65

**Table 3: Parameter settings for sensitivity runs, indicated by check, colour-coded by ISW outflow thickness (bottom row). All other parameters remain as they were for the standard model run.**

		Drag coefficient				
		0.015	0.0175	0.02	0.0225	0.025
Frazil size configuration (mm)	A: (0.2,0.6,0.9,1.2,1.5)	✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓
	1.125×A		✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓	
	1.25×A		✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓	
	1.375×A		✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓	
	1.5×A	✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓
	1.625×A		✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓	
	1.75×A		✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓	
	1.875×A		✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓	
	2×A	✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓		✓✓✓✓✓✓✓✓
	1 km			✓✓✓✓✓✓✓✓		
$W_{int}$ -ISW plume outflow width with constant $D_{ini}(D_{SC}^{ini})$	5 km			✓✓✓✓✓✓✓✓		
$C_i^{ini}$ -Depth-averaged volumetric frazil concentration in outflow	$0.2 \times 10^{-6}$			✓✓✓✓✓✓✓✓		
	$5 \times 10^{-6}$			✓✓✓✓✓✓✓✓		
$V_a$ -Ambient flow speed	0			✓✓✓✓✓✓✓✓		
	$-0.02 \text{ m s}^{-1}$			✓✓✓✓✓✓✓✓		
$\bar{n}$ -Average number of frazil crystals in all size classes per unit volume	$200 \text{ m}^{-3}$			✓✓✓✓✓✓✓✓		
	$5000 \text{ m}^{-3}$			✓✓✓✓✓✓✓✓		
$D_{ini}(D_{SC}^{ini})(m)$ -Constant ISW plume outflow thickness (constant outflow supercooled layer thickness)	30	✓	✓	✓	✓	✓
	50					
	65					
	70					
	78					
	95					
	100					
	110					




Figure 1: Satellite image of McMurdo Sound region on 29 Nov. 2011. Purple and green frames outline the model and ice borehole (Fig. 6) domains, respectively. Colours within the purple frame indicate the steady state supercooled ISW plume thickness calculated by the vertically-modified ISW plume model in the standard run (Fig. 5d). Light gray lines outline McMurdo Ice Shelf front and coastlines. Model boundaries d-a, a-b (except the ISW outflow) and “b-c” are treated as solid walls, while “c-d” is an open boundary. Blue and red dots respectively mark the oceanographic CTD and ice drilling sites, and the blue arrow represents the location of the ISW outflow in the model. The red arrow in the inset (bottom-left) points to the location of the McMurdo Sound region. Location names C, I, W, NN, and FN mean Central, Intermediate, West, Near North, and Far North, respectively. Satellite image: NASA Rapid Response MODIS Subsets (<http://earthdata.nasa.gov/data/near-real-time-data/rapidresponse/modis-subsets>).




Figure 2: (a) Exponential profiles of equilibrium frazil concentration for selected values of  $Z^*$ . Coloured bars at the right and horizontal dashed lines indicate the distribution of supercooling (blue,  $T_{SC} > 0$ ) and overheating (red,  $T_{SC} < 0$ ) for the values of  $\sigma_{SC}$  used in (b). (b) Dependence of integral value of  $I_{gr}$  on  $Z^*$  for suspended frazil ice freezing ( $I_{gr} > 0$ ) and melting ( $I_{gr} < 0$ ) under the supercooling conditions shown in (a). The star denotes the particular conditions under which the integral values of  $I_{gr}$  calculated using VU and VM formulations are equal. Note that different y-axis scales are used for freezing and melting.





Figure 3: Schematic diagram of vertical distribution of thermal forcing and relevant processes within a supercooled ISW plume of homogeneous temperature and salinity. Secondary nucleation is the process by which the frazil ice in the smallest class is supplemented by collisions between other larger frazil ice crystals.



**Figure 4: (a) Time series of  $T_{sc}^0$  simulated by VM (solid lines) and VU (dashed lines) models at five oceanographic sites (colour-coded) in the McMurdo Sound region. (b) Time series of area-averaged  $T_{sc}^0$  (blue), platelet layer thickness (green), and frazil concentration (red) simulated by VM (solid lines) and VU (dashed lines) models over the model domain (purple frame in Fig. 1).**




Figure 5: Spatial patterns, interpolated from model results using Natural Neighbour method, of (a), (b) total, (c), (d) supercooled ISW plume thickness, and (e), (f) depth-averaged frazil concentration at the end of the standard runs of (a), (c), (e) VU and (b), (d), (f) VM models over the domain (purple frame in Fig. 1). Note that the colour scale used in (a-d) is unified.






Figure 6: (a) Platelet layer thickness over green box in Fig. 1 interpolated, using Natural Neighbor method, from ice-core measurements (red dots). (b) and (c) platelet layer thickness derived from (b) VM and (c) VU models, compared with ice-core measurements (colour-coded dots). Note that the colour scale is unified.




Figure 7: Relationship between  $T_{sc}^0$  and thickening rate classified by (a) outflow supercooled layer thickness  $D_{sc}^{ini}$  and (b)  $\overline{Z}_*$  (colour-coded). Numbers in legend of (a) represent the values of  $D_{sc}^{ini}$ . Solid and hollow dots in (b) correspond to the VM and VU model runs, respectively. Coloured lines depict the central trend of the corresponding data points shown in (a). Triangle corresponds to the standard run. The results are from the last 30 days of the model runs.






Figure 8: Comparison of  $T_{sc}^0$  calculated by the VM and VU models. Triangle corresponds to the standard run. The colour scale of  $\overline{Z}_*$  is the same as in Figure 7b.





Figure 9: Same as Fig. 7, but for the relationship between  $T_{sc}^0$  and  $C_l$ .



**Figure 10: Relationship between  $\overline{Z}_*$  and difference of (a)  $C_l$  and (b) thickening rate calculated by VM and VU models (VM minus VU), classified by outflow supercooled layer thickness  $D_{SC}^{int}$ . Numbers in legend represent the values of  $D_{SC}^{int}$ .**