



# Supraglacial pond evolution in the Everest region, central Himalaya, 2015-2018.

Caroline J. Taylor<sup>1</sup>, J. Rachel Carr<sup>1</sup>

<sup>1</sup>School of Geography, Politics and Sociology, Newcastle University, Newcastle Upon Tyne, NE1 7RU, UK.

Correspondence to: Caroline J. Taylor (C.Taylor11@newcastle.ac.uk)

**Abstract.** Supraglacial ponds are characteristic of debris-covered glaciers and greatly enhance local melt rates. They can grow rapidly and coalesce to form proglacial lakes, which represent a major hazard. Here, we use Sentinel-2A satellite imagery (10 m) to quantify the spatiotemporal changes of 6,425 supraglacial ponds for 10 glaciers in the Everest region, Nepal, between 2015 and 2018. During the study period, ponded area increased on all glaciers, but showed substantial temporal and spatial variation, and the rate of pond growth increased substantially relative to 2000-2015 (Watson et al., 2016). Both Imja and Spillway Lake expanded and Khumbu Glacier developed a chain of connected ponds. 54% of ponds were associated with an ice-cliff, but the proportion of ponds with cliffs decreased during the study period. Pond location generally corresponded to lower surface velocity, but this relationship was not ubiquitous. Ponds are now predominantly found at mid-elevations on our study glaciers, suggesting that conditions conducive to pond formation have advanced up-glacier compared to general theory. Results demonstrate the need to utilize high-resolution imagery (< 10 m), as using Landsat (30 m) would miss 55–86 % of the total ponds. Glaciers were classified by stage of development (Komori, 2008; Robertson, 2012). Two glaciers transitioned between stages between 2015 and 2018, suggesting that lakes in the region are evolving rapidly. Some study glaciers displayed characteristics of multiple classes, so we propose an adapted classification system. Overall, our results demonstrate rapid pond expansion in the Everest region and highlight the need for continued monitoring for hazard assessment.

**Key Words:** *Supraglacial ponds, remote sensing, outburst floods, hazards.*

## 1 Introduction

Mass loss from glaciers in the Himalayas has increased rapidly over the past 30 years, in response to climate change (e.g. Bolch et al., 2012; Kääb et al., 2012; Quincey et al., 2009; Gardelle et al., 2013). Here, ‘summer-accumulation type’ glaciers rely on summer-monsoon snowfall for mass gain (Bolch et al., 2012), which is thought to be reducing as temperatures rise (Fujita, 2008). The shrinkage of these freshwater reservoirs will have significant regional- and local-scale impacts (Immerzeel et al., 2010; Bolch et al., 2012; Dehecq et al., 2018). Seasonal monsoonal rainfall is the dominant source of water in the Himalaya, but glacial meltwater provides up to 40% of water supplies during the dry season (Immerzeel et al., 2010). Thus, the consistent negative mass balance of Himalayan glaciers may lead to long-term reductions in perennial flow supplied to



30 major rivers outside of monsoon season (Xu et al., 2009), and could threaten the water and food security of an estimated 70  
million people in the densely populated downstream catchments (Immerzeel et al., 2010). Additionally, increased supraglacial  
meltwater storage will likely increase the frequency of glacier related hazards in the region, particularly from Glacier Lake  
Outburst Floods (GLOFs) (Thompson et al., 2010). GLOFs are highly destructive and Nepal (along with Bhutan), has been  
identified as the most economically vulnerable to these hazards (Carrivick and Tweed, 2016), making it vital to assess how the  
35 GLOF risk will evolve with climate warming.

The Everest region comprises three catchments spanning the Nepal/Tibet border in Eastern Nepal (Fig. 1; King et al., 2017).  
These catchments have experienced atmospheric warming since the mid-1970's (Shrestha et al., 1999; Shrestha and Aryal,  
2011) and weakened summer monsoons, which has reduced glacier accumulation (Salerno et al., 2015). Consequently, glacial  
mass balance in the region has been strongly negative since the 1970s (Bolch et al., 2008; Benn et al., 2012; Kääb et al., 2012).  
40 For example mass balance in the Everest Region of Nepal was  $-0.22 \pm 0.12$  m w.e.a<sup>-1</sup> between 1999 and 2011 (Gardelle et al.,  
2013). Glaciers in the Everest region are characterised by ice-surface rock debris, which is sourced from the surrounding  
hillslopes and covers approximately ~80% of the glaciated area in the region (Fushimi et al., 1980; Sakai et al., 2000; Watson  
et al., 2016). The presence of this surface debris substantially alters the glacier mass balance gradient, in comparison to clean-  
ice glaciers: melt is suppressed close to the termini, due to the presence of thick debris which insulates the underlying ice, and  
45 enhanced up glacier, where debris is thinner and increases melt (e.g. Quincey et al., 2009; Nicholson and Benn, 2013). As a  
result, the glaciers are characterised by high elevation accumulation areas, and lower-elevation ice tongues, which have low  
surface slopes and are near-stagnant, due to the low driving stresses (e.g. Bolch and Kamp, 2006; Quincey et al., 2009; Bolch  
et al., 2011; Dehecq et al., 2015). Consequently, glaciers in the Everest region lose mass by widespread surface lowering (i.e.  
down-wasting), rather than through terminus retreat (Quincey et al., 2007; King et al., 2017).

50 The low surface slopes and slow ice velocities that characterise debris-covered glacier tongues in the Everest region (Quincey  
et al., 2007; King et al., 2017), and the Himalaya more broadly (e.g. Kääb, 2005) facilitate supraglacial pond formation, by  
allowing glacial meltwater and rainfall to accumulate in depressions on the glacier surface (Reynolds, 2000; Sakai et al., 2000;  
Miles et al., 2016 and 2017). These ice surface lakes can then coalesce to form a larger proglacial lake, which carries the risk  
of producing GLOFs (e.g. Richardson and Reynolds, 2000; Quincey et al., 2007; Thompson et al., 2012; Mertes et al., 2017).  
55 Supraglacial ponds are highly variable in character (e.g. in shape, size, turbidity and ice-cliffs), dynamic in nature, and are  
expected to become increasingly prevalent in a warming climate (Thompson et al., 2016; Miles et al., 2017; Watson et al.,  
2016; 2017a and 2017b). Understanding the spatial and temporal patterns of pond growth is therefore vital for accurately  
forecasting proglacial lake growth and the associated hazard of GLOFs (Richardson and Reynolds, 2000; Quincey et al., 2007;  
Benn et al., 2012).

60 As well as posing a hazard through the formation of proglacial lakes, supraglacial ponds represent hotspots for ablation on  
low-gradient debris-covered tongues (Miles et al., 2017; Watson et al., 2017a and 2017b). This is because they have a  
comparatively low albedo and therefore absorb more insolation, which they transmit to surrounding ice, resulting in higher  
melt rates (e.g. Reynolds, 2000; Benn et al., 2001; Röhl, 2008; Miles et al., 2016; Watson et al., 2016; Mertes et al., 2017;



Salerno et al., 2017). These enhanced melt rates cause the ponds to expand both horizontally, through subaerial and sub-  
aqueous melting at the margins, and vertically, via basal melting (e.g. Sakai et al., 2000; Röhl, 2008; Mertes et al., 2017).  
65 Variations in the spatial distribution of ponds across a glacier is thought to be governed by surface slope and velocity (e.g.  
Reynolds, 2000; Bolch et al., 2008; Quincey et al., 2009), and results in differential surface melt rates (Sakai et al., 2000; Benn  
et al., 2001; Miles et al., 2016; Watson et al., 2016; Mertes et al., 2017). Once formed, ponds can persist for months to years,  
or drain via englacial pathways (e.g. Immerzeel et al., 2014; Miles et al., 2017; Watson et al., 2017a). Patterns of pond drainage  
70 are generally governed by their interaction with crevasses and englacial features, which provide efficient drainage outlets  
(Benn and Lehmkuhl, 2000; Miles et al., 2017). Determining whether ponds drain regularly or persist is important, because  
long-duration ponding can cause substantial ice-surface melt, whereas repeated pond drainage events can convey energy into  
the glacier's interior (Miles et al., 2017), resulting in quite different ice loss patterns. Additionally, recent studies (e.g. Brun et  
al., 2017; Buri et al., 2016; Mertes et al., 2017; Watson et al., 2017b) suggest that ice-cliffs play a significant role in pond  
75 formation, by enhancing marginal pond melt and subaerial calving. Furthermore, as ponds expand, ice and debris influx into  
the pond from retreating cliff-tops increases, causing pond turbidity to increase and subsequently reducing albedo, initiating a  
positive feedback of melt (Mertes et al., 2017). These factors complicate predicting future supraglacial pond formation and  
evolution and make them highly dynamic features.

Previous studies have documented changes in supraglacial water storage across the Himalaya (Table 1; e.g. Wessels  
80 et al., 2002; Kattlemann, 2003; Bajracharya and Mool, 2007; Gardelle et al., 2011). However, the use of comparatively coarse  
resolution imagery (e.g. 30m resolution Landsat imagery and 15m resolution ASTER imagery) means substantial water  
volumes may be missed and the low repeat frequency makes differentiating between pond persistence and regular drainage  
difficult, which is important for quantifying the impact of the ponds on mass loss (Immerzeel et al., 2014; Miles et al., 2017).  
Watson et al. (2016), presented the first high temporal and spatial resolution study of supraglacial pond evolution in the Everest  
85 region, for the period 2000-2015, where a total of 9340 ponds were identified. Here, we extend this previous work to quantify  
supraglacial pond evolution between 2015 and 2018. This will provide an up-to-date picture of pond coverage within the  
region. Our main objectives are; (1) characterise the spatial and temporal evolution of supraglacial ponds, to determine the  
magnitude and extent of change since 2015, (2) assess the impacts of using higher resolution imagery versus lower resolution  
imagery on the ability to identify and assess supraglacial ponds (3) examine the impact of local glacier characteristics (glacier  
90 surface slope, ice velocities and the presence of ice cliffs) on pond formation location, area and number and (4) classify the  
stage of proglacial lake development across the region.

## 2 Methods

### 2.1 Study Site and Water-Body Definitions

The study focuses on ten debris-covered glaciers that drain the Dudh Koshi basin, within Sagarmatha National Park, Eastern  
95 Nepal (Fig. 1). The glaciers flow predominantly in a southerly direction, with the exception of Ama Dablam (north flowing)



and Imja (west flowing) glaciers (Fig. 1). All of the study glaciers have extensive debris-covered tongues and high accumulation areas. Ngozumpa, Pangbung and Khumbu glaciers have the greatest length (~15 km, ~13 km and ~11 km respectively) and Imja Glacier the shortest at ~2 km (Table 2). Of the 10 glaciers in this study, 8 were also included in Watson et al. (2016) study for the period 2000-2015 (Table 2). We add Pangbung Glacier and Sumna Glacier to our study, as they are important for downstream water resources and could potentially pose a threat to downstream communities due to GLOFs (Immerzeel et al., 2010). The distinction between glacial ‘pond’ and ‘lake’ remains poorly defined in the Himalayan literature, with Watson et al. (2016) referring to all surface water as ponds unless specifically named otherwise, and other authors switching between the terms ‘lake’ and ‘pond’ (e.g. Gardelle et al., 2011; Nie et al., 2013). Here, we use the term pond to refer to all bodies of water on the glacier surface. Proglacial lakes are discussed separately from ponds, and are defined as all water bodies that are outside of the glacier margin, but are in contact with it. Following this classification, ‘Spillway Lake’, located at the terminus of Ngozumpa Glacier, is classified as a pond as it remains bound by glacier ice on all sides, but here is discussed separately, as its very large area would skew results. Imja Lake, located at the front of the Imja/Lhotse Shar glacier complex is classed as a proglacial lake.

## 2.2 Data Sources

This study used true-colour orthorectified Sentinel 2A imagery (<10 m resolution) (available from USGS at <http://earthexplorer.usgs.gov/>) for the time period December 2015 to April 2018 (SI. Table 1). Images were chosen outside of the monsoon season (June-September), to minimize cloud cover (see SI. Table 1 for percentage cloud cover), and, where possible, images were selected from the same month to avoid inadvertently including seasonal differences into our analysis (SI. Table 1). For 2016-2018, we used imagery from April, but for 2015, we had to use data from December (SI. Table 1), as Sentinel 2A data are only available from November 2015 and we wanted to ensure that our study period continued directly on from that of Watson et al. (2016), to facilitate comparison. Furthermore, our image dates (April and December) are outside of the monsoon and summer season, which should minimise seasonal effects. True colour images were derived from combining the blue (490 nm), green (560 nm) and red (665 nm) bands. For each of the study years, two true-colour Sentinel-2A images from the same date were mosaicked to produce one spatially continuous dataset of the entire region. Glacier outlines were obtained from the Randolph Glacier Inventory 5.0 (available from GLIMS at <http://www.glims.org/maps/glims>) and modified manually to reflect the debris-covered area of each glacier. These were then used as glacier / land masks and only ponds located within this mask were included in the study.

## 2.3 Maximum Likelihood Classification and Manual Editing

A supervised classification technique was used to automatically delineate supraglacial ponds. First, we manually selected training sites that contained the primary land cover classes (e.g. clean ice, water, debris covered ice etc.) from the true-colour Sentinel 2A image. We then performed the Maximum Likelihood Classification (MLC) on the true-colour image, plus bands 5 and 7 (near infrared and thermal wavelengths respectively): both infrared and thermal wavelengths are absorbed by water



bodies so their addition aided the classification substantially. Whilst other classifications can be used (e.g. the Hierarchical Knowledge Based Classifier (HKBC) method), previous studies suggest that MLC is the most accurate classification method for delineating water stores on glaciers (Tiwari et al., 2016). This method was repeated for all image dates. The classification results were assessed manually, by comparing automatically detected pond margins to the underlying imagery. We then manually edited any ponds where the classification had failed to accurately detect the pond margins. In total, we identified 6,533 ponds, and then extracted key statistics for analysis, specifically ponded area and number of ponds.

## 2.4 Controls on Pond Location

We assessed controls on supraglacial pond formation and growth patterns, specifically: glacier surface slope, ice velocities and the presence of ice cliffs. We derived a slope map and glacier elevation profiles from the ASTER DEM (<http://earthexplorer.usgs.gov/>) to identify glaciers with particularly low slope gradients and thus potential areas for future pond development. Glacier velocities were derived from repeat-image feature tracking of Landsat images by Dehecq et al. (2015). The data used in this study are derived from average velocities for 2013-2015 and have a 120 m spatial resolution (Dehecq et al., 2015). Each of the study glaciers were divided into 10 bands, representing 10% glacier surface area, to facilitate comparison of pond locations with surface slope and ice velocities. Ice cliffs were mapped manually using the true-colour Sentinel-2A images for all time periods.

## 3 Results

### 3.1 Supraglacial Pond Change

#### 3.1.1 Regional Ponded Area Change

Across the study region, total ponded area increased on all 10 glaciers between 2015 and 2018 (Fig. 2). The most prominent changes in both pond number and area were observed on the three largest glaciers (SI. Table 2; Ngozumpa, Pangbung and Khumbu glaciers), which contain ~58% of the total 6,533 ponds identified in this study (Fig. 2). Specifically, ponded area increased by 255,849 m<sup>2</sup> (33.3 %) on Ngozumpa Glacier, 191,386 m<sup>2</sup> (36.2 %) on Pangbung Glacier and 134,299 m<sup>2</sup> (43.1 %) on Khumbu Glacier between December 2015 and April 2018 (Fig. 2). This increase mainly resulted from pond coalescence and growth on Ngozumpa and Khumbu glaciers, as the number of ponds decreased but the ponded area increased (SI. Fig. 1): the number of ponds decreased by 60 (13.5% decrease) on Ngozumpa and 3 (1%) on Khumbu (Fig. 2.). The remaining seven smaller glaciers showed an increase in ponded area between 2015 and 2018, ranging from 9,664 m<sup>2</sup> on Imja Glacier (30 % increase) to 120,162 m<sup>2</sup> on Lhotse Glacier (68 % increase). This increase in area was a result of both the establishment of new ponds and the coalescence of existing smaller area ponds (Fig. 2), for instance ponded area on Ama Dablam Glacier increased 38,958 m<sup>2</sup> and pond number increased by 29, demonstrating new pond formation. Generally, the ponded area increased on all 10 study glaciers, but most marked on the three larger glaciers.



160 Whilst total ponded area on all glaciers increased from 2015-2018 (SI. Table 2), the number of ponds found on each glacier varied (Fig. 2). Only 2 of the 10 glaciers showed an increase in pond number: on Pangbung Glacier the number of ponds increased by 4(200 ponds total), and on Ama Dablam Glacier the number of ponds increased by 29(67 ponds total) between December 2015 and April 2018. The remaining 8 glaciers exhibited an overall decrease in the number of ponds, ranging from a reduction of 2 (Lhotse Glacier) to 60 (Ngozumpa Glacier). Glaciers with decreasing pond numbers generally showed an increase in ponded area in 2015, 2017 and 2018 (SI. Fig. 1). The exception to this inverse relationship between pond number and ponded area was in 2016 (SI. Fig. 1). During this year, a large decrease in pond number was observed on 165 glaciers in the eastern part of the study area, particularly Lhotse, Lhotse Shar and Lhotse Nup glaciers (Fig. 2b). At the same time, there was a large increase in pond number on glaciers in the western part of the study area (Fig. 2c). For example Pangbung Glacier increased in pond number from 196 (2015) to 468 (2016), and Sumna Glacier increased from 78 (2015) to 182 (2016). Overall, our results show that ponded area increased on all glaciers during the study period 2015-2018, whereas the number of ponds generally declined, despite the considerable spatial and temporal variability (Fig. 2).

170 Two water bodies in the study area were assessed separately, due to their large size; supraglacial lake ‘Spillway Lake’ on the terminus of Ngozumpa Glacier, and proglacial lake ‘Imja Lake’ fronting the Imja/Lhotse Shar Glacier complex (Fig. 4). Spillway Lake, located at the terminus of Ngozumpa Glacier, underwent a net gain of 40,565 m<sup>2</sup> during the study period (December 2015 - April 2018) and reached a size of 286,367 m<sup>2</sup>. This remains the largest surface water store in the study region. The only proglacial lake identified in this study was Imja Lake, which expanded up-glacier by 455.6 m to reach and 175 area of size of 1,493,142.68 m<sup>2</sup> by 2018. Whilst these two water bodies are currently the only very large water bodies in the study area, our data show substantial growth and coalescence of surface ponds on Pangbung and Khumbu glaciers (Fig. 3b, c).

### 3.1.2 Glacier-scale Pond Changes

180 Despite the overarching trends across the region, changes in supraglacial pond number and area varied from glacier to glacier and across individual glaciers (Fig. 2; Table 2). For the largest three glaciers, most of the ponded area was located near the terminus, and this persisted throughout the study period (Fig. 3a-c). The number of ponds, although demonstrated an overall decrease, showed limited year to year variation, whilst ponded area increased (Fig. 2), which was primarily as a result of pond coalescence on the lower glacier tongues. (Fig. 3a-c). For example, on the terminus of Ngozumpa Glacier, two smaller ponds (Fig. 3a, i and ii) formed new branches of Spillway Lake in April 2017. This reduced the total number of ponds by 2, whilst 185 increasing the total ponded area of Spillway Lake. Similarly, along the eastern margins of Khumbu Glacier, lateral pond expansion between April 2016 and April 2017 (Fig. 3b, i and ii) resulted in coalescence of two major ponds and a reduction in the number of surface ponds by 2. Our data therefore suggests that supraglacial pond expansion on the larger glaciers in our study area predominantly results from pre-existing ponds coalescing, rather than by the formation and growth of new ponds.

190 On the seven smaller glaciers, there was substantial spatial and temporal variability in the number and area of surface ponds, both between the glaciers and across individual glaciers (Fig. 3d and Fig. 4). For example, Lhotse Shar and Imja Glacier



neighbour each other, in the east of the study region (Fig. 1). However in April 2018, Lhotse Shar Glacier had over three times the ponded area (135,420 m<sup>2</sup>) of Imja Glacier (42,603 m<sup>2</sup>) as well as almost three and a half times the number of ponds (96 and 28 respectively). Sumna Glacier in the west of the region showed major variations in ponded area and number over time, as its percentage pond cover ranged from 2.16 % to 4.87 % during the four-year study period. For most of the smaller glaciers, ponds occupied similar locations at each time step (e.g. Ama Dablam Glacier SI. Fig. 2a and Lhotse Glacier SI. Fig. 2b). Sumna Glacier was the exception to this and showed a distinctive change in the spatial pattern of the pond locations during the study period: at the start of the study (2015), it had more ponds at higher elevations, but by the end (2018), ponds were most concentrated near the terminus (Fig. 3d). Overall, our results show an increase in both ponded area and number of ponds on the smaller glaciers, but this showed substantial spatial and temporal variation, even between neighbouring glaciers and on the same glacier.

### 3.2 Controls on Pond Location

#### 3.2.1 Glacier Elevation Profile

The study glaciers have an average slope >18° at the tongue and <25° at higher elevations. In general, areas where the mean overall slope is lower (>10°) contain more ponds. Overall, slope angle tends to decrease closer to the glacier termini, where slopes are generally between 2° and 4° (Fig. 5). However, with the exception of Sumna Glacier (see section 3.1.2.) there is no apparent relationship between elevation and pond number/ area on the study glaciers (Figs. 5 and 6). We assessed this in further detail by dividing each glacier into 10 equal elevation bands (to account for differences in total length and to facilitate direct comparison between glaciers) and calculating the number and area of ponds in each elevation band (Figs. 5 and 6): Bands are numbered from 1 (the glacier head wall) to 10 (terminus). On six of the ten study glaciers, the number of ponds decreased from the head of the glacier to terminus, whilst the pattern was reversed on the remaining 4, so that pond numbers increased with distance up glacier (Fig. 5 and 6). Furthermore, all the study glaciers showed large variations in ponded area and number between individual elevation bands.

For all study glaciers, the largest number of ponds were usually found in the central elevation bands (bands 5-6), as exemplified by Pangbung (32.2 % of ponds), Khumbu (23.1 %) and Lhotse Shar (47.8 %) glaciers (Fig. 5). The exception to this trend can be seen on Imja, Lhotse and Sumna glaciers, where the highest numbers of ponds can be found nearer to the termini (i.e. elevation bands 9 and 10), and Ama Dablam Glacier where most ponds are located nearer the high accumulation zone (bands 1-2; Fig. 5). Furthermore, the number of ponds was much higher on Imja Glacier's terminus (band 10; 6 ponds; 21.4 %) than on any of the other study glaciers. No ponds were identified in the high accumulation zone (band 1) for four of the ten study glaciers; Imja, Nuptse, Lhotse Shar and Sumna (Fig. 5 and 6). Sumna Glacier also displayed distinctive areas of higher frequency ponding (bands 6-8) and lower frequency ponding (bands 2-4) which showed no clear relationship with elevation (Fig. 3d; Fig. 5). A number of breaks in slope were identified on six of the ten study glaciers: Ama Dablam, Imja,



Lhotse, Lhotse Shar, Lhotse Nup and Sumna (Fig. 5). In the band immediately down glacier of the break in slope, there was an increase in pond number on four of the glaciers (Imja, Lhotse, Lhotse Shar, and Sumna; Fig. 5). This was most notable on Lhotse Glacier, where there was a change in slope in band 1 and the number of ponds increased from 1 (band 1) to 20 (Band 2) (Fig. 5). Ama Dablam and Lhotse Nup were the exception to this trend, where on Ama Dablam Glacier there was a decrease in pond number from 8 to 4, following the break in slope in band 3 (Fig. 5) whilst on Lhotse Nup a break in slope in band 6 preceded a decrease in pond number from 7 to 1 (Fig. 5).

### 3.2.2. Glacier Velocity

Generally, glacier velocity decreased from source to terminus, being highest in bands 1-3 and lowest in the bands 8-10 (Fig. 7). Where velocities were higher, the number of surface ponds was generally lower (Fig. 7). For instance, in bands 1 and 2 on Lhotse Shar Glacier there were no ponds recorded (velocity  $>20 \text{ m a}^{-1}$ ). However, further down glacier from band 3, velocities were lower ( $<20 \text{ m a}^{-1}$ ) and the number of ponds increased by 14 (14.6 %) (Fig. 7). The main exception to this trend is Nuptse Glacier, where velocity is low in bands 1-2 ( $> 2 \text{ m a}^{-1}$ ) and there are no ponds, but the area of higher velocity in bands 4 to 5 ( $>14 \text{ m a}^{-1}$ ) contains a total of 34 ponds (Fig. 7). In general, where velocities are lowest, for instance nearer the glacier terminus, pond number increases, (Fig. 7). For example, from band 5 onwards on Lhotse Shar Glacier, there is almost no recordable velocity and pond number reaches its highest (23 ponds) (Fig. 7). The relationship between total pond area and glacier velocity is similar to that for pond number (Fig. 8): higher velocities coincide with lower ponded area, whereas lower velocities have a higher ponded area (Fig. 8). Sumna Glacier displays this relationship clearly: it has 23 % of its ponds in Bands 2-4 and 70 % in bands 6-8. This corresponds to velocities of 3 to  $7 \text{ m a}^{-1}$  and  $< 3 \text{ m a}^{-1}$  respectively. Overall, our results indicate that lower velocities correspond with higher ponded area and pond number, and higher velocities generally relate to fewer ponds.

### 3.2.3. Ice Cliffs

Although ice cliffs were identified on all 10 glaciers 2015-2018, the number of cliffs showed marked spatial and temporal variation during this period (Fig. 9). Ngozumpa and Khumbu glaciers had the highest percentage area of the glacier covered by ice cliffs, with 4.3% and 3.92% respectively, and Sumna glacier the least (1.1%; Fig. 9). The greatest temporal variability was observed on Ama Dablam Glacier, where percentage ice-cliff coverage decreased from 2.59 % in 2015 to 1.3 % in 2016, and then rapidly increased to 3.3 % in 2018 (Fig. 9). The number of supraglacial ponds with a corresponding ice-cliff exceeded the number without: on average, across all of the study glaciers, 54 % of ponds had a coincident ice-cliff (Fig. 9). During the study, the number of ponds without an ice cliff increased on average by 1.6 % of the total glacier surface area and this was most noticeable on the seven smaller glaciers (Fig. 9). For instance, on Sumna Glacier, the area of the ponds with a cliff remained relatively stable ( $\sim 0.8 \%$ ), whereas the area of ponds without a cliff increased from 1 % in 2015 to 1.75 % in 2018 (Fig. 9). In comparison, the number of ponds with an ice cliff increased by just 0.9 % (Fig. 9). For example, on Ngozumpa Glacier, ponds with cliffs increased by 0.84 % over the four year period compared to 1.54 % over the four year period for



ponds without cliffs. This may indicate pond growth can occur irrespective of ice cliff presence, thus whilst ice cliffs continue  
255 to form in the Everest region, ponded area is increasing at a greater rate.

### 3.3 Future Lake Development

Each of the 10 study glaciers was assigned a number, according to the stage of lake development described in established lake  
classification schemes (Komori, 2008; Robertson, 2012; Table 3). In 2015 (Fig. 10b), the study region was dominated by  
glaciers in Stage 2 of lake development, with 60% showing ponds that have coalesced, were ice-dammed and have large  
260 ponded areas (>20,000m<sup>2</sup>). Only Ngozumpa Glacier was defined as Stage 3, due to the presence of the large terminal Spillway  
Lake (Fig. 10). The remaining three glaciers (Sumna, Lhotse Nup and Ama Dablam Glaciers) were all classified as Stage 1,  
with supraglacial ponds forming in their lower ablation zones (Fig. 10b).

Over the four-year study period, (December 2015- April 2018) two glaciers (Ama Dablam and Lhotse Nup) transitioned to a  
new stage of lake development (Fig. 10). Both progressed from Stage 1, where a few supraglacial ponds were identified, to  
265 Stage 2, where ponds had begun to coalesce (Fig. 10). Two glaciers (Pangbung and Lhotse Shar) partially transitioned from  
Stage 1 to Stage 2, and from Stage 2 to Stage 3, respectively (Fig. 10). Features that fit more than one stage were identified on  
these two glaciers, meaning that they could not be assigned a stage using the current classification. For example, on Pangbung  
Glacier, ponds were appearing on the lower ablation zone (characteristic of Stage 1), but some ponds were also beginning to  
coalesce (Stage 2; Fig. 3c). On Lhotse Shar Glacier, coalescing was observed (Stage 2), but there was also stable expansion of  
270 its proglacial lake (Stage 3). As a result, the current proglacial lake development in the study area cannot be captured by  
existing classification schemes.

## 4 Discussion

### 4.1. Changes in Supraglacial Pond Area 2015-2018

The area of supraglacial ponds in the study increased markedly between 2015 and 2018, ranging from a 13.6 % increase on  
275 Sumna Glacier to a 108.1 % increase on Lhotse Nup Glacier, despite showing large inter-annual variations (Fig. 2). These  
increases show a marked acceleration in pond growth, compared to 2000-2015 (Watson et al., 2016). For instance, ponded  
area increased at a rate of 1.6 % a<sup>-1</sup> on Nuptse Glacier between 2000 and 2015 (Watson et al., 2016), whereas our study  
measured a rate of 22.6 % a<sup>-1</sup> between 2015 and 2018 (SI. Table 2). Similarly, the rate of expansion on Lhotse Glacier in our  
study (17.0 % a<sup>-1</sup>) is four times greater than that found by Watson et al. (2016; 4.1 % a<sup>-1</sup>). This is a major concern in terms of  
280 risks to downstream communities, as these very high rates of pond growth will rapidly increase the water volumes available  
for outburst floods and will also encourage pond coalescence and the rate of proglacial lake formation.

One potential explanation for the observed acceleration in pond growth relates to climatic controls: warmer air  
temperatures should increase melt rates and hence encourage pond expansion, whilst increased precipitation could add water  
directly to the ponds. Data on climate trends proximal to our study glaciers are very limited. However, available data (Salerno



285 et al., 2015) suggest that minimum and mean air temperatures have risen in the Everest area between 1994 and 2013, at  
elevations above 5000 m. However, warming was most marked in spring and winter, and would thus have a more limited  
impact on ice melt, and it was concurrent with a reduction precipitation (Salerno et al., 2015), which would decrease direct  
inputs to the ponds. As such, we suggest that the observed increase in ponding may at least partly reflect changes in the  
dynamics of our study glaciers, which may provide the conditions that promote pond formation. Between 2000 and 2017,  
290 glaciers in East Nepal decelerated by  $-1.8 \pm 0.1 \text{ ma}^{-1}$  ( $17.0 \pm 1\% \text{ a}^{-1}$ ) and thinned, which in turn reduced driving stresses  
(Dehecq et al., 2018). Down-wasting and deceleration creates an inverted mass balance gradient and an uneven glacier surface,  
which together facilitate pond formation (e.g. Reynolds, 2000; Miles et al., 2016 and 2017). Furthermore, slow flow is likely  
to reduce the number of crevasses forming and would thus reduce the chance of pond drainage (Immerzeel et al., 2014; Miles  
et al., 2017; Watson et al., 2017a and 2017b). As such, we suggest that recent changes in ice dynamics in the Everest region  
295 are likely to be contributing to observed pond growth and that this may lead to a positive feedback, whereby rapid pond growth  
accelerates down wasting, leading to further pond expansion.

#### 4.2. Glacier-scale Poned Area Patterns

The three larger glaciers (Ngozumpa, Khumbu and Pangbung) increased substantially in ponded area between 2015 and 2018  
(Fig. 2: 33.3 % on Ngozumpa Glacier, 36.2 % on Pangbung Glacier and 43.1 % on Khumbu Glacier). Changes in pond number  
300 show less net change, with only Ngozumpa Glacier demonstrating a clear decline in pond number (decrease of 60) (Fig. 2).  
This suggests that pond area is generally increasing via the expansion of existing ponds and through pond coalescence, as seen  
at the terminus of Ngozumpa Glacier (Fig. 3a) and the eastern margins of Khumbu Glacier (Fig. 3b), with new pond formation  
playing a smaller role, accounting for the year to year variability. This has implications for both glacier lake related hazards  
and ice loss rates in the future. As ponds continue to join, and the area of supraglacial ponds increases, the likelihood of  
305 proglacial lakes formation is increased (Komori, 2008; Robertson, 2012), which in turn increases the risk of GLOFs (e.g.  
Richardson, 2000; Quincey et al., 2007; Benn et al., 2012; Rounce et al., 2017). In addition, the larger ponded surface area  
increases the wind fetch, which may lead to enhanced undercutting at pond margins and thus increase glacier melt rate (Benn  
et al., 2001; Röhl, 2006, 2008; Sakai et al., 2009). Our data demonstrate that the three largest glaciers had low velocities across  
their tongues: on Ngouzmpa and Pangbung glaciers, velocity remained below  $12 \text{ ma}^{-1}$  for all 10 bands, whilst on Khumbu  
310 Glacier velocities started high ( $> 30 \text{ ma}^{-1}$ ) in band 1, but rapidly reduced to below  $12 \text{ ma}^{-1}$  for the remaining 9 bands (Fig. 7  
and 8). We suggest that these slow velocities promoted pond coalescence and growth (Benn et al., 2012, 2017) and also reduced  
the frequency of pond drainage, by limiting the number of open crevasses (e.g. Miles et al., 2017), and thus increasing pond  
area.

Generally, the smaller glaciers in the region do not have large, extensive ponding at their termini, but the ponded area  
315 on all seven of the smaller glaciers increased during the study period (Fig. 2). For example, the ponded area on Ama Dablam  
Glacier increased by  $38,958 \text{ m}^2$  (48 %) 2015 to 2018 (Fig. 2b). Similar to the larger glaciers, pond number on the seven smaller  
glaciers did not undergo net change. Thus, given the overall increasing trend in area of the ponds but variations in the changes



to pond number across the region, we suggest that area increases are also primarily due to the coalescing of smaller area ponds (e.g. Ama Dablam Glacier SI. Fig. 3). Whilst pond number generally shows no clear trends, Ama Dablam Glacier is the exception, and clearly shows an increase in pond number across all study years (Fig. 2c), which suggests area increase on this glacier could be due to the formation of new ponds in addition to pond coalescing, which is consistent with earlier observations (Watson et al., 2016). This coalescing of ponds may enhance melt rates, by increasing the fetch across the pond and increasing the ponded area (Sakai et al., 2009), and potentially lead to proglacial lake development. Ice velocities are generally higher on the smaller glaciers, and more spatially variable, which may limit the opportunity for pond growth and/or encourage crevasses formation and thus promote drainage (e.g. Immerzeel et al., 2014; Miles et al., 2017; Watson et al., 2017a). A regional difference was noted in 2016, where pond number increased markedly on glaciers to the west, remained relatively similar centrally and decreased on glaciers to the east. For example, pond number increased by 272 on Pangbung Glacier and by 104 on Sumna Glacier, whilst Khumbu, Lhotse and Lhotse Shar glaciers experienced decreases of 58, 91 and 54 respectively. Pond number on Ngozumpa Glacier in the centre of the region decreased by just 33. We speculate that topographic influence by the ridge that divides east/west (Fig. 1) may result in basin specific microclimates, which may account for the changes noted here. Generally however, pond number decreased across the region and ponded area increased, subject to interannual variations. For 2016-2018, we used imagery from April, but for 2015, we had to use data from December (SI. Table 1), as Sentinel 2A data are only available from November 2015 and we wanted to ensure that our study period continued directly on from that of Watson et al. (2016), to facilitate comparison. Whilst we acknowledge that using imagery from December 2015 could introduce seasonal differences into our results, our data show that the trend in pond area, and the lack of trend in pond number, are apparent through all years of our study, suggesting that seasonal variations have a limited impact on our results.

#### 4.3. Evaluating the use of Sentinel Data for Remote Sensing Studies

Previous studies of supraglacial and proglacial lakes changes in the Everest region have largely used Landsat imagery (resolution = 30 m, 1 pixel = 900 m<sup>2</sup>; Table 1; e.g. Gardelle et al., 2011; Nie et al., 2013; Zhang et al., 2015). Here we use Sentinel 2, which is 10 m resolution (1 pixel = 100 m<sup>2</sup>). Our results demonstrate that ponds < 100 m<sup>2</sup> (one pixel in Sentinel data) accounted for 3% - 8% and those < 400 m<sup>2</sup> (four pixels in Sentinel data) comprised 28% - 59% of total ponds found in the region (SI. Fig. 4) (translating to 0.08% - 0.35% and 2.23% - 8.90% for ponded area respectively). Of the total number of ponds identified, between 55% and 86% of ponds were under 900 m<sup>2</sup> (i.e. one pixel in Landsat), which equated to between 7.47% and 30.14% of the total ponded area. As such, using Landsat imagery to map pond changes in the region would have missed the majority of the total number of ponds, and a large proportion of the ponded area. Whilst these ponds are comparatively small in area, including them in assessments is vital, as they inform us about where ponds are nucleating, and hence controls on their formation. These data also indicate locations that may become ponded in the future, and therefore subject to enhanced melt rates, and/or areas that may eventually coalesce with other ponded sections. Furthermore, recent work has demonstrated that Sentinel-2 imagery has a better spectral contrast between debris-cover ice and supraglacial ponds than Landsat of RapidEye, which affirms its suitability (Watson et al., 2018).



## 4.4 Controls on Pond Formation

### 4.4.1. Glacier Elevation Profile

Previous work suggests that supraglacial ponds usually begin to form on slopes  $< 10^\circ$  and larger ponds occur where surface gradients are less than  $2^\circ$ , typically found close to the glacier terminus (e.g. Quincey et al., 2007; Bolch et al., 2012). In some areas, large number of ponds coincided with slopes of  $2\text{--}4^\circ$  and pond frequency was highest at the termini of Imja, Lhotse and Sumna glaciers (Figs. 5 and 6). However, contrary to theory, supraglacial ponds were also found in areas with much greater slope gradients ( $> 10^\circ$ ), both at the termini and further up-glacier (Figs. 5 and 6). For instance, the highest number of ponds on Ama Dablam Glacier (76 %) were located in the high accumulation zone (band 1-2). We also observed changes in the number of ponds after breaks in slope (Fig. 5), which may reflect localised areas of extensional / compressional flow that would open or close crevasses, and thus facilitate or reduce pond drainage (Benn et al., 2001, 2012; Gulley and Benn, 2007; Röhl, 2008; Thompson et al., 2012; Mertes et al., 2017).

Our data show that the greatest number and area of ponds occurred in the mid-elevation bands (bands 5-6) and not at the termini (bands 8-10) of our study glaciers (Figs. 5-6). We suggest this is because of the inverted mass balance gradient observed on debris-covered glaciers in the Everest region: thick debris at the terminus suppresses melt, whereas thinner debris further up glacier enhances melt (Bolch et al., 2008; Quincey et al., 2009; King et al., 2018). This pattern of mass balance results in near-stagnant ice velocities over much of the tongue and rapid down-wasting at mid-elevations (Quincey et al., 2007, 2009; Nicholson and Benn, 2013; Juen et al., 2014; King et al., 2017), which produces uneven ‘ablation topography’ (Nicholson and Benn, 2013). Together, this creates ideal conditions for pond formation in mid-elevation areas: the uneven topography allows water to collect; melt rates are at their maximum, due to the comparatively thin debris cover enhancing ice melt; and low ice velocities inhibit drainage via crevasse opening. In contrast, pond formation may be more limited close to the termini, due to the much thicker debris insulating the underlying ice and thus reducing both the amount of meltwater available to fill ponds and the rate at which ablation topography forms to contain the water. Our data therefore suggest that the optimal conditions for pond formation are now predominantly found at mid-elevations (bands 5 to 6) on glaciers in the Everest region, compared to the termini identified in previous studies (Reynolds, 2000; Bolch et al., 2008; Quincey et al., 2009; Sakai and Fujita, 2010). This has important implications for total ice loss rates, as ponds strongly enhance glacier melting (Sakai et al., 2000; Benn et al., 2001; Miles et al., 2016; Watson et al., 2016; Mertes et al., 2017), and could thus expand the area of enhanced ice loss further up glacier.

### 4.4.2. Velocity

Ice velocities strongly influence pond size, pond drainage and pond persistence (Miles et al., 2017). Where velocities are low, little reorganisation of internal water pathways can occur, resulting in ineffective englacial drainage from the glacier surface, which in turn promotes surface ponding (Jordan and Stark, 2001; Benn et al., 2012, 2017). As a result, low surface gradients at glacier tongues generally encourage pond formation and low surface velocities facilitate storage, as opposed to drainage



(Reynolds, 2000; Quincey et al., 2007; Watson et al., 2017a). Our results generally support this, as there were more ponds and a greater ponded area closer to the study glacier termini (bands 8-10), where velocities were lower (Figs. 7 and 8). For example, on Sumna, Imja and Lhoste Shar glaciers no ponds were found in bands 1 or 2 where velocity was  $> 20 \text{ m a}^{-1}$ , but as velocity decreased ( $< 20 \text{ m a}^{-1}$ ) pond numbers began to increase (Fig. 7). However, as with glacier slope, there are exceptions to this relationship. For example, the peak number of ponds corresponds with peak velocities on Lhotse and near-peak velocities on Nupste (Fig. 7). The cause of these velocity patterns is difficult to elucidate, but we suggest that it may relate to the pattern of debris inputs from the surrounding hillslopes (and hence the debris characteristics and distribution), the location of tributary glaciers and/or meltwater inputs from the surrounding slopes. Overall our data indicate that ice velocities influence pond area and number, but that the relationship is far from simple and requires further detailed study.

#### 4.4.3. Ice-Cliffs

All ten of the glaciers included in this study had surface ice-cliffs, and the majority of ponds were associated with adjacent ice cliffs (54 %). This follows theory, as ice cliffs melt rates are three to six times greater than that of debris-covered ice (Kirkbride, 1993; Benn et al., 2001; Röhl, 2008; Watson et al., 2016). Thus, ice cliffs can not only initiate pond formation through surface melt (Sakai et al., 2002; Buri et al., 2016a; Watson et al., 2017b) but also facilitate pond expansion in process of thermo-erosional notching (Josberger, 1978; Röhl, 2006). Despite this, our results show that the proportion of ponds without an ice cliff increased substantially during the study period (2015-2018; Fig. 9). For example on Sumna Glacier, ponded area without an adjacent cliff almost doubled during the study, whereas ponded area with a cliff actually decreased, meaning that there were fewer ponds with an ice cliff than without in 2018 (Fig. 9). Overall, our results show that ponds are expanding and that they are usually associated with ice cliffs (Figs. 2 and 9). However, as ponded area increases, the proportion of the ponds with ice cliffs reduces (Fig. 9). We speculate that this may be because pond growth is outstripping ice cliff formation, however given the rapid growth and decay often observed of ice cliffs (e.g. Sakai et al., 2002; Watson et al., 2017b) this is the opposite trend to what would be expected. It could be suggested that higher resolution ( $< 10 \text{ m}$ ) imagery is required to identify the full range of ice cliffs on the glacier surface, with smaller cliffs not picked up by Sentinel 2A, however this requires further investigation.

#### 4.5. Future Lake Development

##### 4.5.1 Applicability of Lake Classification Models

During this study, two of the ten study glaciers progressed to a new stage of the Komori/Robertson (2008; 2012) proglacial lake classification scheme: Ama Dablam and Lhotse Nup, both progressed from Stage 1 to Stage 2. However, Komori (2008) found glaciers in Bhutan took on average 40 years to pass through Stage 1 and 2, and enter into Stage 3, whilst those in the Aoraki/Mt Cook region took 8-30 years for the same transition (Robertson, 2012). Given the short time-frame of this study (4 years), our results demonstrate that proglacial lakes in the Everest region are evolving rapidly and quicker than other regions, both in the Himalaya and globally. This has important implications for hazard assessments in the region, as these rapid changes



require high temporal resolution monitoring to determine potential changes in GLOF risk and could result in hazardous lakes  
415 forming quickly in new locations. Our results therefore highlight the need for frequent monitoring of the hazards posed by  
glacier lake growth in the Everest region, particularly given that Nepal (along with Bhutan) is at the greatest economic risk  
from GLOFs (Carrivick and Tweed, 2016).

Two of our study glaciers exhibited characteristics of multiple classes of proglacial lake development (Fig. 10).  
Specifically, on Pangbung Glacier, ponds were appearing on the lower ablation zone during the study period, which is  
420 indicative of Stage 1, but some ponds began to coalesce (Stage 2) (Table 3). On Lhotse Shar Glacier, we observed pond  
coalescing (Stage 2), but also the stable expansion of the proglacial lake (Stage 3). As such, the proglacial lake development  
observed in our study region does not fit within the four stage classification system of Komori (2008) and Robertson (2012;  
Table 3). Additionally, changes in ponded area, number of ponds and ice-cliffs were observed on all glaciers (Figs. 2 and 9),  
but these substantial differences were not accounted for in the current model. For example, both Khumbu and Nuptse glaciers  
425 remained in Stage 2 throughout the study period, but experienced increases in percentage area of ice-cliffs (45 % and 76 %  
respectively; Fig. 9). Ngozumpa Glacier remained in Stage 3, but showed a marked increase in both ponded area (33 %) and  
the number of ponds with ice-cliffs (24 %; Fig. 2c and Fig. 9). As such, the current classification model does not account for  
many of the observed changes in pond area and characteristics, which could contribute to proglacial lake growth, and does not  
account for glaciers that bridge different categories. Thus, on the basis of our observations, we propose a six-stage  
430 categorisation which accounts for the ‘in-stage’ changes observed, such as appearance of ice-cliffs and marginal pond  
expansion (Fig. 11). The inclusion of two additional stages (shown in red), now include ‘in-stage’ changes observed in this  
study, such as marginal expansion, ice-cliff formation and pond drainage. The inclusion of these stages would enable us to  
assign our study glaciers to just one stage, making the model more suitable for evaluating and communicating glacier lake  
hazard potential.

#### 435 4.4.2 Lake Development Trajectories and Outburst Risk

A key observation of this study was the coalescing of smaller area ponds on the eastern margins of Khumbu Glacier and the  
Ngozumpa terminus (e.g. Fig. 3a and b, i and ii). Despite this, both glaciers remained in the same stage of proglacial lake  
classification during the period (Stage 3 and Stage 2 respectively). Using our new classification scheme, Khumbu Glacier  
would be classified as Stage 3 (compared to currently assigned Stage 2), suggesting that Khumbu Glacier is further along the  
440 progression of proglacial lake formation than previous classifications have indicated. Our data show that Spillway Lake grew  
by 40,565 m<sup>2</sup> between 2015 and 2018. Previous work showed that Spillway Lake grew by ~ 10% per year (2001 to 2010) to  
reach 258,000 m<sup>2</sup> in December 2009 (Thompson et al., 2012) but more recently shrank by 8,345 m<sup>2</sup> between 2009 and 2015  
(Watson et al., 2016). It was suggested that this reduction in area resulted from supraglacial drainage channel evolution and  
lowering of the hydrological base level, which paused the lake’s expansion (Watson et al., 2016). Given Spillway lake grew  
445 by 40,565 m<sup>2</sup> to a size of 286,367 m<sup>2</sup> (December 2015- April 2018), our results indicate the channel has since stabilised,  
allowing Spillway Lake to expand up-glacier.



During the 4 years of our study, Imja Lake expanded by 456 m up-glacier and Spillway Lake increased in area by ~ 14% a<sup>-1</sup>. At the same time, we saw large ponds form via coalescing on the margins of Khumbu and Ngozumpa glaciers, and pond coalescence begin on Ama Dablam Glacier (Fig. 3 and SI. Fig. 2a). We also observed an increase ponded area increased on all of our study glaciers (Fig. 2). As such, our results show that increasing volumes of water are being stored on, and in front of, glaciers in the Everest region, and that this trend has accelerated, when our data are compared to previous assessments (Watson et al., 2016). The glaciers in our study region appear to be rapidly moving along the trajectory of proglacial lake formation, and may be doing so quicker than other regions (Komori, 2008; Robertson, 2012). These developments have major implications for downstream GLOF risk, as lake volume and speed of expansion are key factors in the hazard potential of proglacial lakes (Rounce et al., 2017b). As such, there is an urgent need for high resolution monitoring of ice-surface water volumes and proglacial lake development in the Everest Region.

## 5 Conclusions

This study represents the most up-to-date assessment of supraglacial ponds in the Everest region of Nepal. All 10 glaciers demonstrated an overall increase in ponded area over the period 2015 to 2018, ranging from 13.6 % (Sumna Glacier) to 108 % (Lhotse Nup Glacier). Given the short time span of this study, this represents a marked acceleration in the rate of pond growth compared to that presented by Watson et al. (2016) for the period 2000 – 2015, which utilised higher resolution imagery (~ 0.5–2 m). This is a major concern for proglacial lake formation and thus future risk to downstream communities. Our data suggests that increases in ponded area are predominantly through pond expansion and coalescence, rather than new pond formation. This suggests a transition towards larger surface ponds and lakes. Our study shows the importance of using high resolution imagery, as had medium resolution imagery (e.g. Landsat at 30 m resolution) been used, we would have missed the majority of ponds identified in this study (in the order of 55 % to 86 %). Our data show the conditions for pond formation are now predominantly found at mid-elevations on glaciers in the Everest region, suggesting the area where ponds can form may be advancing up-glacier from the debris-covered tongues, allowing ponds to establish over a larger percentage of the glacier surface. Generally, ponded area and number correspond to ice velocities, but there are notable variations in this relationship, which may relate to debris input locations and/or the presence of tributary glaciers. Ice cliffs were found on all 10 study glaciers, but the proportion of ponds with ice cliffs reduced during the study period, suggesting that the rate of ice cliff formation may be slower than pond expansion in this region. Two glaciers (Ama Dablam and Lhotse Nup) progressed to a new stage of proglacial lake development between 2015 and 2018 and this transition occurred much faster than observed elsewhere (e.g. Komori, 2008; Robertson, 2012). Our results suggest that existing classification schemes do not adequately capture proglacial evolution in the study region, so we propose a new six-stage model that encompass observed changes on our study glaciers. Overall, our results demonstrate that the volume of water stored in supraglacial ponds and proglacial lakes in the Everest region is increasing rapidly, and highlights the need for high temporal resolution monitoring of these features, in order to accurately determine the near-future GLOF risk in their downstream catchments.



## 6 Conflict of Interest

480 The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## 7 Acknowledgements

We acknowledge two freely available datasets used in this study; EarthExplorer from which the Sentinel-2A data and ASTER DEMs were obtained, and the Randolph Glacier Inventory (GLIMS Glacier Database) for the glacier shapefiles. We are also  
485 grateful to Dehecq et al. (2015) for velocity data.

## References

- Bajracharya, S. R., Mool, P. K. and Shrestha, B. R.: Impact of climate change on Himalayan glaciers and glacial lakes: Case studies on GLOF and associated hazards in Nepal and Bhutan, Kathmandu: International Centre for Integrated Mountain Development, 2007.
- 490 Benn, D. I. and Lehmkuhl, F.: Mass balance and equilibrium-line altitudes of glaciers in high-mountain environments, *Quatern. Int.*, 65, 15-29, [https://doi.org/10.1016/S1040-6182\(99\)00034-8](https://doi.org/10.1016/S1040-6182(99)00034-8), 2000.
- Benn, D.I., Wiseman, S. and Hands, K.A.: Growth and drainage of supraglacial lakes on debris-mantled Ngozumpa Glacier, Khumbu Himal, Nepal, *J. Glaciol.*, 47(159), 626-638, 2001.
- Benn, D.I., Bolch, T., Hands, K., Gulley, J., Luckman, A., Nicholson, L.I., Quincey, D., Thompson, S., Toumi, R. and  
495 Wiseman, S.: Response of debris-covered glaciers in the Mount Everest region to recent warming, and implications for outburst flood hazards, *Earth-Sci Rev.*, 114, 156–174, <https://doi.org/10.1016/j.earscirev.2012.03.008>, 2012.
- Benn, D.I., Thompson, S., Gulley, J., Mertes, J., Luckman, A. and Nicholson, L.: Structure and evolution of the drainage system of a Himalayan debris-covered glacier, and its relationship with patterns of mass loss, *The Cryosphere*, <https://doi.org/10.5194/tc-11-2247-2017>, 2017.
- 500 Bolch, T. and Kamp, U.: Glacier mapping in high mountains using DEMs, Landsat and ASTER data. *Grazer Schriften der Geographie und Raumforschung*, 41, 37-48, 2006.
- Bolch, T., Buchroithner, M. F., Peters, J., Baessler, M., and Bajracharya, S.: Identification of glacier motion and potentially dangerous glacial lakes in the Mt. Everest region/Nepal using spaceborne imagery, *Nat. Hazard. Earth. Sys.*, 8(6), 1329-1340, [www.nat-hazards-earth-syst-sci.net/8/1329/2008/](http://www.nat-hazards-earth-syst-sci.net/8/1329/2008/), 2008.
- 505 Bolch, T., Piezonka, T. and Benn, D. I.: Multi-decadal mass loss of glaciers in the Everest area (Nepal Himalaya) derived from stereo imagery, *The Cryosphere*, 5(2), 349-358, doi:10.5194/tc-5-349-2011, 2011.
- Bolch, T., Kulkarni, A., Käab, A., Huggel, C., Paul, F., Cogley, J.G., Frey, H., Kargel, J.S., Fujita, K., Scheel, M., Bajracharya, S. and Stoffel, M.: The State and Fate of Himalayan Glaciers, *Science*, 336(6079), 310, [10.1126/science.1215828](https://doi.org/10.1126/science.1215828), 2012a.



- Bolch, T., Peters, J., Yegorov, A., Pradhan, B., Buchroithner, M. and Blagoveshchensky, V.: Identification of potentially dangerous glacial lakes in the northern Tien Shan, *Nat. Hazard*, 59(3), 1691-1714, <https://doi.org/10.1007/s11069-011-9860-2>, 2012b.
- Brun, F., Berthier, E., Wagnon, P., Kääh, A. and Treichler, D.: A spatially resolved estimate of High Mountain Asia glacier mass balances from 2000 to 2016, *Nat. Geosci.*, 10(9), 668-673, [10.1038/NGEO2999](https://doi.org/10.1038/NGEO2999), 2017.
- Buri, P., Miles, E.S., Steiner, J.F., Immerzeel, W.W., Wagnon, P. and Pellicciotti, F.: A physically based 3-D model of ice cliff evolution over debris-covered glaciers, *J. Geophys. Res. Earth Surf.*, 121(12), 2471-2493, doi:10.1002/2016JF004039, 2016a.
- Buri, P., Pellicciotti, F., Steiner, J. F., Miles, E. S., and Immerzeel, W. W.: A grid-based model of backwasting of supraglacial ice cliffs on debris-covered glaciers, *Ann. Glaciol.*, 57(71), 199-211, doi: 10.3189/2016AoG71A059, 2016b.
- Carrivick, J. L., and Tweed, F. S.: A global assessment of the societal impacts of glacier outburst floods, *Global Planet. Change*, 144, 1-16, <https://doi.org/10.1016/j.gloplacha.2016.07.001>, 2016.
- Dehecq, A., Gourmelen, N. and Trouvé, E.: Deriving large-scale glacier velocities from a complete satellite archive: Application to the Pamir–Karakoram–Himalaya, *Remote. Sense. Environ.* 162, 55-66, <https://doi.org/10.1016/j.rse.2015.01.031>, 2015.
- Dehecq, A., Gourmelen, N., Gardner, A.S., Brun, F., Goldberg, D., Nienow, P.W., Berthier, E., Vincent, C., Wagnon, P. and Trouvé, E.: Twenty-first century glacier slowdown driven by mass loss in High Mountain Asia, *Nat. Geosci.*, 1, <https://doi.org/10.1038/s41561-018-0271-9>, 2018.
- Fujita, K.: Influence of precipitation seasonality on glacier mass balance and its sensitivity to climate change, *Ann. Glaciol.*, 48, 88-92, <https://doi.org/10.1038/s41561-018-0271-9>, 2008.
- Fushimi, H., Yoshida, M., Watanabe, O. and Upadhyay, B. P.: Distributions and grain sizes of supraglacial debris in the Khumbu glacier, Khumbu Region, East Nepal, *J. Jap. Society. Snow and Ice*, 41(Special), 18-25, 1980.
- Gardelle, J., Arnaud, Y. and Berthier, E.: Contrasted evolution of glacial lakes along the Hindu Kush Himalaya mountain range between 1990 and 2009, *Glob. Planet. Change*, 75(1-2), <https://doi.org/10.1016/j.gloplacha.2010.10.003>, 47-55, 2011.
- Gardelle, J., Berthier, E., Arnaud, Y., and Kaab, A.: Region-wide glacier mass balances over the Pamir-Karakoram-Himalaya during 1999-2011 (vol 7, pg 1263, 2013), *The Cryosphere*, 7(6), 1885-1886, doi:10.5194/tc-7-1263-2013, 2013.
- Gulley, J. and Benn, D. I.: Structural control of englacial drainage systems in Himalayan debris-covered glaciers, *J. Glaciol.*, 53(182), 399-412, 2007.
- Immerzeel, W.W., van Beek, L.P.H. and Bierkens, M.F.P.: Climate Change Will Affect the Asian Water Towers, *Science*, 328(5984), 1382, [10.1126/science.1183188](https://doi.org/10.1126/science.1183188), 2010.
- Immerzeel, W. W., Kraaijenbrink, P. D. A., Shea, J. M., Shrestha, A. B., Pellicciotti, F., Bierkens, M. F. P., and De Jong, S. M.: High-resolution monitoring of Himalayan glacier dynamics using unmanned aerial vehicles, *Remote. Sense. Environ.*, 150, 93-103, <https://doi.org/10.1016/j.rse.2014.04.025>, 2014.
- Iwata, S., Aoki, T., Kadota, T., Seko, K. and Yamaguchi, S.: Morphological evolution of the debris cover on Khumbu Glacier, Nepal, between 1978 and 1995, *IAHS-AISH Pub.*, 3-11, 2000.
- Jordan, R.E. and Stark, J.A.: Capillary tension in rotting ice layers (No. ERDC/CRREL-TR-01-13). ENGINEER RESEARCH AND DEVELOPMENT CENTER HANOVER NH COLD REGIONS RESEARCH AND ENGINEERING LAB, 2001.
- Josberger, E. G.: A laboratory and field study of iceberg deterioration, In *Iceberg Utilization* (pp. 245-264), 1978.




- Juen, M., Mayer, C., Lambrecht, A., Han, H., and Liu, S.: Impact of varying debris cover thickness on ablation: a case study for Koxkar Glacier in the Tien Shan, *The Cryosphere*, 8(2), 377-386, doi:10.5194/tc-8-377-2014, 2014.
- Kääb, A.: Combination of SRTM3 and repeat ASTER data for deriving alpine glacier flow velocities in the Bhutan Himalaya, *Remote. Sense. Environ.*, 94(4), 463-474, <https://doi.org/10.1016/j.rse.2004.11.003>, 2012.
- Kääb, A., Berthier, E., Nuth, C., Gardelle, J. and Arnaud, Y.: Contrasting patterns of early twenty-first-century glacier mass change in the Himalayas, *Nature*, 488, 495, doi:10.1038/nature11324, 2012.
- Kattelmann, R.: Glacial lake outburst floods in the Nepal Himalaya: a manageable hazard?, *Nat. Hazards*, 28(1), 145-154, 2003.
- 555 King, O., Quincey, D. J., Carrivick, J. L. and Rowan, A. V.: Spatial variability in mass loss of glaciers in the Everest region, central Himalayas, between 2000 and 2015, *The Cryosphere*, 11(1), 407-426, doi:10.5194/tc-11-407-2017, 2017.
- Kirkbride, M.P.: The temporal significance of transitions from melting to calving termini at glaciers in the central Southern Alps of New Zealand, *Holocene*, 3(3), 232-240, 1993.
- Komori, J.: Recent expansions of glacial lakes in the Bhutan Himalayas, *Quatern. Int.*, 184(1), 177-186, 560 <https://doi.org/10.1016/j.quaint.2007.09.012>, 2008.
- Mertes, J. R., Thompson, S. S., Booth, A. D., Gulley, J. D. and Benn, D. I.: A conceptual model of supra-glacial lake formation on debris-covered glaciers based on GPR facies analysis, *Earth. Surf. Processes. Landf.*, 42(6), 903-914, DOI: 10.1002/esp.4068, 2017.
- Miles, E.S., Willis, I.C., Arnold, N.S., Steiner, J. and Pellicciotti, F.: Spatial, seasonal and interannual variability of supraglacial ponds in the Langtang Valley of Nepal, 1999–2013, *J. Glaciol.*, 63(237), 88-105, doi: 10.1017/jog.2016.120, 2016.
- Miles, E.S., Steiner, J., Willis, I., Buri, P., Immerzeel, W.W., Chesnokova, A. and Pellicciotti, F.: Pond Dynamics and Supraglacial-Englacial Connectivity on Debris-Covered Lirung Glacier, Nepal, *Front. Earth. Sci.*, 5(69), <https://doi.org/10.3389/feart.2017.00069>, 2017a.
- Nicholson, L., and Benn, D. I.: Properties of natural supraglacial debris in relation to modelling sub-debris ice ablation, *Earth. Surf. Processes. Landf.*, 38(5), 490-501, DOI: 10.1002/esp.3299, 2013.
- 570 Nie, Y., Liu, Q. and Liu, S.: Glacial lake expansion in the Central Himalayas by Landsat images, 1990–2010, *PLoS One*, 8(12), e83973, <https://doi.org/10.1371/journal.pone.0083973>, 2013.
- Quincey, D.J., Richardson, S.D., Luckman, A., Lucas, R.M., Reynolds, J.M., Hambrey, M.J. and Glasser, N.F.: Early recognition of glacial lake hazards in the Himalaya using remote sensing datasets, *Global. Planet. Change*, 56(1), 137-152, 575 <https://doi.org/10.1016/j.gloplacha.2006.07.013>, 2007.
- Quincey, D.J., Luckman, A. and Benn, D.: Quantification of Everest region glacier velocities between 1992 and 2002, using satellite radar interferometry and feature tracking, *J. Glaciol.*, 55(192), 596-606, 2009.
- Reynolds, J.M.: On the formation of supraglacial lakes on debris-covered glaciers, *IAHS publ.*, 153-164, 2000.
- Richardson, S.D. and Reynolds, J.M.: An overview of glacial hazards in the Himalayas, *Quatern. Int.*, 65-66, 31-47, 580 [https://doi.org/10.1016/S1040-6182\(99\)00035-X](https://doi.org/10.1016/S1040-6182(99)00035-X), 2000.



- Robertson, C. M.: Temporal evolution of the termini and subaqueous morphologies of lake-calving glaciers in Aoraki/Mount Cook National Park, New Zealand: a thesis presented in partial fulfilment of the requirements for the degree of Doctor of Philosophy in Geography at Massey University, Palmerston North, New Zealand, Doctoral dissertation, Massey University, 2012.
- 585 Röhl, K.: Thermo-erosional notch development at fresh-water-calving Tasman Glacier, New Zealand, *J. Glaciol.*, 52(177), 203-213, 2006.
- Röhl, K.: Characteristics and evolution of supraglacial ponds on debris-covered Tasman Glacier, New Zealand, *J. Glaciol.*, 54(188), 867-880, 2008.
- Rounce, D. R., Byers, A. C., Byers, E. A. and McKinney, D. C.: Brief communication: Observations of a glacier outburst flood from Lhotse Glacier, Everest area, Nepal, *The Cryosphere*, 11(1), 443, doi:10.5194/tc-11-443-2017, 2017a.
- 590 Rounce, D. R., Watson, C. S. and McKinney, D. C.: Identification of hazard and risk for glacial lakes in the Nepal Himalaya using satellite imagery from 2000–2015, *Remote. Sense.*, 9(7), 654, 654; doi:10.3390/rs9070654, 2017b.
- Sakai, A., Takeuchi, N., Fujita, K. and Nakawo, M.: Role of supraglacial ponds in the ablation process of a debris-covered glacier in the Nepal Himalayas, *IAHS PUBL.*, 119-132, 2000.
- 595 Sakai, A., Nakawo, M., and Fujita, K.: Distribution characteristics and energy balance of ice cliffs on debris-covered glaciers, Nepal Himalaya. *Arct. Antarct. Alp. Res.*, 34(1), 12-19, DOI: 10.1080/15230430.2002.12003463, 2002.
- Sakai, A., Nishimura, K., Kadota, T. and Takeuchi, N.: Onset of calving at supraglacial lakes on debris-covered glaciers of the Nepal Himalaya, *J. Glaciol.*, 55(193), 909-917, 2009.
- Salerno, F., Guyennon, N., Thakuri, S., Viviano, G., Romano, E., Vuillermoz, E., Cristofanelli, P., Stocchi, P., Agrillo, G., Ma, Y. and Tartari, G.: Weak precipitation, warm winters and springs impact glaciers of south slopes of Mt. Everest (central Himalaya) in the last 2 decades (1994–2013), *The Cryosphere*, 9(3), 1229-1247, doi:10.5194/tc-9-1229-2015, 2015.
- 600 Salerno, F., Thakuri, S., Tartari, G., Nuimura, T., Sunako, S., Sakai, A. and Fujita, K.: Debris-covered glacier anomaly? Morphological factors controlling changes in the mass balance, surface area, terminus position, and snow line altitude of Himalayan glaciers, *Earth Planet. Sci. Lett.*, 471, 19-31, <https://doi.org/10.1016/j.epsl.2017.04.039>, 2017.
- 605 Shrestha, A. B., Wake, C. P., Mayewski, P. A. and Dibb, J. E.: Maximum temperature trends in the Himalaya and its vicinity: an analysis based on temperature records from Nepal for the period 1971–94, *J. Climate*, 12(9), 2775-2786, 1999.
- Shrestha, A.B. and Aryal, R.: Climate change in Nepal and its impact on Himalayan glaciers, *Reg. Environ. Change.*, 11(1), 65-77, <https://doi.org/10.1007/s10113-010-0174-9>, 2011.
- Thompson, S., Benn, D., Luckman, A. and Kulesa, B.: The evolution of supraglacial lakes, Ngozumpa Glacier, Khumbu Himal, Nepal, In *EGU General Assembly Conference Abstracts* (Vol. 12, p. 11396), 2010.
- 610 Thompson, S. S., Benn, D. I., Dennis, K., and Luckman, A.: A rapidly growing moraine-dammed glacial lake on Ngozumpa Glacier, Nepal, *Geomorphology*, 145, 1-11, <https://doi.org/10.1016/j.geomorph.2011.08.015>, 2012.
- Thompson, S., Benn, D.I., Mertes, J. and Luckman, A.: Stagnation and mass loss on a Himalayan debris-covered glacier: processes, patterns and rates, *J. Glaciol.*, 62(233), 467-485, doi: 10.1017/jog.2016.37, 2016.
- 615 Tiwari, R. K., Garg, P. K., Saini, V. and Shukla, A.: Comparisons of different methods for debris covered glacier classification, In *Khanbilvardi, R., Ganju, A., Rajawat, A.S. and Chen, J.M, Land Surface and Cryosphere Remote Sensing III, International Society for Optics and Photonics*, 9877-64, 2016.



- Watson, C.S., Quincey, D.J., Carrivick, J.L. and Smith, M.W.: The dynamics of supraglacial ponds in the Everest region, central Himalaya, *Global Planet. Change.*, 142(Supplement C), 14-27, <https://doi.org/10.1016/j.gloplacha.2016.04.008>, 2016.
- 620 Watson, C. S., Quincey, D. J., Carrivick, J. L., and Smith, M. W.: Ice cliff dynamics in the Everest region of the Central Himalaya, *Geomorphology*, 278, 238-251, <https://doi.org/10.1016/j.geomorph.2016.11.017>, 2017.
- Watson, C.S., King, O., Miles, E.S. and Quincey, D.J.: Optimising NDWI supraglacial pond classification on Himalayan debris-covered glaciers, *Remote Sense. Environ.*, 217, 414-425, <https://doi.org/10.1016/j.rse.2018.08.020>, 2018.
- 625 Wessels, R. L., Kargel, J. S. and Kieffer, H. H.: ASTER measurement of supraglacial lakes in the Mount Everest region of the Himalaya, *Ann. Glaciol.*, 34(1), 399-408, 2002.
- Xu, J., Grumbine, R.E., Shrestha, A., Eriksson, M., Yang, X., Wang, Y.U.N. and Wilkes, A.: The Melting Himalayas: Cascading Effects of Climate Change on Water, Biodiversity, and Livelihoods, *Conserv. Biol.*, 23(3), 520-530, <https://doi.org/10.1111/j.1523-1739.2009.01237.x>, 2009.
- 630 Zhang, G., Yao, T., Xie, H., Wang, W. and Yang, W.: An inventory of glacial lakes in the Third Pole region and their changes in response to global warming, *Global Planet. Change.*, 131, 148-157, <https://doi.org/10.1016/j.gloplacha.2015.05.013>, 2015.



635 **Figure 1: Location of the 10 study glaciers (blue outline) within the Everest region. Arrows dictate direction of ice flow. Glacier names are as follows: S- Sumna Glacier, P- Pangbung Glacier, N- Ngozumpa Glacier, K- Khumbu Glacier, Nu- Nuptse Glacier, LN- Lhotse Nup Glacier, L- Lhotse Glacier, LS- Lhotse Shar Glacier, I- Imja Glacier, AD- Ama Dablam Glacier. Inset: Location of the Everest Region within Nepal. Background image is surface elevation, derived from ASTER DEM (available at <http://earthexplorer.usgs.gov/>).**



Reference	Date	Imagery	Resolution	Focus
Iwata et al. (2000)	1978-1995	SPOT	Not specified	-Sketch map from SPOT imagery compared to 1987 field survey.
Wessels et al. (2002)	2000	ASTER	15m	-Water delineated for a single time period.
Bolch et al. (2008)	1962-2005	ASTER Landsat	15m 30m	-Normalized Difference Water Index (NDWI) and manual delineation used to classify water bodies.
Gardelle et al. (2011)	1990-2009	Landsat	30m	-Decision tree used to classify lakes using the NDWI.
Salerno et al. (2012)	2008	AVNIR-2	10m	-Manual digitalisation of water bodies.
Thompson et al. (2012)	1984-2010	Aerial Photos ASTER Landsat	<1m 15m 30m	-Multi-temporal analysis of Spillway Lake expansion. Glacier area change not reported.
Nie et al. (2013)	1990-2010	Landsat	30m	-NDWI based classification, but no area changes reported.
Zhang et al. (2015)	1990-2010	Landsat	30m	-Water bodies manually digitalised but ponded area changes not reported.
Mertes et al. (2016)	2014	GeoEye	<1m	-Conceptual model of supraglacial lake evolution based on Ground Penetrating Radar (GPR) facies analysis of Spillway Lake.
Watson et al. (2016)	2000-2015	GoogleEarth WorldView 1&2, GeoEye, QuickBird-2	2m 0.5-0.6m	-Water bodies semi-automatically or manually digitised. Area changes are quantified.


Table 1: Previous remote sensing studies of supraglacial water storage in the Everest region (After Watson et al., 2016).







665

Glacier	Area (km <sup>2</sup> )	Length (km)	Min-Max Elevation (range)
			(m)
Ama Dablam	2.13	4.40	4769 – 5084 (315)
Imja	1.08	2.46	5023 – 5187 (164)
Khumbu	6.64	10.82	4956 – 5246 (290)
Lhotse	5.74	6.69	4715 – 5245 (530)
Lhotse Nup	1.38	3.82	4954 – 5310 (356)
Lhotse Shar	3.00	4.03	5008 – 5429 (421)
Ngozumpa	15.10	15.76	4868 – 5541 (673)
Nuptse	3.06	6.20	4662 – 5354 (692)
Pangbung*	11.34	13.14	4742 – 5380 (638)
Sumna*	5.34	7.39	4888 – 5502 (614)

680 **Table 2: Characteristics of the 10 study glaciers, showing area, length, elevation and direction of flow. \*Indicates glaciers not included in the Watson et al. (2016) study.**




**Figure 2: Changes in supraglacial ponds observed on all 10 glaciers March 2015- April 2018; (a) pondered area as a percentage of total glacier area, (b) total number of ponds and (c) total pondered area. Bars are colour coded by date. AD-Ama Dablam Glacier, I-Imja Glacier, K-Khumbu Glacier, N-Ngozumpa Glacier, NU-Nuptse Glacier, L-Lhotse Glacier, LN-Lhotse Nup Glacier, LS-Lhotse Shar Glacier, P-Pangbung Glacier, S-Sumna Glacier.**







690 **Figure 3: Poned area change for the Ngozumpa (a) Khumbu (b) Pangbung (c) and Sumna (d) Glaciers, for the period 2015-2018. Inset (a) up-glacier expansion of Spillway Lake, inset (b) pond expansion along the lower eastern margins of Khumbu Glacier, inset (c) pond expansion on the glacier tongue, (d) pond proliferation on the lower ablation zone 2015-2017 and drainage along the western margins 2017-2018. Arrows indicate direction of ice flow. Background image is 2018 Sentinel-2 imagery (from USGS at <http://earthexplorer.usgs.gov/>).**




695 **Figure 4: Area change of Imja Lake, in the southeast of the region for the period 2015-2018. Arrows indicate direction of ice flow. Background image is 2018 Sentinel-2 imagery (freely available from USGS at <http://earthexplorer.usgs.gov/>).**



**Figure 5:** Number of ponds found on the glacier surface April 2018 compared to glacier elevation/slope at 10% glacier area distance bands (0-10 = band 1, 10-20= band 2 etc.) Dashed red line indicates breaks of slope.




**Figure 6: Poned Area found on the glacier surface April 2018 compared to glacier elevation/slope at 10% glacier area distance bands (0-10 = band 1, 10-20= band 2 etc.)**




705

**Figure 7: Number of ponds found on the glacier surface April 2018 compared to glacier velocity at 10% glacier area distance bands (0-10 = band 1, 10-20 = band 2 etc.**



710 **Figure 8: Pondered Area found on the glacier surface April 2018 compared to glacier velocity at 10% glacier area distance bands (0-10 = band 1, 10-20 = band 2 etc.)**



**Figure 9: Percentage of ponds with and without ice-cliffs, displayed as percentage of total glacier surface area. AD- Ama Dablam Glacier, I- Imja Glacier, K- Khumbu Glacier, N- Ngozumpa Glacier, NU- Nuptse Glacier, L- Lhotse Glacier, LN- Lhotse Nup Glacier, LS- Lhotse Shar Glacier, P- Pangbung Glacier, S- Sumna Glacier.**



Stage of Lake Development	Description
0	No supraglacial ponds. <span style="float: right;">720</span>
1	Appearance and growth of supraglacial ponds in the lower ablation zones
2	Coalescing of supraglacial ponds to form larger, ice dammed ponds.
3	Stable expansion of ponds up-glacier and shift to moraine dammed proglacial lake. <span style="float: right;">725</span>
4	Glacier retreats out of the proglacial lake.

730 **Table 3: Stage of lake development following Komori (2008) and Robertson (2012).**





Figure 10: Stage of lake development change observed over the study period, from 2015 (bottom) to 2018 (top). AD- Ama Dablam Glacier, I- Imja Glacier, K- Khumbu Glacier, N- Ngozumpa Glacier, NU- Nuptse Glacier, L- Lhotse Glacier, LN- Lhotse Nup Glacier, LS- Lhotse Shar Glacier, P- Pangbov Glacier, S- Sumna Glacier. Background image is 2018 Sentinel-2 image (available from USGS at <http://earthexplorer.usgs.gov/>).

760



775 **Figure 11: Updated Komori (2008) and Robertson (2012) conceptual lake classification model, to include 2 additional stages accounting for ‘in-stage’ changes observed in this study; Stage 2: recognises the role of ice-cliffs and changes in surface debris leading to pond formation and decay; Stage 3: gives marginal based expansion of ponds and pond drainage events their own category (After Komori, 2008; Robertson, 2012; Mertes et al., 2016).**