



## Year-round Impact of Winter Sea Ice Thickness Observations on Seasonal Forecasts

Beena Balan-Sarjini<sup>1</sup>, Steffen Tietsche<sup>1</sup>, Michael Mayer<sup>1,2</sup>, Magdalena Balmaseda<sup>1</sup>, Hao Zuo<sup>1</sup>, Patricia de Rosnay<sup>1</sup>, Tim Stockdale<sup>1</sup>, and Frederic Vitart<sup>1</sup>

<sup>1</sup>The European Centre for Medium-Range Weather Forecasts, , Shinfield Rd, Reading RG2 9AX, UK

<sup>2</sup>Department of Meteorology and Geophysics, , University of Vienna, Vienna, Austria

**Correspondence:** Beena Balan-Sarjini (beena.sarjini@ecmwf.int)

**Abstract.** Nowadays many seasonal forecasting centres provide dynamical predictions of sea ice. While initializing sea ice by assimilating sea ice concentration (SIC) is common, constraining initial conditions of sea ice thickness (SIT) is only at its early stages. Here, we make use of the availability of Arctic-wide winter SIT observations covering 2011-2016 to constrain SIT in the ECMWF (European Centre for Medium-Range Weather Forecasts) ocean–sea-ice analysis system with the aim of improving the initial conditions of the coupled forecasts. The impact of the improved initialization on the predictive skill of Arctic sea ice for lead times of up to 7 months is investigated in a low-resolution analogue of the currently operational ECMWF seasonal forecasting system SEAS5.

By using winter SIT information merged from CS2 and SMOS (CS2SMOS: CryoSat2 Soil Moisture and Ocean Salinity), substantial changes of sea ice volume and thickness are found in the ocean–sea-ice analysis, including damping of the overly strong seasonal cycle of sea ice volume. Compared with the reference experiment, which does not use SIT information, forecasts initialized using SIT data show a reduction of the excess sea ice bias and an overall reduction of seasonal sea ice area forecast errors of up to 5% at lead months 2 to 5. Using the Integrated Ice Edge Error (IIEE) metric, we find significant improvement of up to 28% in the September sea ice edge forecast started from April. However, sea ice forecasts for September started in spring still exhibit a positive sea ice bias, which points to too slow melting in the forecast model. A slight degradation in skill is found in the early freezing season sea ice forecasts initialized in July and August, which is related to degraded initial conditions during these months. Both the ocean reanalyses, with and without SIT constraint, show strong melting in the middle of the melt season compared to the forecasts. This excessive melting related to positive net surface radiation biases in the atmospheric flux forcing of the ocean reanalyses remains and consequently degrades analysed summer SIC. The impact of thickness initialization is also visible in the sea surface and near-surface temperature forecasts. While positive forecast impact is seen in near-surface temperature forecasts of early freezing season initialized in May (when the sea ice initial conditions have been observationally constrained in the preceding winter months), negative impact is seen for the same season when initialised in August month when the sea ice initial conditions are degraded. We conclude that the strong thinning by CS2SMOS initialization mitigates or enhances seasonally dependent forecast model errors in sea ice and near-surface temperatures in all seasons.



25 The results indicate that the memory of SIT in the spring initial conditions last into autumn, influencing forecasts of the peak summer melt and early freezing seasons. Our results demonstrate the usefulness of new sea ice observational products in both data assimilation and forecasting systems, and strongly suggest that better initialization of SIT is crucial for improving seasonal sea ice forecasts.

## 1 Introduction

30 Sea ice is an integral part of the Earth system as it regulates the heat, moisture and momentum flux exchange between the polar oceans and the atmosphere. Decline in Arctic sea ice is a visible indicator of the changing climate. Forecasting Arctic sea ice has advanced significantly in the last decade, with most forecasting centres using prognostic sea ice models operationally, allowing us to explore the sea ice forecast skill on long lead times from weeks to months to seasons. Possibilities of economically viable shorter shipping routes across the Arctic in the summer are constantly being explored. Monthly and seasonal outlooks of sea  
35 ice products are therefore in great demand especially by the Arctic communities, maritime and resource extraction industries.

Moreover, there is increasing scientific evidence that warming and sea ice loss in the Arctic due to climate change affect the European weather and climate (Balmaseda et al. (2010), Mori et al. (2014), Overland et al. (2016), Ruggieri et al. (2016)). Unlike sea ice concentration and extent, long records of satellite observations of sea ice thickness are sorely lacking (Laxon et al. (2003), Kwok and Rothrock (2009), Haas et al. (2010), Meier et al. (2014), Sallila et al. (2019), Scarlat et al. (2020)).

40 Since reliable estimates of long-term, basin-wide sea ice extent and volume are needed for understanding climate change and for initializing numerical weather forecasts, there is growing interest in using improved and new types of sea ice observations in data assimilation systems (Lindsay et al. (2008), Blanchard-Wrigglesworth et al. (2011), Tietsche et al. (2013), Sigmond et al. (2013), Balmaseda et al. (2015)). Earlier studies propose that long-term memory in the winter sea ice thickness can potentially improve summer sea ice extent forecasts (Guemas et al. (2016), Tietsche et al. (2014), Day et al. (2014)). They concluded that  
45 potential predictability mainly originates from the persistence or advection of sea ice thickness anomalies, interaction with ocean and atmosphere and changes in the radiative forcing.

While assimilation of sea ice concentration (SIC) is routinely done in operational sea ice forecasting, assimilation of sea ice thickness (SIT) is at its early stages (Allard et al. (2018), Xie et al. (2018), Mu et al. (2018), Fritzner et al. (2019)). These studies have found that SIT initialization improves sea ice forecasts in forced ocean–sea-ice forecasting systems which were  
50 run for short time periods spanning from 3 months up to 3 years. Blockley and Peterson (2018) reported for the first time the positive impact of winter SIT initialization on the skill of seasonal forecasts for summer sea ice forecasts using a fully-coupled atmosphere–ocean–sea-ice model. All of these studies used either European Space Agency’s Cryosat-2 (CS2) radar altimeter freeboard SIT measurements alone (Laxon et al. (2013), Hendricks et al. (2016)) or merged with SMOS radiometric measurements (Kaleschke et al. (2012), Tian-Kunze et al. (2014)) in a dataset called CS2SMOS (Ricker et al. (2017)).

55 Currently SIC is the only sea ice variable assimilated in the ECMWF ocean–sea–ice data assimilation system. Although the ECMWF sea ice reanalysis and reforecasts compare well with other systems (Chevallier et al. (2017), Uotila et al. (2018), Zampieri et al. (2018)), they are affected by noticeable errors (Tietsche et al. (2018)). There are large biases in sea ice forecasts



60 from months to seasons, pointing to uncertainties in both the models and observations used in the assimilation and forecasting systems. Here we explore the pathway to improve the initialization using observations of sea ice thickness which covers both the thick and thin ice regions of the Arctic. We then assess the impact of the changed sea ice initial condition on the forecast skill on long lead times of months to seasons. Compared to Blockley and Peterson (2018), who looked only at summer forecast skills, our study for the first time assesses the forecast impact of SIT initialization on all seasons using a fully-coupled seasonal forecasting system. We use the ECMWF coupled ensemble seasonal forecasting system SEAS5 and CS2SMOS thickness observations.

65 The rest of the article is organised as follows. Section 2 describes the methodology of sea ice thickness initialization and forecasting, including a brief description of ocean–sea-ice models, the assimilation system, the atmosphere-ocean–sea-ice coupled forecasting system, observations used and the experimental set-up. Section 3 presents the main results: the impact of new SIT observations on the analysed sea ice state and the impact of the changed sea ice initialization on seasonal range forecasts. Finally, Section 4 provides the summary of the findings with concluding remarks.

## 70 2 Observations and Methods

The procedure followed here to assess the impact of SIT information follows a twin experiment approach. Each of the experiments consists of two distinctive steps: 1) the production of a set of ocean and sea ice initial conditions by conducting twin ocean–sea-ice assimilation experiments (ocean–sea-ice reanalyses; abbreviated as ORA), which only differ in the use of SIT information ; and 2) the production of a set of twin retrospective seasonal forecast (reforecast) experiments, initialized from the  
75 respective ORA. The ORA twin reanalyses are a low resolution variant of the currently operational ORAS5 (Zuo et al. (2019)). The seasonal forecast experiments are also low resolution versions of the operational ECMWF seasonal forecasting system SEAS5 (Stockdale et al. (2018), Johnson et al. (2018)). The impact of SIT in the ocean initial conditions and seasonal forecast is then evaluated, using verification against observational datasets and other more specific diagnostics. The verification will also use fields from ORAS5 and ERA-5 (ECMWF atmospheric Re-Analysis-5); Hersbach et al. 2019) reanalyses. Although  
80 the datasets used for verification are not strictly independent, it is relevant as cross-check variables evaluation, for instance between SIC and SIT assimilation. SIT verification is also conducted as a sanity check of the nudging approach compared to the reference experiment. In this section we first describe the sea ice information used for both initialization and verification, and then offer a brief description of the experimental set-up.

In addition to the sea ice data sets described below, the initialization step uses ocean observations: sea surface temperature, sea-level anomalies from altimeter and in-situ temperature and salinity, which are the same as those used in ORAS5, as  
85 described in Zuo et al. (2019).



## 2.1 Sea Ice Observational information

### 2.1.1 Sea Ice Concentration Product: OSI-401-b

The two ocean–sea-ice reanalysis experiments presented here assimilate the sea ice concentration product of the EUMETSAT Ocean and Sea Ice Satellite Application Facility (OSI SAF, [www.osi-saf.org](http://www.osi-saf.org); product identifier OSI-401-b (Tonboe et al. (2017))). These Level-3 OSI SAF SIC product (OSI-401-b) is produced as daily-mean fields with only a few hours latency. In contrast to the operational ORAS5 system, which uses Level-4 SIC data, experiments presented in this study use Level-3 SIC data. The main difference is that Level-4 products rely on gap-filling, whereas Level-3 products have missing data, for instance if the satellite has a temporary malfunction, or if certain areas like the North Pole are not observed. The OSI-401-b SIC observational estimate is based on SSMIS (Special Sensor Microwave Imager / Sounder) measurements. SIC is provided as the percentage of an ocean grid point covered by sea ice. The product comes in a polar stereographic grid of 10km horizontal resolution with varying pole hole size.

The impact of Level-3 SIC observations in the initialization is reported to have neutral forecast impact on seasonal sea ice forecasts and positive impact on sub-seasonal range (Balan-Sarajini et al. (2019)). The OSISAF OSI-401-b SIC data set is also used for verification of SIC and sea ice edge.

### 2.1.2 Sea Ice Thickness product: CS2SMOS

A recent initiative led by the Alfred Wegener Institute provides a merged product of Arctic-wide winter ice thickness that combines thick-ice retrievals by CryoSat2 (CS2) satellite and thin-ice retrievals by the Soil Moisture and Ocean Salinity (SMOS) satellite. This merged sea ice thickness observational product, CS2SMOS (<https://spaces.awi.de/display/CS2SMOS>, Ricker et al. (2017)), is the first ever multi-sensor ice thickness product for the Arctic. CS2 (Hendricks et al. (2016)) measures freeboard (the height of the ice or snow surface above the water level) using altimetry, whereas SMOS (Tian-Kunze et al. (2014)) measures brightness temperatures in the L-band microwave frequencies. Both measurements are converted to ice thickness in metres. Due to their different measurement principles, SMOS retrievals should be reliable for ice thinner than about 1 m and CS2 retrievals for ice thicker than 1 m. The merged product can hence represent the entire thickness range covering the whole Arctic with reasonable accuracy (Ricker et al. (2017)). CS2 and SMOS are merged using an optimal interpolation scheme to produce the CS2SMOS product, which is available on a weekly basis on an Equal-Area Scalable Earth Grid version 2 (EASE2) grid with 25km horizontal resolution. Both the CS2 and SMOS retrievals are not possible in the melt season due to signal contamination owing to the presence of melt ponds, and wet and warm snow and ice surfaces. It is only available for 5 full months from November to March of the ice growth season every year.

The CS2SMOS SIT information has been assimilated in one of the twin ORA experiments, during the November–March period. It has also been used for verification of initialization in those months. We emphasize that this dataset does not provide SIT information during the period April–October. Nevertheless, there is still substantial impact in the April–October period from constraining sea ice thickness during the November–March period, as we will see in Section 3 – a truly year-round impact.



## 2.2 Methods

### 120 2.2.1 Ocean–sea-ice Reanalysis Experiments

In order to assess the impact of new sea ice thickness observations on the assimilation, we carry out two ORAs as shown in Table 1. They are 1) a reference experiment with SIC assimilation (ORA-REF), and 2) an experiment with SIC assimilation and sea ice thickness constraint (ORA-SIT). Experiments ORA-REF and ORA-SIT are run for the time period January 2011 to December 2016, because these are the full years for which CS2SMOS observations were available at the time of experimenta-  
125 tion. Note that ORA-REF is a continuation of a longer experiment which started in 2005 and ORA-SIT starts from ORA-REF on the 1st of January, 2011.

Experiment name	SIC constraint	SIT constraint	Time period	Description
ORA-REF	Yes	No	2011-2016	SIC assimilation
ORA-SIT	Yes	Yes	2011-2016	SIC assimilation and SIT nudging

**Table 1.** Specifications of the ocean–sea-ice assimilation experiments

Our reanalysis experiments are forced by near-surface air temperature, humidity and winds as well as surface radiative fluxes from the atmospheric reanalysis ERA-Interim (ERA-I) (Dee et al. (2011)) until 2015 and from the ECMWF operational analysis from 2015 to 2016. We use the same set-up of NEMOVAR (Variational data assimilation system for NEMO (Nucleus  
130 for European Modelling of the Ocean) ocean model) used in ORAS5 (Zuo et al. (2019)) – in particular, almost the same observations are assimilated. The only differences are the following: a) a coarser model resolution as described below, b) different assimilated SIC observations compared to the current operational one and, c) a longer assimilation window of 10 days instead of 5 days.

The ocean general circulation model used in these experiments is NEMO version 3.4 (Madec (2008)) with a horizontal  
135 resolution of approximately  $1^\circ$  and 42 vertical layers. The grid is tripolar, with the poles over Northern Canada, Central Asia and Antarctica enabling higher resolution across the Arctic than at the equator. The first model layer is 10 m thick, and the upper 25 levels represent approximately the top 880 m. Both the horizontal and vertical resolution in our setup is lower than that of the current operational system, which has a horizontal resolution of approximately  $0.25^\circ$  and 75 vertical levels. The time step is one hour.

140 The prognostic thermodynamic-dynamic sea ice model used is LIM2 (Louvain-la-Neuve Sea Ice Model) (Fichefet and Maqueda (1997)). The vertical growth and decay of ice due to thermodynamic processes is modelled according to the three-layer (one layer for snow and two layers for ice) Semtner scheme (Semtner (1976)). The ice velocity is calculated from a momentum balance considering sea ice as a two-dimensional continuum in dynamical interaction with the atmosphere and



ocean. Internal stress within the ice for different states of deformation is computed following the viscous-plastic (VP) rheology  
145 proposed by Hibler III (1979). LIM2 has a single sea ice category to represent sub-grid scale ice thickness distribution, and  
open water areas like leads and polynyas are represented using ice concentration. Melt ponds are not modelled. LIM2 has a  
time step of 1 hour and is coupled to the ocean at every time step.

As for ORAS5, both experiments here use the variational data assimilation using NEMOVAR in a 3D-Var FGAT (First  
Guess at Appropriate Time) configuration as described in Mogensen et al. (2012). The length of the assimilation window is  
150 10 days in our experiments. Assimilated observations comprise temperature and salinity profiles, altimeter-derived sea level  
anomalies and sea ice concentration. Sea-surface temperature is constrained to observations by a strong relaxation. A global  
freshwater correction is added to reproduce the observed global-mean sea-level change. The assimilation of the SIC is done  
separately from the ocean variables, and is described in Tietsche et al. (2015) and Zuo et al. (2017).

In addition to the observations assimilated via NEMOVAR, the SIT in experiment ORA-SIT is constrained to the CS2SMOS  
155 via a linear nudging technique (Tietsche et al. (2013), Tang et al. (2013)). The relationship between the modelled and observed  
sea ice thickness in a grid point is described by the following equation:

$$SIT^n = SIT^m - \left[ \frac{\Delta t}{\tau} (SIT^m - SIT^o) \right] \quad (1)$$

where  $SIT^n$  is the nudged thickness,  $SIT^m$  is the modelled thickness,  $SIT^o$  is the observed thickness,  $\Delta t$  is the sea ice  
model time step of 1 hour, and  $\tau$  is the nudging coefficient corresponding to a relaxation time scale of 10 days. The choice  
160 of a 10-day relaxation time scale makes sense as a first trial, since it is consistent with the length of the assimilation window.  
To facilitate the nudging, the CS2SMOS weekly observations in EASE2 grid have been interpolated to daily gridded fields in  
ORCA 1° grid. The weekly to daily interpolation is done by appropriately weighting two adjacent weekly-mean fields. We  
have also tested the sensitivity to different nudging strengths by running variants of ORA-SIT with a relaxation time scale of  
20, 30 and 60 days, but in this study we only use the experiment with the strongest constraint (10-day relaxation time) for  
165 initializing the ensemble reforecasts.

### 2.2.2 Coupled Reforecast Experiments

In order to assess the impact of CS2SMOS sea ice thickness initialization on sea ice forecasts, we performed 2 sets of twin  
coupled ocean–sea-ice-atmosphere reforecast experiments as shown in Table 2, which only differ on the ocean–sea-ice initial  
conditions, provided by the data assimilation experiments shown in Table 1. The reference reforecast (FC-REF) is initialized  
170 by ORA-REF, and reforecast experiment FC-SIT is initialized by ORA-SIT. Comparison of results from these two sets of  
reforecasts allows quantifying the impact of SIT information on the seasonal forecasts.

The reforecast experiments are carried out using a version of the ECMWF coupled seasonal forecasting system. The coupled  
model consists of the same ocean and sea ice model (NEMO3.4/LIM2) used for our reanalysis experiments, and is coupled to  
the ECMWF atmospheric model, Integrated Forecast System (IFS) version 43r3. It is run with a horizontal resolution of 36 km,  
175 corresponding to a cubic octahedral reduced Gaussian grid at truncation TCo319 and 91 vertical levels (SEAS5 is run with IFS  
cycle 43r1 at the same atmospheric resolution, but with 0.25° horizontal resolution and 75 vertical levels in the ocean). The



Experiment name	Start years	Lead mon	Ens. size	Initial condition	Description
FC-REF	2011–2016	7	25	ORA-REF	SIC initialization
FC-SIT	2011–2016	7	25	ORA-SIT	SIC and SIT initialization

**Table 2.** Overview of the reforecast experiments. For each of the start years, forecasts are started on the 1st of every calendar month.

coupled model also includes the land surface model HTESSSEL (Hydrology Tiled ECMWF Scheme for Surface Exchanges over Land) and the ocean surface wave model WAM. The coupling of the atmosphere and ocean is done using a Gaussian interpolation method, and the coupling frequency is 1 hour. For more details on SEAS5 see (Stockdale et al. (2018), Johnson et al. (2018)).

Both reforecasts are started from the 1st of each month of each year 2011–2016, resulting in 72 forecast start dates overall. Note that out of all months of each year in the 2011–2016 period only winter (December–April) months are directly constrained by November–March observations as the CS2SMOS data is only available for those 5 full months. The initial conditions for the remaining 7 start months (May–November) of each year are indirectly affected by the thickness constraint applied earlier in the ice growth season in the reanalysis. The non-availability of the observations for the melt season in a way provides an opportunity to test the predictability of winter SIT from summer initial conditions. The forecast initialized from each start date has 25 ensemble members for both sets of reforecasts.

### 3 Results


Here we first assess the impact of sea ice thickness observations on the estimation of sea ice properties in the ORA initial conditions, and then we evaluate the impact on the skill of seasonal forecast of sea ice area, sea ice edge, sea ice volume and 2m temperature. When possible, we use the observational datasets for verification. However, as mentioned above, sea ice thickness and volume (SIV) can not be verified properly for the months April–October, due to the lack of sea ice thickness observations. In those cases, we will describe the impact in terms of differences between experiments.

#### 3.1 Impact of Sea Ice Thickness Initialization on the Sea Ice Reanalysis

Figure 1 shows the SIT bias with respect to the CS2SMOS observations for ORA-REF (Figure 1a, c) and ORA-SIT (Figure 1b, d), for March (Figure 1a, b) and November (Figure 1c, d). The ORA-REF suffers from large ice thickness bias of up to 1.4 m. The predominant bias pattern is an underestimation of ice thickness by more than 1 m in the central Arctic, and an overestimation in the Beaufort Gyre and the Canadian Archipelago of the order of 1 m. This pattern is present for all the months when CS2SMOS is available. In March, widespread overestimation in the coastal Arctic seas is also present. These biases are much reduced or absent in ORA-SIT. Most of the large-scale pattern of underestimation and overestimation of sea ice




## Bias in Sea-Ice Thickness



**Figure 1.** Bias in monthly-mean (2011-2016) sea ice thickness (m) in experiment a) ORA-REF and b) ORA-SIT, for March (a, b) and November (c, d). The reference is CS2SMOS observations. ORA-REF is the ocean–sea-ice assimilation experiment with no sea ice thickness constraint. ORA-SIT is the assimilation experiment with a thickness relaxation time scale of 10 days.



## Change in Sea-Ice Thickness, ORA-SIT - ORA-REF



**Figure 2.** Difference in monthly-mean (2011-2016) sea ice thickness (m) between experiments ORA-SIT and ORA-REF for a) March and b) July and for c) September and d) November months.



in ORA-REF is not present in ORA-SIT in March. However, slight underestimation over the central Arctic and overestimation over the Canadian Archipelago still remain in November. This is probably caused by the lack of SIT observations during the months preceding November. In contrast, the estimation of the March conditions benefit from the availability of SIT information in the preceding winter.

205 Figure 2 shows the difference in SIT between ORA-SIT and ORA-REF for March, July, September and November. The difference patterns between ORA-SIT and ORA-REF are quite consistent for all the months, characterized by a thickening of the thick ice over the Central Arctic and North of Greenland, and a thinning of the thin ice area over the Beaufort and Siberian Seas, thus enhancing the spatial gradients on sea ice thickness distribution. The largest impact occurs in March, probably because at this month the SIT observations have been assimilated during the preceding 5 months. The impact of SIT winter  
210 information lasts well into the summer months, with a slight clockwise displacement of the thinning, and a reduction of the thickening, which by September has roughly halved. The impact during March and November is consistent with a reduction of the bias in ORA-REF (Figure 1a and c). Since basin-scale SIT observations are not available for the end of melt season, biases are unknown.


The thickness constraint also affects the biases in SIC. Figure 3 shows the SIC bias w.r.t. OSI-401-b SIC as well as the  
215 SIC difference between ORA-REF and ORA-SIT. In March, ORA-REF shows mostly an overestimation of SIC all around the sea ice edge, over the Davis Strait, northeast of Greenland, Bering Sea and Okhotsk Sea. In ORA-SIT this bias is uniformly reduced up to 10%. In November (Figure 3g, h and i) ORA-REF has similar SIC overestimation biases over the sea ice edge, but this time the SIT constraint has very little impact on SIC biases. The ORA-REF biases in July are characterized by a weak underestimation of SIC. Notably, in ORA-SIT there is an increase of the negative SIC bias of more than 10% over the Pacific  
220 and Siberian Arctic sectors towards the end of melt season, with July and August (not shown) months being the most affected.

To gain some insight into the degradation of the July SIC bias in ORA-SIT we look at the mean annual cycle of the SIC assimilation increments. The assimilation increments are indicative of the corrections that the assimilation of SIC observations exerts to compensate for errors in the sea ice state. Figure 4 shows the mean annual cycle of the area-averaged assimilation increments in ORA-REF (blue) and ORA-SIT (green). In both experiments, the assimilation increments exhibit a clear seasonal  
225 cycle, with large positive increments from May to October, indicative of strong underestimation of SIC in the ORAs, and weak negative increments from December to March. The differences in SIC increments over the Arctic between the two experiments peaks during July, with ORA-SIT increments about 9% per month higher than in ORA-REF. The results in this figure indicates that 1) both ORAs melt sea ice too fast during the summer months, as shown by negative SIC biases at marginal seas of the Arctic Ocean where thin sea ice resides during the summer months (Figure 3d and e); and 2) the SIT assimilation exacerbates  
230 the summer SIC biases in ORA-SIT (as seen in eg: Figure 3e) due to corrected but thinner sea ice at the beginning of the melt season in almost all marginal seas of the Arctic Ocean (Figure 2a).


From January to May and from November to December, on an average less ice is being taken away by the increments in ORA-SIT (green) analysis than that in ORA-REF (Figure 4). These results clearly show the long-lasting effect of the SIT information: the SIT constraint was only applied during the growth season from November to March (grey shading), but its  
235 impact is evident in sea ice concentration throughout the melting season even in the presence of SIC assimilation.



### Bias and change in Sea Ice Concentration



**Figure 3.** Bias in monthly-mean (2011-2016) sea ice concentration w.r.t. OSI-401-b observations for ORA-REF (a, d, g), ORA-SIT (b, e, h), and the difference between ORA-SIT and ORA-REF for (c, f, i). Panels (a, b, c) are for March, (d, e, f) for July, and (g, h, i) for November.




**Figure 4.** Annual cycle of the mean of the SIC increments in ORA-SIT (green), and ORA-REF (blue), averaged over north of  $70^{\circ}N$  during 2011-2016. The grey shading shows months (January to March, and November to December) with CS2SMOS SIT nudging.

### 3.2 Impact of ice thickness initialization on sea ice forecasts

Figure 5a gives an overview of bias in sea ice area in the FC-REF reforecast w.r.t. ORAS5 reanalysis as a function of forecast start and lead months (units are  $10^6 \text{ km}^2$ ). ORAS5 is preferred to OSISAF for the verification of integrated sea ice area because of its complete spatial coverage. The figure shows the forecast bias for different forecast lead times (y-axis) as a function of forecast starting month (x-axis). Errors at lead month 1 are generally small throughout the year. However, for longer lead times, there is a strong over-prediction of sea ice area in summer months, and a moderate under-prediction of autumn sea ice conditions, consistent with too slow melting and refreeze respectively. The forecast biases are generally small in winter months.

These three bias regimes, in general – low bias in winter, positive bias in summer and negative bias in autumn – seem to be mostly independent of start months. These biases shown in FC-REF are quite similar to those in SEAS5 (not shown) which are discussed in more detail in Stockdale et al. (2018). The positive biases in the melt season forecasts is considerably reduced with the SIT initialisation in FC-SIT started in January to June and the negative biases in the forecasts is worsened in FC-SIT started in July to October (Figure 5b). The forecasts for winter months remain unbiased in FC-SIT. Note that the bias regimes in the forecasts are very different from the bias regimes in the reanalysis (Section 3.1), which tends to have too much ice in winter and too little ice in summer.

Impact of thickness initialization has not only improved the biases in summer SIC forecasts, but it has also improved the sea ice extent forecasts as measured by the Integrated Ice Edge Error (IIEE) (Figure 6). Seasonal forecasts of ice edge are in great




**Figure 5.** Bias in the forecast of Arctic sea ice area ( $\times 10^{12} \text{m}^2$ ) w.r.t. ORAS5 as a function of start and lead month for 2011–2016, a) in the reference reforecast FC-REF and b) in the SIT-initialised reforecast FC-SIT. Red colour denotes over-prediction of sea ice area, and blue colour denotes under-prediction.


demand for exploring economically viable Arctic shipping routes. IIEE is one of the recent user-relevant sea ice metrics on ice extent or ice edge (Goessling et al. (2016), Bunzel et al. (2017)). Since it can be decomposed into over- and under-prediction, it is more useful than the traditional basin-wide sea ice extent error. It is calculated using uncalibrated SIC forecasts.

255 IIEE for all lead months and start months verified against OSI-401-b suggests reduced error in sea ice edge (blue colours) in FC-SIT overall. The most striking feature is the significant improvement in summer forecast error for lead months 2–7 in FC-SIT compared to FC-REF. The main contribution to the error reduction of up to 30% in summer forecasts comes from the reduction of the model bias leading to consistent over-prediction (not shown). For the traditional September sea ice extent forecast starting in April, an improvement of 28% is found. Forecast verification in October and November from July and  
 260 August starts show a slight degradation, caused by under-prediction (not shown). This could again be due to the indirect effect of a thinner starting point in FC-SIT (Figure 2b) and a lower, degraded SIC in the ORA-SIT reanalysis (Figure 3e), combined with the already existing slow refreeze nature of the model.

Figures 5 and 6 point out that the impact of ice thickness initialization on the forecast bias and errors is strongly dependent on season and lead time. Most seasons and lead times are improved but some are, perhaps inevitably, deteriorated. To measure the  
 265 overall impact on forecast error and make a statement about potential skill improvements that to be expected for operational forecasts, we aggregate FC-SIT and FC-REF for all start months from January 2011 to December 2016 and compute the area-integrated mean absolute forecast error (MAE) of sea ice concentration for each lead month.



**Figure 6.** Difference in Integrated Ice Edge Error in  $10^{11} \text{ m}^2$  between FC-SIT and FC-REF reforecasts 2011–2016 w.r.t. OSI-401-b observations. Blue colour denotes reduced error in sea ice edge in FC-SIT and red colour denotes increased error in FC-SIT. Black triangles represent statistically significant (DelSole and Tippett (2016)) changes at 5% level.



**Figure 7.** Spatially integrated SIC mean absolute error over lead month for all FC-REF and FC-SIT forecasts (no. of forecast start months,  $n = 72$ ; January 2011 to December 2016) w.r.t OSI-401-b observations. Panel a) shows the error without bias correction, panel b) the error after bias correction. Lead months for which the reduction of forecast error in FC-SIT passes the DelSole and Tippett (2016) significance test at the 5% level are marked by filled circles, insignificant changes are marked by crosses.

Averaged over all start dates and grid points, Figure 7 shows that the MAE of sea ice area is substantially improved in FC-SIT. When no bias correction is applied prior to computing the MAE (Figure 7a), FC-SIT forecasts are significantly better in each lead month, with maximum error reduction of about 10%.


However, skill assessments of seasonal forecasts are conventionally made after a forecast calibration where the impact of the forecast bias is removed. By this measure, a reduction of forecast bias does not by itself count as an improvement. As Figure 7b shows, removing the respective bias of FC-SIT and FC-REF prior to computing the MAE results in a smaller error reduction: errors in FC-SIT are significantly lower only in lead months 2–5, by up to 5%. Figure 7 demonstrates that, although the thickness initialization predominantly reduces the bias, it also leads to an improvement in the skill of sea ice area forecasts that is relevant for operational forecasting systems.

Finally, we analyse the impact of SIT initialization on forecasts of pan-Arctic sea ice volume. Figure 8 shows the sea ice volume forecast climate at different lead month for the forecasts starting in May (top) and August (bottom). The forecast climate is computed by averaging the reforecast starting at a given calendar month for the years 2011–2015. Seven months forecasts started in August lead to February of the following year. Since the ORAs are not available in January and February, 2017, the year 2016 is not accounted for in this figure. For reference, the sea ice volume estimates of ORA-REF and ORA-SIT reanalyses are also shown. It is remarkable that the shape of the seasonal cycle is largely preserved between FC-REF and FC-SIT, maintaining the initial offset during the whole forecast range. The figure clearly shows that FC-SIT starts from a thinner ice state than FC-REF in both initial months.


The May starts show large differences between the forecasts and the ORAs: Both FC-SIT and FC-REF show a slower SIV decrease (lower melt rate) than the ORAs from June to September, and also a slower refreeze during October and November.



### Time evolution of mean Sea Ice Volume forecasts



a) Ensemble-mean Sea-Ice Volume Forecasts, May start months



b) Ensemble-mean Sea-Ice Volume Forecasts, August start months

**Figure 8.** Time series of ensemble-mean sea ice volume (units are  $10^{12} \text{ m}^3$ ) forecasts averaged over 2011–2015, for May start date (a) and August start date (b) in reference reforecast (FC-REF, dashed blue line) and reforecast with thickness initialization (FC-SIT, dashed green line) compared to their own reanalyses, ORA-REF (solid blue line), and ORA-SIT (solid green line).



The melt rate of the ORAs here are consistent with those in ORAS5 (see Uotila et al. (2018) or Mayer et al. (2019)). Compared to the May starts, differences between FC-SIT and FC-REF is smaller in the August start, and so is their agreement with the ORAs. Again, the FC-SIT shows smaller values than FC-REF from the beginning, and both forecast sets exhibit a parallel SIV  
290 evolution. The shape of the seasonal cycle in the forecasts is different from the ORAs; the forecasts initialized in August show a slower refreeze during October than the ORAs. However, after October, the SIV increases faster in the forecasts than in ORA-SIT, and it continues increasing more or less at the same rate until the end of January in the forecasts, while in ORA-SIT (solid green line) the freezing rate slows down after November. As a result by the end of January the forecast SIV is higher than in ORA-SIT. ORA-REF without the thickness constraint has the highest SIV in the ice growth season. In the next section  
295 we examine the discrepancies in SIV changes between ORAs and FCs in more detail.

### 3.3 Linking sea ice analysis and forecast errors to the Arctic surface energy budget

In order to investigate the physical causes of sea ice errors in the ORAs and forecasts, the Arctic surface energy budget is considered. We estimate melt energy tendency (MET), which is the energy used to melt sea ice and energy released in the process of freezing, and is proportional to SIV changes. It is defined as in Mayer et al. (2019):


$$300 \quad MET = L_f \rho \left( \frac{dSIT}{dt} \right) \quad (2)$$

where  $L_f$  denotes latent heat of fusion ( $-0.3337 \times 10^6 \text{ J kg}^{-1}$ ),  $\rho$  represents sea ice density (assumed constant at  $928 \text{ kg m}^{-3}$ ), and  $SIT$ , the grid-point averaged sea ice thickness. Thickness changes are computed as exact monthly differences. Figure 9 shows the MET mean annual cycle (2011-2015) north of  $70^\circ N$  for ORA-REF, ORA-SIT, FC-REF, and FC-SIT. The values for the forecasts are compiled from one-month forecasts from every calendar month. Assimilation increments of SIC  
305 proportionally affect SIV in the ORAs (Tietsche et al. (2013), Tietsche et al. (2015)). The resulting MET increments are shown for both ORA-REF and ORA-SIT as well. We note that the MET annual cycle of ORA-REF is very similar to that of ORAS5 (not shown).

Figure 9 clearly shows that ORA-REF exhibits the most pronounced annual cycle of MET, with strongest melting in summer and strongest freezing in winter. Earlier studies have shown that the MET annual cycle is exaggerated in ORAS5 (Uotila et al.  
310 2019; Mayer et al. 2019) and hence also in ORA-REF. ORA-SIT has a damped MET annual cycle, as the thickness constraint during winter prevents overly strong SIV accumulation. Lower SIV at the end of winter consequently leads to weaker melting in summer. However, summer melt in ORA-SIT is likely still too strong, as this experiment features a negative SIC bias in summer despite realistic SIT and SIC earlier in the year, when CS2SMOS data is available (see Figure 3e).

Both FC-REF and FC-SIT agree very well with each other and exhibit a much weaker MET annual cycle than the ORAs  
315 (Figure 9). The difference between the forecasts and the ORAs in May and June melting cannot be explained by the MET increments (neutral impact at that time), which points to differences in energy fluxes into the sea ice as a cause.

We therefore compare the mean annual cycle of surface net radiation ( $Rad_S$ ) over ocean north of  $70^\circ N$ . Figure 10a shows the 2011-2015 annual cycle of  $Rad_S$  from FC-REF, FC-SIT, ERA-I, ERA5, and the satellite-based product Clouds and Earth's



**Figure 9.** Mean annual cycle of MET over ocean area north of  $70^{\circ}N$  in ORA-REF, ORA-SIT, FC-REF (lead month 1), FC-SIT (lead month 1). MET increments for ORA-REF and ORA-SIT are shown as well.

Radiant System – Energy-Balanced and Filled Surface edition 4.0 (CERES-EBAF; Kato et al. (2018)), which we use as  
 320 reference.


We consider  $Rad_S$  from ERA-I as a good proxy for  $Rad_S$  seen by the ORAs, due to two reasons: 1) ORAs use ERA-I forcing during most of the study period, and 2) ORAs does not output  $Rad_S$  term; although it is not exactly identical e.g. due to different albedo in the ORAs. ERA-I exhibits a positive  $Rad_S$  bias in summer, peaking in June at  $15 \text{ W/m}^2$ , while FC-REF and FC-SIT agree quite well with CERES-EBAF, especially in May and June, when MET discrepancies with the ORAs are large (Figure 9). Thus the  $Rad_S$  bias of ERA-I can explain a large fraction of the overly strong MET in the ORAs during May and June, and the discrepancy between the ORAs and the forecasts.  
 325

The mean deviation of  $Rad_S$  from CERES-EBAF (Figure 10b) clearly indicates that forecasts are closer to the observational product than the atmospheric reanalyses in May and June. This positive  $Rad_S$  bias of ERA-I should be considered alongside




Mean annual cycle of surface net radiation

mean annual cycle



mean deviation from CERES-EBAF



**Figure 10.** a) Mean annual cycle of surface net radiation,  $Rad_S$  ( $Wm^2$ ) over ocean area north of  $70^\circ N$  from ERA-I, ERA5, FC-REF (lead month 1), FC-SIT (lead month 1), and CERES-EBAF, and b) Mean deviation of  $Rad_S$  from CERES-EBAF for FC-REF, FC-SIT, ERA-I and ERA5.



the results by Hogan et al. (2017), who found a negative bias in downwelling shortwave radiation in ERA-I due to excessive  
330 low-level clouds. Our results can be explained by the positive bias in downwelling longwave radiation in ERA-I outweighing  
the shortwave flux bias. Figure 10 also shows results for ERA5, which is closer to CERES-EBAF than ERA-I, which indicates  
a reduced cloud bias in this more recent atmospheric reanalysis and gives rise to the expectation of improved MET in future  
ocean reanalyses forced by this product.

Additional information on the realism of summer MET in the forecasts can be obtained from the sea ice area forecast bias  
335 of FC-SIT, as displayed in Figure 5b. It shows that FC-SIT May starts exhibit a strongly reduced positive bias compared to  
FC-REF. The bias reduction can be attributed to the improved initial conditions in FC-SIT, but the fact that the sea ice area  
bias remains positive from July onward indicates that MET in the forecasts is too low in summer. Figure 10b suggests that  
 $Rad_S$  is too low in the forecasts in July and August, which probably contributes to the positive SIA bias remaining in FC-SIT  
(Figure 5b).

340 The October MET (Figure 9) indicates stronger refreeze in the ORAs (lower MET values) compared to the forecasts. This  
is consistent with negative MET increments present in the ORAs, which act to speed up refreeze in the reanalyses (see Fig-  
ure 9). The less negative MET values of the forecasts in October are consistent with the too weak freezing and consequent  
underestimation of sea ice in autumn in the August starts.


Area-averaged net radiation of all considered products agrees well with CERES-EBAF in October (see Figure 10), and  
345 also difference maps show only a weakly positive  $Rad_S$  bias of the reanalyses and forecasts compared to CERES-EBAF (not  
shown). Hence, errors in other physical terms such as ocean-ice fluxes must play an important role in fall, but more detailed  
investigations are beyond the scope of this paper.

### 3.4 Impact of ice thickness initialization on forecasts of atmospheric variables

To discuss the impact of the sea ice thickness constraint on the atmosphere, we first assess the differences in the forecast means  
350 (or biases) between FC-SIT and FC-REF. Figure 11a shows the bias in 2m temperature ( $t_{2m}$ ) (averaged over  $50-90^\circ N$ ) in FC-  
REF as a function of start dates and lead months. Significant cold biases are present in forecasts for most of the start months  
and lead months except for non-significant warm biases in November forecasts started in August, September and October  
months. Mean differences in  $t_{2m}$  (Figure 11b) are generally positive with very few and non-significant exceptions, which is  
expected from the generally reduced sea ice cover in FC-SIT. Strongest warming with area averages of  $0.5K$  can be found  
355 during fall for forecasts started between March and September. February and March start dates show a moderate but significant  
warming at short lead times, but otherwise changes are relatively small for October to February start dates. Also, changes in  
summer temperatures are small compared to those in fall. Inspection of temperature difference patterns between FC-SIT and  
FC-REF indicates that differences in summer are confined to areas around the sea ice edge (not shown), while changes in fall  
are more widespread (see Figure 11c). The warming pattern in fall appears as a diagonal feature in Figure 11b, which suggests  
360 that changes depend more on season than on forecast lead time. Therefore, to gain more insight into the spatial structure of the  
changes, Figure 11c and d show forecast differences in 2m temperature and mean sea level pressure in SON, respectively. To



**Difference in mean t2m and mean sea level pressure forecasts**



**Figure 11.** Mean forecast differences between FC-SIT and FC-REF 2011-2016: a) bias in mean 2m temperature north of  $50^{\circ}N$  w.r.t. ERA5, as a function of start dates and lead months, in FC-REF, b) similar to a), but difference in mean 2m temperature between FC-SIT and FC-REF. Triangles denote significant changes according to DelSole and Tippett (2016) test at the 5% level. Mean forecast difference (FC-SIT - FC-REF) for SON aggregated from May, June, July, August start dates of c) 2m temperature and d) mean sea level pressure. Dots indicate areas of significant changes on the 95% level according to Komolgorov-Smirnov test.



find robust changes, the differences are aggregated from forecasts started between May and August, yielding samples of 600 forecasts. Moreover, aggregation along the diagonal maximizes the signal (compare to Figure 11b).

Widespread temperature differences  $>1\text{K}$  can be seen over the Arctic Ocean and the Canadian Archipelago in SON (Figure 11c), but significant warming spreads also south to North America and Eurasia. Solar radiation in the Arctic is very weak for SON. Hence, the warming in FC-SIT must stem from enhanced fluxes of heat from the ocean to the atmosphere, with a possible positive feedback from increased atmospheric water vapour. The fluxes are enhanced in FC-SIT due to larger areas of open waters and increased SSTs, both a result of reduced sea ice concentration. Furthermore, we find warming over the Northwest Atlantic, which is related to the warmer SSTs present already in the initial conditions from ORA-SIT (not shown). Another area of significant warming in FC-SIT relative to FC-REF can be seen over Eastern Europe and Western Russia. This warming seems consistent with patterns of mean sea level pressure differences shown in Figure 11d. They show lower pressure in FC-SIT over Scandinavia and higher pressure over central Russia, which together suggest more southerly winds in the region of warmer temperatures. Furthermore, mean sea level pressure changes indicate lower pressure over the Arctic Ocean and the Canadian Archipelago, i.e. in areas of maximum warming. In addition, there are positive pressure differences southeast of Greenland. Altogether, the patterns in sea level pressure difference resemble a wave-like response, but it should be kept in mind that only some parts of these changes are statistically significant. Nevertheless, we note that qualitatively similar and significant changes are also found in 500hPa geopotential forecasts for SON (not shown), suggesting that the features seen in Figure 11d are indeed robust.


We now turn to the question whether changes in the forecast mean constitute a forecast improvement or a forecast deterioration in the sense that they lead to an overall reduction of model biases. Since forecast bias is strongly dependent on region, season and lead time, aggregating over many seasons and lead months hampers physical understanding of the impact of thickness initialization. We therefore focus only on forecasts for the September–November (SON) season, where the impact on 2m temperature is strongest.

As Figure 12a and b show, the 2m-temperature forecast bias for the SON season have a strong dependence on the start and lead month. Cold biases are clearly dominating the entire hemisphere in May forecasts, whereas a mixture of warm and cold biases is visible in August forecasts, with predominantly warm biases over the ice edge. As discussed previously, the thickness initialization leads to a homogeneous warming of 2m temperature (Figure 11c), which is not very sensitive to the time of initialization.

To determine whether the mean change leads to an increase or a reduction in the bias, we compute changes in mean absolute error (MAE) of 2m-temperature forecasts without the usual calibration. This is shown in Figure 12c and d. Mean absolute forecast errors are substantially reduced in SON (by more than  $1\text{K}$ ) over the entire ice cover and some adjacent regions (Figure 12c). In this case, the thickness initialization helps to mitigate the model bias. Conversely, when initializing forecasts in August mean absolute forecast errors are increased over the marginal Seas of the Arctic Ocean and the Canadian Archipelago (Figure 12d). This points to an exacerbation of the model biases by the thickness initialization. However, the negative impact for August start dates is not as significant as the positive impact for May start dates.



### Bias and difference in MAE in t2m forecasts



**Figure 12.** Verification of 2m temperature against ERA5 for SON forecasts started in May (a,c) and August (b,d): Bias (in K) of FC-REF is shown on the top (a,b), and MAE difference (in K) between FC-SIT and FC-REF at the bottom (c,d). Differences significant at the 5% level according to the test by DelSole and Tippett (2016) are stippled. The homogeneous warming of FC-SIT w.r.t. FC-REF shown in Figure 11c results in MAE for SON t2m being reduced for May start dates c) and increased for August start dates d).



Forecast skill changes on other atmospheric fields have been explored as well. The picture for circulation-related fields such as mean sea-level pressure and 500 hPa, geopotential height (not shown) is less clear compared to 2m-temperature, indicating that much of the statistically significant changes found at the near-surface temperature in the Arctic are due to local thermodynamic effects.

#### 400 4 Summary and Concluding Remarks

In this paper we use 6 years of Arctic-wide sea ice thickness observations of January, February, March, November and December months during 2011 to 2016 to constrain the modelled sea ice thickness, and study the impact on the ocean–sea-ice reanalysis. Coupled forecasts of the ocean–sea-ice-wave-land-atmosphere are initialized using these data assimilation experiments, and the forecast skill of Arctic sea ice for lead times up to 7 months is investigated. To our knowledge this study  
405 provides the first comprehensive assessment of coupled seasonal sea ice forecasts with thickness initialization for all the seasons. It complements to the study by Blockley and Peterson (2018), who reported the positive forecast impact on summer season only. This paper does not delve into the technical implementation of sea ice observational information in the ECMWF systems as reported in Balan-Sarajini et al. (2019), but instead it focuses on 1) collating the relevant scientific results on the impact of sea ice thickness information alone on seasonal forecasts, 2) conducting targeted diagnostics to gain understanding  
410 of the results, and 3) providing a more thorough discussion on the impact.

Constraining initial conditions by nudging to CS2SMOS ice thickness results in a substantial reduction of sea ice volume and thickness in the ocean–sea-ice analysis. This reduces some of the existing forecast biases in SEAS5 and improves forecast skill in the melt season, but in turn increases the errors during autumn, when the existing sea ice forecast bias is negative.

The impact of sea ice thickness initialization on seasonal forecast skill for Arctic sea ice variables, namely sea ice cover, sea  
415 ice thickness, sea ice volume and sea ice edge, is mostly positive for seasonal forecasts started from January to June start dates. We find significant improvement of up to 28% in the traditional September sea ice edge forecasts started from April start dates as shown by Integrated Ice Edge Error. However, sea ice forecasts for September started in spring still exhibit a positive sea ice bias, which points to too slow melting in the forecast model. Neutral forecast impact for November and December start dates is found. However, a slight degradation is seen in autumn forecasts started from July and August start dates, which is shown  
420 to be due to errors in the sea ice initial conditions. Both the ocean reanalyses, with and without SIT constraint, show strong melting in the middle of the melt season compared to the forecasts. This excessive melting is shown to be due to positive net surface radiation biases in the atmospheric flux forcings of the ocean reanalyses. Compared to the forecasts, strong freezing is seen throughout the freeze season in the ocean reanalysis without SIT constraint. With SIT constraint applied from November to March, the existing strong freezing is somewhat damped in the late freeze season. The exact causes of the differences in  
425 freezing between the reanalyses and forecasts require further investigation. Aggregating all the forecasts started in January to December, positive forecast impact of up to 5% skill improvement for integrated SIC is found at 2-5 lead months. Thinning of sea ice by CS2SMOS mitigates or enhances seasonally dependent forecast model error.



The impact of sea ice thickness initialization on atmospheric variables has also been investigated. Changes in ensemble mean 2m-temperature over the Arctic region are significant for SON forecasts initialized from May to August start dates. The impact is also seen in mean sea level pressure and to certain extent in 500hPa geopotential height and is mostly local and thermodynamically driven, except for some remote impact over the north west Atlantic ocean. Similar to the sea ice edge forecasts, positive forecast impact is seen for 2m-temperature forecasts for the early freezing season, SON, started in May and negative impact for the same season is seen when started in August when the initial conditions are degraded. Statistically significant changes in 2m-temperature mean absolute error are predominantly due to corresponding local changes in errors in the sea surface temperature and sea ice variables. There is no clear change in forecast skill of upper atmospheric circulation in our experiments. Our results illustrate that information on sea ice thickness is relevant for identifying model errors and for exploiting the long-term memory present in ice thickness for seasonal forecasts of sea ice and near-surface temperatures. Constraining SIT in the initialisation alters biases arising due to both errors in the forcing and the sea-ice model. Though the SIT assimilation is not expected to solve these underlying problems per se, by moving the model state closer to reality, it helps us to better understand the errors in our system, as well as improving forecast skill scores in the meantime. As atmospheric forecast errors are dominated by biases, we are yet to demonstrate the benefit of interannual varying data on bias-corrected forecast scores. Robustness of impact on upper atmospheric variables and possible teleconnections need to be further assessed which would require a longer study period and larger sample size.

These findings demonstrate that making use of recently-available, spatially and temporally rich sea ice thickness observations from satellites for the ice growth season has the potential to significantly improve 1) the sea ice state in currently operational ocean–sea-ice reanalyses and, 2) the seasonal forecasts in operational forecasting systems. Our study also emphasizes the potential of future sea ice satellite missions for Earth System reanalysis and forecasts.

*Data availability.* The EUMETSAT OSI SAF sea ice concentration product (OSI-401-b) is from [www.osi-saf.org](http://www.osi-saf.org) and the CS2SMOS sea ice thickness product is from <https://spaces.awi.de/display/CS2SMOS>. The ERA5 atmospheric reanalysis data is available at the Copernicus Climate Change Service Climate Data Store (<http://cds.climate.copernicus.eu>) and the ORAS5 ocean reanalysis data is available at the Integrated Climate Data Center portal (<http://icdc.cen.uni-hamburg.de/thredds/catalog/ftpthredds/EASYInit/oras5/catalog.html>).

*Author contributions.* BBS led the manuscript writing with input and feedback from all the co-authors. MAB and ST conceived the work plan. ST, HZ and BBS created the experimental set-up. BBS, MM and ST carried out the computations and analyzed the results. MAB, PDR, TS and FV contributed to discussions and interpretation of the results.

*Competing interests.* The authors declare that they have no conflict of interest.



*Acknowledgements.* We acknowledge the European Union Horizon 2020 SPICES (Space-borne observations for detecting and forecasting sea ice cover extremes) project (640161) for funding this work. MM's work was partially supported by the Austrian Science Fund project 33177. The production of the merged CryoSat2-SMOS sea ice thickness data was funded by the ESA project SMOS and CryoSat-2 Sea Ice Data Product Processing and Dissemination Service, and data from 01/11/2011 to 31/12/2016 were obtained from Alfred Wegener Institute  
460 Helmholtz Centre for Polar and Marine Research (AWI).



## References

- Allard, R. A., Farrell, S. L., Hebert, D. A., Johnston, W. F., Li, L., Kurtz, N. T., Phelps, M. W., Posey, P. G., Tilling, R., Ridout, A., et al.: Utilizing CryoSat-2 sea ice thickness to initialize a coupled ice-ocean modeling system, *Advances in Space Research*, 62, 1265–1280, 2018.
- 465 Balan-Sarojini, B., Tietsche, S., Mayer, M., Balmaseda, M., and Zuo, H.: Towards improved sea ice initialization and forecasting with the IFS, <https://doi.org/10.21957/mt6m6rpwt>, <https://www.ecmwf.int/node/18918>, 2019.
- Balmaseda, M., Hernandez, F., Storto, A., Palmer, M., Alves, O., Shi, L., Smith, G., Toyoda, T., Valdivieso, M., Barnier, B., Behringer, D., Boyer, T., Chang, Y.-S., Chepurin, G., Ferry, N., Forget, G., Fujii, Y., Good, S., Guinehut, S., Haines, K., Ishikawa, Y., Keeley, S., Köhl, A., Lee, T., Martin, M., Masina, S., Masuda, S., Meyssignac, B., Mogensen, K., Parent, L., Peterson, K., Tang, Y., Yin, Y.,  
470 Vernieres, G., Wang, X., Waters, J., Wedd, R., Wang, O., Xue, Y., Chevallier, M., Lemieux, J.-F., Dupont, F., Kuragano, T., Kamachi, M., Awaji, T., Caltabiano, A., Wilmer-Becker, K., and Gaillard, F.: The Ocean Reanalyses Intercomparison Project (ORA-IP), *Journal of Operational Oceanography*, 8, s80–s97, <https://doi.org/10.1080/1755876X.2015.1022329>, <http://www.tandfonline.com/doi/full/10.1080/1755876X.2015.1022329>, 2015.
- Balmaseda, M. A., Ferranti, L., Molteni, F., and Palmer, T. N.: Impact of 2007 and 2008 Arctic ice anomalies on the atmospheric circulation: Implications for long-range predictions, *Quarterly Journal of the Royal Meteorological Society*, 136, 1655–1664,  
475 <https://doi.org/10.1002/qj.661>, <http://doi.wiley.com/10.1002/qj.661>, 2010.
- Blanchard-Wrigglesworth, E., Armour, K. C., Bitz, C. M., and DeWeaver, E.: Persistence and inherent predictability of Arctic sea ice in a GCM ensemble and observations, *J. Climate*, 24, 231–250, <https://doi.org/10.1175/2010JCLI3775.1>, <http://journals.ametsoc.org/doi/abs/10.1175/2010JCLI3775.1>, 2011.
- 480 Blockley, E. W. and Peterson, K. A.: Improving Met Office seasonal predictions of Arctic sea ice using assimilation of CryoSat-2 thickness, *The Cryosphere*, 12, 3419–3438, 2018.
- Bunzel, F., Notz, D., and Tietsche, S.: Definition of a new set of observation based metrics relevant for regional applications. SPICES Deliverable, D8.2 Report, 2017.
- Chevallier, M., Smith, G. C., Dupont, F., Lemieux, J.-F., Forget, G., Fujii, Y., Hernandez, F., Msadek, R., Peterson, K. A., Storto, A., et al.:  
485 Intercomparison of the Arctic sea ice cover in global ocean–sea ice reanalyses from the ORA-IP project, *Climate Dynamics*, 49, 1107–1136, 2017.
- Day, J. J., Hawkins, E., and Tietsche, S.: Will Arctic sea ice thickness initialization improve seasonal forecast skill?, *Geophysical Research Letters*, 41, 7566–7575, <https://doi.org/10.1002/2014GL061694>, <http://doi.wiley.com/10.1002/2014GL061694>, 2014.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer,  
490 P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, <http://doi.wiley.com/10.1002/qj.828>, 2011.
- 495 DelSole, T. and Tippett, M. K.: Forecast Comparison Based on Random Walks, *Monthly Weather Review*, 144, 615–626, <https://doi.org/10.1175/MWR-D-15-0218.1>, <http://journals.ametsoc.org/doi/10.1175/MWR-D-15-0218.1>, 2016.



- Fichefet, T. and Maqueda, M. A. M.: Sensitivity of a global sea ice model to the treatment of ice thermodynamics and dynamics, *Journal of Geophysical Research*, 102, 12 609–12 646, <https://doi.org/10.1029/97JC00480>, <http://doi.wiley.com/10.1029/97JC00480>, 1997.
- 500 Fritzner, S. M., Graverson, R., Christensen, K. H., Rostosky, P., and Wang, K.: Impact of assimilating sea ice concentration, sea ice thickness and snow depth in a coupled ocean-sea ice modelling system, 2019.
- Goessling, H. F., Tietsche, S., Day, J. J., Hawkins, E., and Jung, T.: Predictability of the Arctic sea ice edge, *Geophysical Research Letters*, 43, 1642–1650, <https://doi.org/10.1002/2015GL067232>, <http://doi.wiley.com/10.1002/2015GL067232>, 2016.
- Guemas, V., Blanchard-Wrigglesworth, E., Chevallier, M., Day, J. J., Déqué, M., Doblas-Reyes, F. J., Fučkar, N. S., Germe, A., Hawkins, E., Keeley, S., et al.: A review on Arctic sea-ice predictability and prediction on seasonal to decadal time-scales, *Quarterly Journal of the Royal Meteorological Society*, 142, 546–561, 2016.
- 505 Haas, C., Hendricks, S., Eicken, H., and Herber, A.: Synoptic airborne thickness surveys reveal state of Arctic sea ice cover, *Geophysical Research Letters*, 37, 2010.
- Hendricks, S., Ricker, R., and Helm, V.: User Guide-AWI CryoSat-2 Sea Ice Thickness Data Product (v1. 2), 2016.
- Hibler III, W. D.: A Dynamic Thermodynamic Sea Ice Model, *J. Phys. Oceanogr.*, 9, 815–846, 1979.
- 510 Hogan, R., Ahlgrim, M., Balsamo, G., Beljaars, A., Berrisford, P., Bozzo, A., pe, F. D. G., Forbes, R., Haiden, T., Lang, S., Mayer, M., Polichtchouk, I., Sandu, I., Vitart, F., and Wedi, N.: Radiation in numerical weather prediction, ECMWF Technical Memorandum, <https://doi.org/10.21957/2bd5dkj8x>, <https://www.ecmwf.int/node/17771>, 2017.
- Johnson, S. J., Stockdale, T. N., Ferranti, L., Balmaseda, M. A., Molteni, F., Magnusson, L., Tietsche, S., Decremmer, D., Weisheimer, A., Balsamo, G., Keeley, S., Mogensen, K., Zuo, H., and Monge-Sanz, B.: SEAS5: The new ECMWF seasonal forecast system, *Geoscientific Model Development Discussions*, 2018, 1–44, <https://doi.org/10.5194/gmd-2018-228>, <https://www.geosci-model-dev-discuss.net/gmd-2018-228/>, 2018.
- 515 Kaleschke, L., Tian-Kunze, X., Maaß, N., Mäkynen, M., and Drusch, M.: Sea ice thickness retrieval from SMOS brightness temperatures during the Arctic freeze-up period, *Geophysical Research Letters*, 39, L05 501, <https://doi.org/10.1029/2012GL050916>, <http://doi.wiley.com/10.1029/2012GL050916>, 2012.
- 520 Kato, S., Rose, F. G., Rutan, D. A., Thorsen, T. J., Loeb, N. G., Doelling, D. R., Huang, X., Smith, W. L., Su, W., and Ham, S.-H.: Surface irradiances of edition 4.0 clouds and the Earth’s radiant energy system (CERES) energy balanced and filled (EBAF) data product, *Journal of Climate*, 31, 4501–4527, 2018.
- Kwok, R. and Rothrock, D. A.: Decline in Arctic sea ice thickness from submarine and ICESat records: 1958–2008, *Geophys. Res. Lett.*, 36, L15 501, <https://doi.org/10.1029/2009GL039035>, 2009.
- 525 Laxon, S., Peacock, N., and Smith, D.: High interannual variability of sea ice thickness in the Arctic region, *Nature*, 425, 947, 2003.
- Laxon, S. W., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., Hendricks, S., Krishfield, R., Kurtz, N., Farrell, S., and Davidson, M.: CryoSat-2 estimates of Arctic sea ice thickness and volume, *Geophysical Research Letters*, 40, 732–737, <https://doi.org/10.1002/grl.50193>, <http://doi.wiley.com/10.1002/grl.50193>, 2013.
- Lindsay, R. W., Zhang, J., Schweiger, A. J., and Steele, M. A.: Seasonal predictions of ice extent in the Arctic Ocean, *J. Geophys. Res.*, 113, C02 023, <https://doi.org/10.1029/2007JC004259>, <http://onlinelibrary.wiley.com/doi/10.1029/2007JC004259/abstract>, 2008.
- 530 Madec, G.: NEMO ocean engine, Tech. rep., Institut Pierre-Simon Laplace (IPSL), <http://www.nemo-ocean.eu/About-NEMO/Reference-manuals>, 2008.
- Mayer, M., Tietsche, S., Haimberger, L., Tsubouchi, T., Mayer, J., and Zuo, H.: An improved estimate of the coupled Arctic energy budget, *Journal of Climate*, 32, 7915–7934, 2019.



- 535 Meier, W. N., Hovelsrud, G. K., Van Oort, B. E., Key, J. R., Kovacs, K. M., Michel, C., Haas, C., Granskog, M. A., Gerland, S., Perovich, D. K., et al.: Arctic sea ice in transformation: A review of recent observed changes and impacts on biology and human activity, *Reviews of Geophysics*, 52, 185–217, 2014.
- Mogensen, K., Balmaseda, M. A., and Weaver, A.: The NEMOVAR ocean data assimilation system as implemented in the ECMWF ocean analysis for System 4, Tech. Rep. 668, European Centre for Medium-Range Weather Forecasts, 2012.
- 540 Mori, M., Watanabe, M., Shiogama, H., Inoue, J., and Kimoto, M.: Robust Arctic sea-ice influence on the frequent Eurasian cold winters in past decades, *Nature Geoscience*, 7, 869–873, <https://doi.org/10.1038/ngeo2277>, <http://dx.doi.org/10.1038/ngeo2277>, 2014.
- Mu, L., Yang, Q., Losch, M., Losa, S. N., Ricker, R., Nerger, L., and Liang, X.: Improving sea ice thickness estimates by assimilating CryoSat-2 and SMOS sea ice thickness data simultaneously, *Quarterly Journal of the Royal Meteorological Society*, 144, 529–538, <https://doi.org/10.1002/qj.3225>, <http://doi.wiley.com/10.1002/qj.3225>, 2018.
- 545 Overland, J. E., Dethloff, K., Francis, J. A., Hall, R. J., Hanna, E., Kim, S.-J., Screen, J. A., Shepherd, T. G., and Vihma, T.: Nonlinear response of mid-latitude weather to the changing Arctic, *Nature Climate Change*, 6, 992, 2016.
- Ricker, R., Hendricks, S., Kaleschke, L., Tian-Kunze, X., King, J., and Haas, C.: A weekly Arctic sea-ice thickness data record from merged CryoSat-2 and SMOS satellite data, *The Cryosphere*, 11, 1607–1623, <https://doi.org/10.5194/tc-11-1607-2017>, <https://www.the-cryosphere.net/11/1607/2017/>, 2017.
- 550 Ruggieri, P., Buizza, R., and Visconti, G.: On the link between Barents-Kara sea ice variability and European blocking, *Journal of Geophysical Research: Atmospheres*, 121, 5664–5679, 2016.
- Sallila, H., Farrell, S. L., McCurry, J., and Rinne, E.: Assessment of contemporary satellite sea ice thickness products for Arctic sea ice, *The Cryosphere*, 13, 1187–1213, 2019.
- Scarlat, R. C., Spreen, G., Heygster, G., Huntemann, M., Pañilea, C., Toudal Pedersen, L., and Saldo, R.: Sea Ice and Atmospheric Parameter Retrieval From Satellite Microwave Radiometers: Synergy of AMSR2 and SMOS Compared With the CIMR Candidate Mission, *Journal of Geophysical Research: Oceans*, p. e2019JC015749, 2020.
- 555 Semtner, A. J.: A Model for the Thermodynamic Growth of Sea Ice in Numerical Investigations of Climate, *J. Phys. Oceanogr.*, 6, 379–389, 1976.
- Sigmond, M., Fyfe, J. C., Flato, G. M., Kharin, V. V., and Merryfield, W. J.: Seasonal forecast skill of Arctic sea ice area in a dynamical forecast system, *Geophysical Research Letters*, 40, 529–534, <https://doi.org/10.1002/grl.50129>, <http://onlinelibrary.wiley.com/doi/10.1002/grl.50129/abstract>, 2013.
- 560 Stockdale, T., Alonso-Balmaseda, M., Johnson, S., Ferranti, L., Molteni, F., Magnusson, L., Tietsche, S., Vitart, F., Decremet, D., Weisheimer, A., Roberts, C. D., Balsamo, G., Keeley, S., Mogensen, K., Zuo, H., Mayer, M., and Monge-Sanz, B.: SEAS5 and the future evolution of the long-range forecast system, ECMWF Technical Memorandum, <https://doi.org/10.21957/z3e92di7y>, <https://www.ecmwf.int/node/18750>, 2018.
- 565 Tang, Y. M., Balmaseda, M. A., Mogensen, K. S., Keeley, S. P. E., and Janssen, P. E. A. M.: Sensitivity of sea ice thickness to observational constraints on sea ice concentration, Tech. Rep. 707, European Centre for Medium-Range Weather Forecasts, Reading, 2013.
- Tian-Kunze, X., Kaleschke, L., Maaß, N., Mäkynen, M., Serra, N., Drusch, M., and Krumpfen, T.: SMOS-derived thin sea ice thickness: algorithm baseline, product specifications and initial verification, *The Cryosphere*, 8, 997–1018, <https://doi.org/10.5194/tc-8-997-2014>, <http://www.the-cryosphere.net/8/997/2014/http://www.the-cryosphere.net/8/997/2014/tc-8-997-2014.html>, 2014.
- 570



- Tietsche, S., Notz, D., Jungclaus, J. H., and Marotzke, J.: Assimilation of sea-ice concentration in a global climate model – physical and statistical aspects, *Ocean Science*, 9, 19–36, <https://doi.org/10.5194/os-9-19-2013>, <http://www.ocean-sci.net/9/19/2013/os-9-19-2013.html>, 2013.
- 575 Tietsche, S., Day, J. J., Guemas, V., Hurlin, W. J., Keeley, S. P. E., Matei, D., Msadek, R., Collins, M., and Hawkins, E.: Seasonal to interannual Arctic sea-ice predictability in current global climate models, *Geophysical Research Letters*, 41, 1035–1043, <https://doi.org/10.1002/2013GL058755>, <http://onlinelibrary.wiley.com/doi/10.1002/2013GL058755/abstract>, 2014.
- Tietsche, S., Balmaseda, M. a., Zuo, H., and Mogensen, K.: Arctic sea ice in the global eddy-permitting ocean reanalysis ORAP5, *Climate Dynamics*, <https://doi.org/10.1007/s00382-015-2673-3>, <http://link.springer.com/10.1007/s00382-015-2673-3>, 2015.
- 580 Tietsche, S., Balmaseda, M., Rosnay, P., Zuo, H., Tian-Kunze, X., and Kaleschke, L.: Thin Arctic sea ice in L-band observations and an ocean reanalysis, *The Cryosphere*, 12, 2051–2072, <https://doi.org/10.5194/tc-12-2051-2018>, <https://www.the-cryosphere.net/12/2051/2018/>, 2018.
- Tonboe, R., Lavelle, J., Pfeiffer, R., and Howe, E.: Product User Manual for OSI SAF Global Sea Ice Concentration (Product OSI-401-b), 2017.
- 585 Uotila, P., Goosse, H., Haines, K., Chevallier, M., Barthelemy, A., Bricaud, C., Carton, J., Fuckar, N., Garric, G., Iovino, D., Kauker, F., Korhonen, M., Lien, V. S., Marnela, M., Massonnet, F., Mignac, D., Peterson, K. A., Sadikni, R., Shi, L., Tietsche, S., Toyoda, T., Xie, J., and Zhang, Z.: An assessment of ten ocean reanalyses in the polar regions, *Climate Dynamics*, (in press), 1–64, 2018.
- Xie, J., Counillon, F., and Bertino, L.: Impact of assimilating a merged sea-ice thickness from CryoSat-2 and SMOS in the Arctic reanalysis, *The Cryosphere*, 12, 3671–3691, 2018.
- 590 Zampieri, L., Goessling, H. F., and Jung, T.: Bright prospects for Arctic sea ice prediction on subseasonal time scales, *Geophysical Research Letters*, 45, 9731–9738, 2018.
- Zuo, H., Balmaseda, M. A., and Mogensen, K.: The new eddy-permitting ORAP5 ocean reanalysis: description, evaluation and uncertainties in climate signals, *Climate Dynamics*, 49, 791–811, <https://doi.org/10.1007/s00382-015-2675-1>, <http://link.springer.com/article/10.1007/s00382-015-2675-1>, 2017.
- 595 Zuo, H., Balmaseda, M. A., Tietsche, S., Mogensen, K., and Mayer, M.: The ECMWF operational ensemble reanalysis-analysis system for ocean and sea-ice: a description of the system and assessment, *Ocean Science*, 2019, 779–808, <https://doi.org/10.5194/os-15-779-2019>, <https://doi.org/10.5194/os-15-779-2019/>, 2019.